



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

8430.16

Vaporisateur contre les mites et
les poux pour tout type d'oiseaux

Aerosol contra ácaros y piojos
para todo tipo de pájaros

UltraCare™

Mite & Lice Bird Spray

For All Birds

For Use on Caged Birds
to Kill Mites and Lice

ACTIVE INGREDIENTS:

Pyrethrins 0.03%
Piperonyl Butoxide, "Technical" . . . 0.30%

OTHER INGREDIENTS: 99.67%

TOTAL: 100.00%

EPA Reg. No. 87703-2

EPA Est. No. 87703-VA-01

*Equivalent to 0.240%

(butylcarbityl)(6-propylpiperonyl)
ether and 0.06% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional
precautionary statements

8 in 1

NET 8 FL. OZ. (237 mL)

UltraCare™ Mite & Lice Bird Spray

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Avoid inhalation of vapors. Avoid contact with skin. Wash contaminated skin with soap and water. Avoid spraying food and foodstuffs. Cover fish aquariums before spraying.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product inconsistent with its labeling. For control of bird lice on parakeets, canaries and other caged birds, leave birds in cage. Remove food and foodstuffs. Spray directly on birds from a distance of 12 to 18 inches. Avoid the bird's eyes and face, spray lightly utilizing 3 or 4 bursts. For cage mite control, remove bird(s), food and water cups, etc., and spray cage thoroughly by holding unit at least one foot from surface utilizing 5 or 6 bursts. To control mites and bird lice, may repeat treatment on cage or bird no more than two times a week. Bird cages should be cleaned in hot water as part of weekly regimen.

STORAGE AND DISPOSAL

Pesticide Storage: Store in cool dry place away from heat, open flame, food and water supplies that is inaccessible to children and pets.

Pesticide Disposal and Container Handling: Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

FIRST AID

If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If in eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If inhaled

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

HOTLINE

Have the product container or label with you when calling
poison control center or doctor or going for treatment. For
emergency information, call (800) 645-5154.

8 in 1 Made in U.S.A. United Pet Group, Inc.
3001 Commerce St. Blacksburg, Va 24060-6671
(800) 645-5154 www.eightinone.com

ITEM# C338 / 041-0340-06

® Registered Trademark and TM Trademark of United
Pet Group, Inc. (UPG) © 2014 UPG

