

18.5 inches Width



LICENSED

LICENSED

PERIOD 2011-2013 LIC. NO.

Storage and Disposal

PERIOD 2014-2016 LIC. NO. 8447.2

Cat. No. S5N

Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Store in original container in a cool, well-ventilated, preferably locked area inaccessible to children and pets. Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site, or at an approved waste disposal facility. Container Disposal: Non-refillable container; do not reuse or refill this container. To avoid wastes, use all material in this container by application according to label direction. When empty, fill the container about 10 percent full of water. Agitate vigorously and dispose of contents on site or at an approved waste disposal facility. Offer the container for recycling, if available; otherwise dispose of container in trash.

First Aid

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. If on Skin or Clothing: Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-535-5053 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapors and contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash clothing before reuse.

Physical or Chemical Hazards: Do not use or store near heat or open flame.

Weiman Products, LLC
755 Tri-State Parkway
Gurnee, IL 60031
(800) 837-8140



(01) 0 0041598 00009 6



Germicidal Solution

Cleaner, Disinfectant & Deodorizer
Bactericidal, Tuberculocidal, Fungicidal and Virucidal

Active Ingredients:

Table with 2 columns: Ingredient name and percentage. Includes n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18), dimethyl benzyl ammonium chloride (0.154%), n-Alkyl (68% C12, 32% C14), dimethyl ethylbenzyl ammonium chloride (0.154%), Isopropanol (21.000%), Other Ingredients (78.692%), and Total (100.000%).

Keep Out Of Reach Of Children

CAUTION

See side panel for First Aid and additional Precautionary Statements.

EPA Reg. No. 1130-15

EPA Est. 1130-IL-1

Net Contents: 1 Gallon (3.78 Liters)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Weiman® Germicidal Solution is a general purpose cleaner, deodorizer and disinfectant to be used where control of cross-contamination hazards is of prime importance. Weiman® Germicidal Solution provides a way to effectively clean and disinfect hard, non-porous environmental surfaces. Weiman® Germicidal Solution aids in the reduction of cross-contamination hazards. Easy to use; no mixing or diluting except for ultrasonic cleaning. Ready-to-use.

Use Weiman® Germicidal Solution on all hard, non-porous environmental surfaces/articles including: Chrome, Formica® Brand, glass, painted surfaces, plastics (such as polycarbonate, polyvinylchloride, polypropylene and polystyrene), Plexiglas®, glazed porcelain and stainless steel.

Use Weiman® Germicidal Solution to clean, deodorize and disinfect such items as: Ambulance equipment/surfaces, counters/countertops, CPR training mannequins, dental chairs/countertops/instruments/implements/OR surfaces, exercise equipment/benches/mats, hardhats, headphones, laboratory equipment/surfaces, medical instruments/implements, nail care implements, operating room lights/tables, respiratory therapy equipment, respirators/face pieces, tanning beds, telephones and veterinary care surfaces/instruments/implements.

This product has demonstrated effectiveness against Influenza Strain A2/HK virus\* and Avian Influenza A Virus\* and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called "swine flu").

Cleaning, Precleaning and Deodorizing: Prior to disinfection, preclean all surfaces with Weiman® Germicidal Solution, making sure to remove all gross filth and heavy soil. Apply solution to surface and allow surface to remain wet for 30 seconds prior to wiping with a paper towel or clean towel/sponge.

Ultrasonic Cleaning: Thoroughly pre-rinse instruments under running water to remove visible gross debris. Using 1 ounce of Weiman® Germicidal Solution per liter of water in ultrasonic unit, immerse instruments into solution and activate ultrasonic unit for 5 minutes (or longer if necessary). Remove and rinse instruments. Follow with appropriate cleaning and disinfection process. Change solution as needed when it becomes diluted or visibly soiled. To Disinfect Hard, Nonporous Surfaces/Articles: To Disinfect Acinetobacter baumannii, Aspergillus niger, Avian Influenza A Virus, Candida albicans, Canine Parvovirus, Escherichia coli O157:H7, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex Virus Type 2, HIV-1 (AIDS Virus), Influenza Strain A2/HK Virus, Mycobacterium bovis BCG, Pandemic 2009

H1N1 Influenza A Virus\*, Polio I Virus, Pseudomonas aeruginosa, Rhinovirus, Rotavirus Strain WA, Salmonella enterica, Staphylococcus aureus, Methicillin-resistant Staphylococcus aureus (MRSA), Trichophyton mentagrophytes and Vancomycin-resistant Enterococcus faecalis (VRE): Using Weiman® Germicidal Solution as is (do not dilute), apply solution to thoroughly wet the precleaned surfaces. Allow to remain wet for 6 minutes prior to wiping or allow surfaces to air dry. For disinfection of Avian Influenza A, use this product only for surfaces that are not conducive to treatment by immersion or excess liquid. Note: Disinfected food contact surfaces must be rinsed with potable water after cleaning and/or disinfection and before reuse. After disinfection, critical and semi-critical devices must undergo appropriate terminal sterilization/high-level disinfection.

Special Instructions for Cleaning and Decontamination Against HBV, HCV and HIV-1 (AIDS Virus) for Surfaces or Objects Soiled with Blood or Body Fluids

Kills HBV, HCV and HIV-1 (AIDS Virus) on precleaned environmental surfaces or objects previously soiled with blood or body fluids at room temperature (20-25°C) in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HBV, HCV and the human immunodeficiency virus Type 1 (HIV-1) associated with AIDS. Personal Protection: When using the germicidal solution, wear disposable latex gloves, protective gown, face mask and eye coverings, as appropriate, when handling HBV, HCV and HIV-1 infected blood or body fluids. Cleaning Procedure: All blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the germicidal solution. Contact Time: Thoroughly wet surface. Allow surface to remain wet for 6 minutes prior to wiping or allow surface to air dry. Disposal of Infectious Materials: Dispose of used solution in accordance with local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

5 inches Height

0.25

0.25

0.25

10.625 inches Width

0.25

0.25



LICENSED

8447.2

PERIOD 2014-2016 LIC. NO.

Cat. No. S5NQ

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, well-ventilated, preferably locked area inaccessible to children and pets. Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site, or at an approved waste disposal facility. Container Disposal: Non-refillable container; do not reuse or refill this container. To avoid wastes, use all material in this container by application according to label directions. When empty, fill the container about 10 percent full of water. Agitate vigorously and dispose of contents on site or at an approved waste disposal facility. Offer the container for recycling, if available; otherwise dispose of container in trash.

First Aid

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. If on Skin or Clothing: Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-535-5053 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapors and contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash clothing before reuse.

Physical or Chemical Hazards: Do not use or store near heat or open flame.



(01) 0 0041596 00010 2 Weiman Products, LLC 755 Tri-State Parkway Gurnee, IL 60031 (800) 837-8140



Department of Agriculture STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.



Germicidal Solution

Cleaner, Disinfectant & Deodorizer Bactericidal, Tuberculocidal, Fungicidal and Virucidal

Active Ingredients:

- n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chloride..... 0.154% n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chloride.. 0.154% Isopropanol ..... 21.000%

- Other Ingredients ..... 78.692% Total ..... 100.000%

Keep Out Of Reach Of Children

CAUTION

See side panel for First Aid and additional Precautionary Statements.

EPA Reg. No. 1130-15 EPA Est. 1130-IL-1

Net Contents: 1 Quart (0.95 L)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Weiman® Germicidal Solution is a general purpose cleaner, deodorizer and disinfectant to be used where control of cross-contamination hazards is of prime importance. Weiman® Germicidal Solution provides a way to effectively clean and disinfect hard, non-porous environmental surfaces. Weiman® Germicidal Solution aids in the reduction of cross-contamination hazards. Easy to use, no mixing or diluting except for ultrasonic cleaning. Ready-to-use.

Use Weiman® Germicidal Solution on all hard, non-porous environmental surfaces/articles including: Chrome, Formica® Brand, glass, painted surfaces, plastics (such as polycarbonate, polyvinylchloride, polypropylene and polystyrene), Plexiglas®, glazed porcelain and stainless steel.

Use Weiman® Germicidal Solution to clean, deodorize and disinfect such items as: Ambulance equipment/surfaces, counters/countertops, CPR training mannequins, dental chairs/countertops/instruments/implements/OR surfaces, exercise equipment/benches/mats, hardhats, headphones, laboratory equipment/surfaces, medical instruments/implements, nail care implements, operating room lights/tables, respiratory therapy equipment, respirators/facepieces, tanning beds, telephones and veterinary care surfaces/instruments/implements.

This product has demonstrated effectiveness against Influenza Strain A2/HK virus\* and Avian Influenza A Virus\* and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called "swine flu").

Cleaning, Precleaning and Deodorizing: Prior to disinfection, preclean all surfaces with Weiman® Germicidal Solution, making sure to remove all gross filth and heavy soil. Apply solution to surface and allow surface to remain wet for 30 seconds prior to wiping with a paper towel or clean towel/sponge. Ultrasonic Cleaning: Thoroughly pre-rinse instruments under running water to remove visible gross debris. Using 1 ounce of Weiman® Germicidal Solution per liter of water in ultrasonic unit, immerse instruments into solution and activate ultrasonic unit for 5 minutes (or longer if necessary). Remove and rinse instruments. Follow with appropriate cleaning and disinfection process. Change solution as needed when it becomes diluted or visibly soiled. To Disinfect Hard, Nonporous Surfaces/Articles: To Disinfect Acinetobacter baumannii, Aspergillus niger, Avian Influenza A Virus, Candida albicans, Canine Parvovirus, Escherichia coli O157:H7, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex Virus Type

2, HIV-1 (AIDS Virus), Influenza Strain A2/HK Virus, Mycobacterium bovis BCG, Pandemic 2009 H1N1 Influenza A Virus\*, Polio I Virus, Pseudomonas aeruginosa, Rhinovirus, Rotavirus Strain WA, Salmonella enterica, Staphylococcus aureus, Methicillin-resistant Staphylococcus aureus (MRSA), Trichophyton mentagrophytes and Vancomycin-resistant Enterococcus faecalis (VRE): Using Weiman® Germicidal Solution as is (do not dilute), apply solution to thoroughly wet the precleaned surfaces. Allow to remain wet for 6 minutes prior to wiping or allow surfaces to air dry. For disinfection of Avian Influenza A, use this product only for surfaces that are not conducive to treatment by immersion or excess liquid. Note: Disinfected food contact surfaces must be rinsed with potable water after cleaning and/or disinfection and before reuse. After disinfection, critical and semi-critical devices must undergo appropriate terminal sterilization/high-level disinfection.

Special Instructions for Cleaning and Decontamination Against HBV, HCV and HIV-1 (AIDS Virus) for Surfaces or Objects Soiled with Blood or Body Fluids

Kills HBV, HCV and HIV-1 (AIDS Virus) on precleaned environmental surfaces or objects previously soiled with blood or body fluids at room temperature (20-25°C) in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HBV, HCV and the human immunodeficiency virus Type 1 (HIV-1) associated with AIDS. Personal Protection: When using the germicidal solution, wear disposable latex gloves, protective gown, face mask and eye coverings, as appropriate, when handling HBV, HCV and HIV-1 infected blood or body fluids. Cleaning Procedure: All blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the germicidal solution. Contact Time: Thoroughly wet surface. Allow surface to remain wet for 6 minutes prior to wiping or allow surface to air dry. Disposal of Infectious Materials: Dispose of used solution in accordance with local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

4.625 inches Height

0.25

0.25

8 inches Width

0.25

0.25

Department of Agriculture  
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

8447.2

# WEIMAN®

## Germicidal Solution

### Cleaner, Disinfectant & Deodorizer

### Bactericidal, Tuberculocidal, Fungicidal and Virucidal

#### Active Ingredients:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)	
dimethyl benzyl ammonium chloride.....	0.154%
n-Alkyl (68% C12, 32% C14)	
dimethyl ethylbenzyl ammonium chloride.....	0.154%
Isopropanol .....	21.000%
<b>Other Ingredients</b> .....	<b>78.692%</b>
<b>Total</b> .....	<b>100.000%</b>

## Keep Out Of Reach Of Children CAUTION

See side panel for First Aid and additional Precautionary Statements.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Weiman® Germicidal Solution is a general purpose cleaner, deodorizer and disinfectant to be used where control of cross-contamination hazards is of prime importance. Weiman® Germicidal Solution provides a way to effectively clean and disinfect hard, non-porous environmental surfaces. Weiman® Germicidal Solution aids in the reduction of cross-contamination hazards. Easy to use, no mixing or diluting except for ultrasonic cleaning. Ready-to-use.

Use Weiman® Germicidal Solution on all hard, non-porous environmental surfaces/articles including: Chrome, Formica® Brand, glass, painted surfaces, plastics (such as polycarbonate, polyvinylchloride, polypropylene and polystyrene), Plexiglas®, glazed porcelain and stainless steel.

Use Weiman® Germicidal Solution to clean, deodorize and disinfect such items as: Ambulance equipment/surfaces, counters/countertops, CPR training mannequins, dental chairs/countertops/instruments/implents/OR surfaces, exercise equipment/benches/mats, hardhats, headphones, laboratory equipment/surfaces, medical instruments/implents, nail care implements, operating room lights/tables, respiratory therapy equipment, respirators/face pieces, tanning beds, telephones and veterinary care surfaces/instruments/implents.

This product has demonstrated effectiveness against Influenza Strain A2/HK virus® and Avian Influenza A Virus® and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called "swine flu").

EPA Reg. No. 1130-15

EPA Est. 1130-IL-1

**Net Contents: 55 Gallons (208 Liters)**

Cat. No. S5N55

**DIRECTIONS FOR USE (Continued): Cleaning, Precleaning and Deodorizing:** Prior to disinfection, preclean all surfaces with Weiman® Germicidal Solution, making sure to remove all gross filth and heavy soil. Apply solution to surface and allow surface to remain wet for 30 seconds prior to wiping with a paper towel or clean towel/sponge. **Ultrasonic Cleaning:** Thoroughly pre-rinse instruments under running water to remove visible gross debris. Using 1 ounce of Weiman® Germicidal Solution per liter of water in ultrasonic unit, immerse instruments into solution and activate ultrasonic unit for 5 minutes (or longer if necessary). Remove and rinse instruments. Follow with appropriate cleaning and disinfection process. Change solution as needed when it becomes dulled or visibly soiled. **To Disinfect Hard, Nonporous Surfaces/Articles:** To Disinfect: *Acinetobacter baumannii*, *Aspergillus niger*, Avian Influenza A Virus, *Candida albicans*, Canine Parvovirus, *Escherichia coli* O157:H7, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex Type 2, HIV-1 (AIDS Virus), Influenza Strain A2/HK Virus, *Mycobacterium bovis* BCG, Pandemic 2009 H1N1 Influenza A Virus®, Polio I Virus, *Pseudomonas aeruginosa*, Rhinovirus, Rotavirus Strain WA, *Salmonella enterica*, Methicillin-resistant *Staphylococcus aureus* (MRSA), *Staphylococcus aureus*, *Trichophyton mentagrophytes* and Vancomycin-resistant *Enterococcus faecalis* (VRE): Using Weiman® Germicidal Solution as is (do not dilute), apply solution to thoroughly wet the precleaned surfaces. Allow to remain wet for 6 minutes prior to wiping or allow surfaces to air dry. For disinfection of Avian Influenza A, use this product only for surfaces that are not conducive to treatment by immersion or excess liquid. **Note:** Disinfected food contact surfaces must be rinsed with potable water after cleaning and/or disinfection and before reuse. After disinfection, critical and semi-critical devices must undergo appropriate terminal sterilization/high-level disinfection.

#### Special Instructions for Cleaning and Decontamination Against HBV, HCV and HIV-1 (AIDS Virus) for Surfaces or Objects Soiled with Blood or Body Fluids

Kills HBV, HCV and HIV-1 (AIDS Virus) on precleaned environmental surfaces or objects previously soiled with blood or body fluids at room temperature (20-25 °C) in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HBV, HCV and the human immunodeficiency virus Type 1 (HIV-1) associated with AIDS. **Personal Protection:** When using the germicidal solution, wear disposable latex gloves, protective gown, face mask and eye coverings, as appropriate, when handling HBV, HCV and HIV-1 infected blood or body fluids. **Cleaning Procedure:** All blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the germicidal solution. **Contact Time:** Thoroughly wet surface. Allow surface to remain wet for 6 minutes prior to wiping or allow surface to air dry. **Disposal of Infectious Materials:** Dispose of used solution in accordance with local regulations for infectious waste disposal.

*This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.*

## Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in original container in a cool, well-ventilated, preferably locked area inaccessible to children and persons unfamiliar with its use. **Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site, or at an approved waste disposal facility. **Container Disposal:** Non-refillable container, do not reuse or refill this container. Offer for recycling, if available. Triple rinse (or equivalent) container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinse into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

#### First Aid

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. **If on Skin or Clothing:** Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. **If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-535-5053 for emergency medical treatment information.


#### Precautionary Statements

##### Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapors and contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash clothing before reuse.

**Physical or Chemical Hazards:** Do not use or store near heat or open flame.

Weiman Products, LLC  
755 Tri-State Parkway  
Gurnee, IL 60031  
(800) 837-8140



(01) 0 0041598 00049 2

5 inches Height

Department of Agriculture  
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

0.25