

LICENSED

PERIOD 2015-2017 LIC. NO.

8456.5

Use GymWipes® / CareWipes® Antibacterial Force Wipes to clean and disinfect hard non-porous surfaces of gym equipment, exercise machines, wrestling mats, examining tables. For use in hospital and healthcare settings, schools, health clubs, athletic facilities and daycare centers. Rinse all equipment that comes in contact with skin before reuse with clean warm water, and allow to air dry.

TO OPEN PACKAGE: Remove cover. Pull up wipe corner from center of roll. Thread through hole in the cover. Replace cover. Open center cap on cover to remove wipe. When not in use, close center cap to prevent moisture loss.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO CLEAN AND DEODORIZE: Wipe surface with wipe. Let air dry. Discard used wipe in trash. Do not flush in toilet.

TO CLEAN AND DISINFECT (DISINFECTION / VIRUCIDAL* / FUNGICIDAL / MOLD AND MILDEW CONTROL DIRECTIONS):

Wipe surface with wipe until surface is visibly wet. Allow surface to remain wet for 10 minutes. Let air dry. If surfaces are extremely dirty, clean first with another wipe before disinfecting. Discard used wipe in trash. Do not flush. Rinse food contact surfaces with clean, potable water after disinfecting. Do not use to disinfect dishes, glassware or utensils. Do not use as a diaper wipe or for personal cleansing. (Note: Do not use on unpainted wood, natural marble or brass.)

For fungicidal activity, GymWipes® / CareWipes® Antibacterial Force Wipes is an effective fungicide against Tricophyton mentagrophytes (the athlete's foot fungus) in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. GymWipes® / CareWipes® Antibacterial Force Wipes is a one-step fungicide.

For mold and mildew, GymWipes® / CareWipes® Antibacterial Force Wipes will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more if new growth appears.

***KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

2XL-400 Buckets / 2XL-401 Refill

EPA REG. NO. 6836-313-82144
EPA EST. NO. 50757-WI-1



**A Fast and Effective
Antibacterial Formula
That Cleans, Disinfects
and Deodorizes**

- Virucidal
- Bactericidal
- Bleach Free
- Fungicidal
- Ammonia-Free



800

**LARGE 8" x 6"
TOWELETTES**

NET CONTENTS
10 lbs.

ACTIVE INGREDIENTS:

Octyl decyl dimethyl ammonium chloride	0.069%
Dioctyl dimethyl ammonium chloride	0.028%
Didecyl dimethyl ammonium chloride	0.042%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%)	
dimethyl benzyl ammonium chloride	0.093%

OTHER INGREDIENTS:	99.768%
TOTAL	100.000%

*Weight of wipes plus liquid.

KEEP OUT OF REACH OF CHILDREN
CAUTION

LICENSED

PERIOD 2012-2014 LIC. NO.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS.

Personal protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning procedure: Blood and other body fluids containing HIV or HBV or HCV must thoroughly be cleaned from surfaces and objects before application of GymWipes® / CareWipes® Antibacterial Force Wipes. GymWipes® / CareWipes® Antibacterial Force Wipes can be used for this purpose.

Contact time: Leave surface wet for 1 minute (60 seconds) for HIV-1 and 10 minutes for HBV and HCV. Use a 10-minute contact time for disinfection against all other viruses, bacteria and fungi claimed.

Disposal of infectious material: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

*This product is for use on hard non-porous surfaces not conducive to treatment by immersion or excess liquid against Avian Influenza A.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

STORAGE AND DISPOSAL:

Store product in original container in areas inaccessible to children. Keep securely closed. Refill empty container only with GymWipes® / CareWipes® Antibacterial Force Wipes. Discard empty container in trash or recycle.



Sold by: 2XL Corporation

2415 Braga Drive • Broadview, IL 60155 • (888) 977-3726

www.2xllcorp.com



LICENSED

PERIOD 2009-2011 LIC. NO.

APPROVED

By Andrea Chutjian at 9:24 am, Jun 02, 2009

Per EPA label 2/27/08 & 8/1/08 Notification.
LONZA / REGULATORY

GymWipes® / CareWipes® Antibacterial Force Wipes are effective against:

****[Bacteria]

Community Associated Methicillin-Resistant Staphylococcus aureus [CA-MRSA] NRS384, USA300

Community Associated Methicillin-Resistant Staphylococcus aureus [CA-MRSA] NRS123, USA400

Staphylococcus aureus (Staph)

Salmonella enterica (Salmonella)

Pseudomonas aeruginosa [Pseudomonas]

Campylobacter jejuni [Campylobacter]

Acinetobacter baumannii

Brevibacterium ammoniagenes [Brevibacterium]

Campylobacter jejuni

Enterobacter aerogenes

Enterococcus faecalis [Enterococcus]

Enterococcus faecalis- Vancomycin Resistant [VRE]

Escherichia coli [E. coli]

ESBL Escherichia coli – [Extended spectrum beta-lactamase producing E. coli]

Klebsiella pneumoniae [Klebsiella]

Legionella pneumophila

Pseudomonas cepacia [Pseudomonas]

Salmonella schottmuelleri [Salmonella]

Salmonella typhi [Salmonella]

Serratia marcescens [Serratia]

Shigella dysenteriae [Shigella]

Staphylococcus aureus – Methicillin-Resistant [MRSA]

Staphylococcus aureus – Multi-Drug Resistant

Staphylococcus aureus -- Vancomycin Intermediate Resistant [VISA]

Streptococcus pyogenes [Strep] [a cause of scarlet fever]

Vibrio cholerae

[Viruses:]

* Rotavirus (Diarrhea-causing virus)

* Hepatitis B Virus [HBV]

* Hepatitis C Virus [HCV]

* Herpes Simplex Virus Type 1

* Herpes Simplex Virus Type 2

* HIV-1 [AIDS VIRUS]

* Human Coronavirus

* Influenza Type A/Brazil [Influenza]

* Norwalk Virus [Norovirus]

* Respiratory Syncytial Virus [RSV]

* SARS Associated Coronavirus [SARS]

[cause of Severe Acute Respiratory Syndrome]

* Vaccinia [Pox Virus]

[Animal Viruses:]

Avian Influenza [H5N1]

Avian Influenza [H3N2]

Avian Infectious Bronchitis Virus

Canine Distemper Virus

Feline Calicivirus

Newcastle's Disease Virus

Pseudorabies Virus

[Fungi:]

Aspergillus niger

Trichophyton metagrophytes [the Athlete's

Foot Fungus]

[A cause of Ringworm of the foot]



www.2xlcorp.com

