

For Barn And Farm Use

This product is a one-step disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, automatic feeders, buckets, brushes, exterior surfaces of watering appliances, farm vehicles (hard, non-porous surface of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs or any hard non-porous washable (non-food contact) surfaces where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Spray 6-8 inches from surface.
3. Allow surface to remain wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

Animal Housing Facilities

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, stables, etc.

1. Remove all animals and feed from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty (or cover) all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Saturate all surfaces by spraying solution until wet. To disinfect, all surfaces must remain wet for 1 minute.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and water appliances, spray with solution and let stand for 1 minute. Then thoroughly scrub all treated surfaces with soap or detergent, and rinse with potable water before reuse.

For Use as a Cleaner and/ or Deodorizer

Spray 6-8 inches from hard, non-porous surfaces. Allow to air dry.

VAH026 (16/034)

8477.10

INTERvention®

Farm Animal Care Disinfectant Cleaner & Deodorizer

Ready To Use

ACTIVE INGREDIENT

Hydrogen Peroxide0.5%

OTHER INGREDIENTS99.5%

Total100.0%

KEEP OUT OF REACH OF CHILDREN

EPA Reg. No. 74559-9
EPA Est. No. 74559-CAN-1

Part No. 24101

See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use

FOR BARN AND
FARM USE

For Barn And Farm Use

Claims

See reference sheet for a complete list of organisms.

Bactericidal in 1 Minute:

Staphylococcus aureus

Staphylococcus aureus (Methicillin Resistant (MRSA))

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Klebsiella pneumoniae

Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))

Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))

Klebsiella pneumoniae (Carbapenem-Resistant (CRE))

Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))

Staphylococcus aureus (Vancomycin Resistant (VRSA))

Staphylococcus epidermidis (Methicillin Resistant (MRSE))

Enterococcus faecalis (Vancomycin Resistant (VRE))

Enterococcus faecium

Bordetella bronchiseptica

Streptococcus pneumoniae (Penicillin Resistant (PRSP))

Pasteurella multocida

Serratia marcescens

Shigella dysenteriae

*Virucidal in 30 Seconds, unless otherwise noted:

Polio Type 1 (Strain Brunhilde)

Avian Influenza A Virus (H7N9)

Avian Influenza A Virus (H3N2)

Animal premise *Virucidal Activity in 1 Minute, unless otherwise noted:

Bovine Viral Diarrhea Virus – 30 seconds

Duck Hepatitis B virus – 30 seconds

Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

Fragrance
Free Formula



US PATENT:
7,632,523
& 7,354,604

Net Contents:

1 U.S. Quart
(946 ml)

Animal Premise *Virucidal 1 Minute

*Virucidal
30 Seconds

Bactericidal
1 Minute

Fungicidal
30 Seconds, unless
otherwise noted

Broad-Spectrum
Non-Food Contact Sanitizer
10 Seconds

READ S.D.S. BEFORE
USING PRODUCT

SDS available online
at www.virox.com



Manufactured by
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com

This product has demonstrated effectiveness against Influenza Type A_v (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Fungicidal in 30 Seconds, unless otherwise noted:

Trichophyton mentagrophytes

Microsporum canis – 1 minute

Tuberculocidal in 1 Minute: *Mycobacterium bovis* (BCG)

Broad-Spectrum Non-Food Contact Sanitizer in 10 seconds against:

Klebsiella pneumoniae

Staphylococcus aureus

Enterobacter aerogenes

Acinetobacter baumannii

Listeria monocytogenes

Campylobacter jejuni

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Pseudomonas aeruginosa

Escherichia coli O157:H7

Vancomycin Resistant *Enterococcus faecalis* (VRE)

Methicillin Resistant *Staphylococcus aureus* (MRSA)

Storage and Disposal

PESTICIDE STORAGE: Store in original container in areas inaccessible to small children. Keep securely closed.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill empty container. Wrap and put in trash or offer for recycling if available.

ENVIRONMENTAL HAZARDS: This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

UPC CODE
0-68460-50313-4

INTERvention®



For Barn And Farm Use

This product is a one-step disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, buckets, brushes, automatic feeders, exterior surfaces of watering appliances, farm vehicles (hard, non-porous surface of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs or any hard non-porous washable (non-food contact) surfaces where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Spray 6-8 inches from surface.
3. Allow surface to remain wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

Animal Housing Facilities

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, stables, etc.

1. Remove all animals and feed from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty (or cover) all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Saturate all surfaces by spraying solution until wet. To disinfect, all surfaces must remain wet for 1 minute.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and water appliances, spray with solution and let stand for 1 minute. Then thoroughly scrub all treated surfaces with soap or detergent, and rinse with potable water before reuse.

For Use as a Cleaner and/or Deodorizer

Spray 6-8 inches from hard, non-porous surfaces. Allow to air dry.

INTERVENTION[®]

INTERVENTION[®]

Farm Animal Care Disinfectant Cleaner & Deodorizer

Ready To Use

ACTIVE INGREDIENT
Hydrogen Peroxide 0.5%
OTHER INGREDIENTS 99.5%
Total 100.0%

EPA Reg. No. 74559-9
EPA Est. No. 74559-CAN-1

Part No. 24105

KEEP OUT OF REACH OF CHILDREN

See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use

FOR BARN AND FARM USE

Fragrance Free Formula

AHP
US PATENT:
7,632,523
& 7,354,604

Net Contents:
1 Gallon (3.78L) **1G**

For Barn And Farm Use

Claims

See reference sheet for a complete list of organisms.

Bactericidal in 1 Minute:

Staphylococcus aureus

Staphylococcus aureus
(Methicillin Resistant (MRSA))

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Klebsiella pneumoniae

Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))

Klebsiella pneumoniae (New Delhi metallo- β -lactamase 1 (NDM-1))

Klebsiella pneumoniae
(Carbapenem-Resistant (CRE))

Staphylococcus aureus
(Vancomycin Intermediate Resistant (VISA))

Staphylococcus aureus
(Vancomycin Resistant (VRSA))

Staphylococcus epidermidis
(Methicillin Resistant (MRSE))

Enterococcus faecalis
(Vancomycin Resistant (VRE))

Enterococcus faecium

Bordetella bronchiseptica

Streptococcus pneumoniae
(Penicillin Resistant (PRSP))

Pasteurella multocida

Serratia marcescens

Shigella dysenteriae

*Virucidal in 30 Seconds, unless otherwise noted:

Polio Type 1 (Strain Brunhilde)

Avian Influenza A Virus (H7N9)

Avian Influenza A Virus (H3N2)

Animal premise *Virucidal Activity in 1 Minute, unless otherwise noted:

Bovine Viral Diarrhea Virus – 30 seconds

Duck Hepatitis B virus – 30 seconds

Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

This product has demonstrated effectiveness against Influenza Type A₂ (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Fungicidal in 30 Seconds, unless otherwise noted:

Trichophyton mentagrophytes

Microsporum canis – 1 minute

Tuberculocidal in 1 Minute:

Mycobacterium bovis (BCG)

Broad-Spectrum Non-Food Contact Sanitizer in 10 seconds against:

Klebsiella pneumoniae

Staphylococcus aureus

Enterobacter aerogenes

Acinetobacter baumannii

Listeria monocytogenes

Campylobacter jejuni

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Pseudomonas aeruginosa

Escherichia coli O157:H7

Vancomycin Resistant *Enterococcus faecalis* (VRE)

Methicillin Resistant *Staphylococcus aureus* (MRSA)

Storage and Disposal

PESTICIDE STORAGE: Store in original container in areas inaccessible to small children. Keep securely closed.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill empty container. Wrap and put in trash or offer for recycling if available.

ENVIRONMENTAL HAZARDS: This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

Animal Premise *Virucidal
1 Minute

*Virucidal
30 Seconds

Bactericidal
1 Minute

Fungicidal
30 Seconds, unless
otherwise noted

Broad-Spectrum
Non-Food Contact Sanitizer
10 Seconds

READ S.D.S. BEFORE
USING PRODUCT

SDS available online
at www.virox.com

VIROX
TECHNOLOGIES INC.

Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com

UPC CODE
0-68460-50393-6

For Barn And Farm Use

This product is a one-step disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, buckets, brushes, automatic feeders, exterior surfaces of watering appliances, farm vehicles (hard non-porous surfaces of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs or any hard non-porous washable (non-food contact) surfaces where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Spray 6-8 inches from surface.
3. Allow surface to remain wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

Animal Housing Facilities

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, stables, etc.

1. Remove all animals and feed from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty (or cover) all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Saturate all surfaces by spraying solution until wet. To disinfect, all surfaces must remain wet for 1 minute.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and water appliances, spray with solution and let stand for 1 minute. Then thoroughly scrub all treated surfaces with soap or detergent, and rinse with potable water before reuse.

For Use as a Cleaner and/ or Deodorizer

Spray 6-8 inches from hard, non-porous surfaces. Allow to air dry.

READ S.D.S. BEFORE USING PRODUCT

SDS available online at www.virox.com

INTERvention®

INTERvention®

Farm Animal Care Disinfectant Cleaner & Deodorizer

Ready To Use

ACTIVE INGREDIENT
Hydrogen Peroxide 0.5%
OTHER INGREDIENTS 99.5%
Total 100.0%

EPA Reg. No. 74559-9
EPA Est. No. 74559-CAN-1

Part No. 24109

KEEP OUT OF REACH OF CHILDREN

See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use

FOR BARN AND FARM USE

Net Contents:
5 Gallon (18.9L)

Fragrance Free Formula

AHP®
US PATENT:
7,632,523
& 7,354,604

5G

For Barn And Farm Use

Claims

See reference sheet for a complete list of organisms.

Bactericidal in 1 Minute:

Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Bordetella bronchiseptica
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Pasteurella multocida
Serratia marcescens
Shigella dysenteriae

*Virucidal in 30 Seconds, unless otherwise noted:

Polio Type 1 (Strain Brunhilde)
Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)

Animal premise *Virucidal Activity in 1 Minute, unless otherwise noted:

Bovine Viral Diarrhea Virus – 30 seconds
Duck Hepatitis B virus – 30 seconds
Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

This product has demonstrated effectiveness against Influenza Type A₂ (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Fungicidal in 30 Seconds, unless otherwise noted:

Trichophyton mentagrophytes
Microsporum canis – 1 minute

Tuberculocidal in 1 Minute:

Mycobacterium bovis (BCG)

Broad-Spectrum Non-Food Sanitizer in 10 seconds against:

Klebsiella pneumoniae
Staphylococcus aureus
Enterobacter aerogenes
Acinetobacter baumannii
Listeria monocytogenes
Campylobacter jejuni
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Pseudomonas aeruginosa
Escherichia coli O157:H7
Vancomycin Resistant *Enterococcus faecalis* (VRE)
Methicillin Resistant *Staphylococcus aureus* (MRSA)

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in securely closed container in areas inaccessible to small children. Store away from direct sunlight and heat.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. For more information on the disposal of unused, unwanted product and the cleanup of spills, contact the EPA Regional Office or the Manufacturer for guidance.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill; or incinerate, or, if allowed by state and local authorities, burn. If burned, stay out of smoke.

ENVIRONMENTAL HAZARDS: This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

Animal Premise *Virucidal 1 Minute

*Virucidal 30 Seconds

Bactericidal 1 Minute

Fungicidal 30 Seconds, unless otherwise noted

Broad-Spectrum Non-Food Contact Sanitizer 10 Seconds

VIROX®
TECHNOLOGIES INC.

Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com

UPC CODE
0-68460-50484-1

For Barn And Farm Use

This product is a one-step disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, automatic feeders, buckets, brushes, exterior surfaces of watering appliances, farm vehicles (hard non-porous surfaces of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs or any hard non-porous washable (non-food contact) surfaces where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Spray 6-8 inches from surface.
3. Allow surface to remain wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

Animal Housing Facilities

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, stables, etc.

1. Remove all animals and feed from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty (or cover) all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Saturate all surfaces by spraying solution until wet. To disinfect, all surfaces must remain wet for 1 minute.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and water appliances, spray with solution and let stand for 1 minute. Then thoroughly scrub all treated surfaces with soap or detergent, and rinse with potable water before reuse.

For Use as a Cleaner and/or Deodorizer

Spray 6-8 inches from hard, non-porous surfaces. Allow to air dry.

READ S.D.S. BEFORE USING PRODUCT

SDS available online at www.virox.com

INTERvention®

INTERvention®

Farm Animal Care

Disinfectant Cleaner & Deodorizer

Ready To Use

ACTIVE INGREDIENT
Hydrogen Peroxide 0.5%
OTHER INGREDIENTS 99.5%
Total 100.0%

EPA Reg. No. 74559-9
EPA Est. No. 74559-CAN-1

Part No. 24110

KEEP OUT OF REACH OF CHILDREN

See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use

FOR BARN AND FARM USE

Net Contents:
30 Gallon (113.6L)



Fragrance Free Formula

30G

For Barn And Farm Use

Claims

See reference sheet for a complete list of organisms.

Bactericidal in 1 Minute:

Staphylococcus aureus

Staphylococcus aureus (Methicillin Resistant (MRSA))

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Klebsiella pneumoniae

Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))

Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))

Klebsiella pneumoniae (Carbapenem Resistant (CRE))

Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))

Staphylococcus aureus (Vancomycin Resistant (VRSA))

Staphylococcus epidermidis (Methicillin Resistant (MRSE))

Enterococcus faecalis (Vancomycin Resistant (VRE))

Enterococcus faecium

Bordetella bronchiseptica

Streptococcus pneumoniae (Penicillin Resistant (PRSP))

Pasteurella multocida

Serratia marcescens

Shigella dysenteriae

*Virucidal in 30 Seconds, unless otherwise noted:

Polio Type 1 (Strain Brunhilde)

Avian Influenza A Virus (H7N9)

Avian Influenza A Virus (H3N2)

Animal premise *Virucidal Activity in 1 Minute, unless otherwise noted:

Bovine Viral Diarrhea Virus – 30 seconds

Duck Hepatitis B virus – 30 seconds

Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

This product has demonstrated effectiveness against Influenza Type A₂ (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Fungicidal in 30 Seconds, unless otherwise noted:

Trichophyton mentagrophytes

Microsporum canis – 1 minute

Tuberculocidal in 1 Minute:

Mycobacterium bovis (BCG)

Broad-Spectrum Non-Food Sanitizer in 10 seconds against:

Klebsiella pneumoniae

Staphylococcus aureus

Enterobacter aerogenes

Acinetobacter baumannii

Listeria monocytogenes

Campylobacter jejuni

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Pseudomonas aeruginosa

Escherichia coli O157:H7

Vancomycin Resistant *Enterococcus faecalis* (VRE)

Methicillin Resistant *Staphylococcus aureus* (MRSA)

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in securely closed container in areas inaccessible to small children. Store away from direct sunlight and heat.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. For more information on the disposal of unused, unwanted product and the cleanup of spills, contact the EPA Regional Office or the Manufacturer for guidance.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill; or incinerate, or, if allowed by state and local authorities, burn. If burned, stay out of smoke.

ENVIRONMENTAL HAZARDS: This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.


Animal Premise *Virucidal
1 Minute

*Virucidal
30 Seconds

Bactericidal
1 Minute

Fungicidal
30 Seconds, unless
otherwise noted

Broad-Spectrum
Non-Food Contact
Sanitizer
10 Seconds



Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com

UPC CODE
0-68460-50485-8

For Barn And Farm Use

This product is a one-step disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, buckets, brushes, automatic feeders, exterior surfaces of watering appliances, farm vehicles (hard non-porous surfaces of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs or any hard non-porous washable (non-food contact) surfaces where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Spray 6-8 inches from surface.
3. Allow surface to remain wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

Animal Housing Facilities

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, stables, etc.

1. Remove all animals and feed from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty (or cover) all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Saturate all surfaces by spraying solution until wet. To disinfect, all surfaces must remain wet for 1 minute.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and water appliances, spray with solution and let stand for 1 minute. Then thoroughly scrub all treated surfaces with soap or detergent, and rinse with potable water before reuse.

For Use as a Cleaner and/ or Deodorizer

Spray 6-8 inches from hard, non-porous surfaces. Allow to air dry.

INTERvention®

INTERvention®

Farm Animal Care Disinfectant Cleaner & Deodorizer

Ready To Use

ACTIVE INGREDIENT
Hydrogen Peroxide 0.5%
OTHER INGREDIENTS 99.5%
Total 100.0%

EPA Reg. No. 74559-9
EPA Est. No. 74559-CAN-1

Part No. 24111

KEEP OUT OF REACH OF CHILDREN

See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use

FOR BARN AND FARM USE



Net Contents:
55 Gallon (208.2L)

55G

Fragrance Free Formula

For Barn And Farm Use

Claims

See reference sheet for a complete list of organisms.

Bactericidal in 1 Minute:

Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Bordetella bronchiseptica
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Pasteurella multocida
Serratia marcescens
Shigella dysenteriae

*Virucidal in 30 Seconds, unless otherwise noted:

Polio Type 1 (Strain Brunhilde)
Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)

Animal premise *Virucidal Activity in 1 Minute, unless otherwise noted:

Bovine Viral Diarrhea Virus – 30 seconds
Duck Hepatitis B virus – 30 seconds
Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

This product has demonstrated effectiveness against Influenza Type A₂ (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Fungicidal in 30 Seconds, unless otherwise noted:

Trichophyton mentagrophytes
Microsporum canis – 1 minute

Tuberculocidal in 1 Minute:

Mycobacterium bovis (BCG)

Broad-Spectrum Non-Food Sanitizer in 10 seconds against:

Klebsiella pneumoniae
Staphylococcus aureus
Enterobacter aerogenes
Acinetobacter baumannii
Listeria monocytogenes
Campylobacter jejuni
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Pseudomonas aeruginosa
Escherichia coli O157:H7
Vancomycin Resistant *Enterococcus faecalis* (VRE)
Methicillin Resistant *Staphylococcus aureus* (MRSA)

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in securely closed container in areas inaccessible to small children. Store away from direct sunlight and heat.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. For more information on the disposal of unused, unwanted product and the cleanup of spills, contact the EPA Regional Office or the Manufacturer for guidance.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill; or incinerate, or, if allowed by state and local authorities, burn. If burned, stay out of smoke.

ENVIRONMENTAL HAZARDS: This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

Animal Premise *Virucidal
1 Minute

*Virucidal
30 Seconds

Bactericidal
1 Minute

Fungicidal
30 Seconds, unless otherwise noted

Broad-Spectrum
Non-Food Contact Sanitizer
10 Seconds

READ S.D.S. BEFORE
USING PRODUCT
SDS available online
at www.virox.com



Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com

UPC CODE
0-68460-50486-5

INTERvention[®]

Farm Animal Care
Disinfectant Cleaner & Deodorizer

Ready to Use

KEEP OUT OF REACH
OF CHILDREN

**FOR BARN
& FARM USE**

EPA Reg. No. 74559-09
EPA Est. No. 74559-CAN-1

This product is a one-step disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, buckets, brushes, automatic feeders, exterior surfaces of watering appliances, farm vehicles (hard, non-porous surface of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs or any hard non-porous washable (non-food contact) surfaces where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Claims

BACTERICIDAL IN 1 MINUTE:

Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo- β -lactamase 1(NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Bordetella bronchiseptica
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Pasteurella multocida
Serratia marcescens
Shigella dysenteriae

TUBERCULOCIDAL IN 1 MINUTE AGAINST:

Mycobacterium bovis (BCG)

FUNGICIDAL IN 30 SECONDS, UNLESS OTHERWISE NOTED AGAINST:

Trichophyton mentagrophytes
Candida albicans – 1 minute
Microsporium canis – 1 minute

Reference Sheet

Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578
virox.com | viroxanimalhealth.com



*VIRUCIDAL IN 30 SECONDS:

Adenovirus Type 8
Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)
Cytomegalovirus
Feline Calicivirus
Hepatitis B Virus (HBV)
Hepatitis C Virus (HCV)
Herpes Simplex Virus Type 1
Herpes Simplex Virus Type 2
HIV Type 1 Strain HTLV III_B
Human Coronavirus
Influenza A Virus VR-544 (Hong Kong Strain)
Influenza B Virus
Norovirus (Feline Calicivirus as the surrogate)
Pandemic Influenza A Virus H1N1
Parainfluenza Virus Type 3
Respiratory Syncytial Virus (RSV)
Rhinovirus R37
Rotavirus (WA strain)
Vaccinia Virus

ANIMAL PREMISE *VIRUCIDAL ACTIVITY IN 1 MINUTE, UNLESS OTHERWISE NOTED

Bovine Viral Diarrhea Virus – **30 seconds**
Duck Hepatitis B virus – **30 seconds**
Equine Influenza A Virus (H3N8) (EIV) – **30 seconds**

This product has demonstrated effectiveness against Influenza Type A₂ (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Mold/Mildew (Mildewcidal Activity) – kills the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mold/Mildew (Mildewstatic Activity) – controls and prevents the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Malodors Activity – eliminates odors and odor-causing bacteria on hard, non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

SOFT SURFACE SANITIZATION AGAINST THE FOLLOWING ORGANISMS IN 30 SECONDS:

Staphylococcus aureus
Enterobacter aerogenes
Methicillin Resistant *Staphylococcus aureus* (MRSA)
Vancomycin Resistant *Enterococcus faecalis* (VRE)

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER IN 10 SECONDS AGAINST:

Staphylococcus aureus

Enterobacter aerogenes

Acinetobacter baumannii

Listeria monocytogenes

Campylobacter jejuni

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Pseudomonas aeruginosa

Escherichia coli O157:H7

Vancomycin Resistant *Enterococcus faecalis* (VRE)

Klebsiella pneumoniae

Methicillin Resistant *Staphylococcus aureus* (MRSA)

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Spray 6-8 inches from surface.
3. Allow surface to remain wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

Animal Housing Facilities

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, stables, etc.

1. Remove all animals and feed from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty (or cover) all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Saturate all surfaces by spraying solution until wet. To disinfect, all surfaces must remain wet for 1 minute.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and water appliances, spray with solution and let stand for 1 minute. Then thoroughly scrub all treated surfaces with soap or detergent, and rinse with potable water before reuse.

For Use as a Cleaner and/or Deodorizer

Spray 6-8 inches from hard, non-porous surfaces. Allow to air dry.

***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS Virus), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces / objects before application of this product.

Contact Time: Allow surface to remain wet for 30 seconds to kill HIV-1, HCV and HBV.

Disposal of Infectious Material: Blood / body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

To Disinfect Non-Critical[‡], Pre-Cleaned Instruments:

1. Instruments must be thoroughly pre-cleaned to remove excess organic debris, rinsed, and rough dried.
2. Clean and rinse lumens of hollow instruments before spraying with this product.
3. Spray all surfaces of instruments with this product until thoroughly wet.
4. Let stand for 1 minute.
5. Wipe with a clean, damp cloth or paper towel and allow to air dry.

[‡]Non-Critical medical devices are items that come in contact only with intact skin.

For use as a Broad Spectrum Non-Food Contact Sanitizer: Pre-clean heavily soiled areas. Spray product directly onto non-porous surfaces until completely wet. Allow the surface to remain visibly wet for 10 seconds. Wipe surfaces dry or allow to air dry.

To Kill Fungi: Pre-clean heavily soiled areas. Spray product directly onto non-porous surfaces until completely wet. Allow surface to remain wet for 30 seconds. Wipe surfaces and let air dry.

To Kill Mold and Mildew: Pre-clean surfaces. Spray product directly onto non-porous surfaces until completely wet. Allow the surface to remain visibly wet for 4 minutes. Allow to air dry.

For use as sanitizer on Soft Surfaces[‡] or Fabrics[‡]:

1. Hold bottle upright 6"-8" from surfaces. 2. Spray until fabric is wet. Do not saturate. 3. Let stand for 30 seconds. Allow to air dry. 4. For difficult odors or heavy fabrics, repeat application. Reapply as necessary.

[‡]composed of cotton or polyester

Allergen Destruction: To destroy specified allergens: Spray, wait 1 minute, and wipe excess. Allow to air dry.

Allergens:

- Cockroach matter -or- particles
- Dust mite matter -or- particles
- Pet dander -or- dog and cat dander