

For Barn And Farm Use

For Use in Barns and Farms

This product is a one-step germicidal disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, and stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, automatic feeders, buckets, brushes, exterior surfaces of watering appliances, farm vehicles (hard, non-porous surface of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs and any hard non-porous washable (non-food contact) surface where disinfection is required. It also reduces odor-

causing bacteria leaving surfaces smelling clean and fresh.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces
3. Allow surface to remain visibly wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

For Use as a Cleaner and/or Deodorizer

Pull towelette from canister and wipe hard, non-porous environmental surfaces. Allow to air dry.

READ S.D.S. BEFORE USING PRODUCT

SDS available online at www.virox.com

8477.12

INTERVENTION®

INTERVENTION®

Farm Animal Care Disinfectant Cleaner & Deodorizer

Ready To Use Wipes

ACTIVE INGREDIENT
Hydrogen Peroxide 0.5%
OTHER INGREDIENTS 99.5%
Total 100.0%

EPA Reg. No. 74559-10
EPA Est. No. 74559-CAN-1
Part No. 24221

KEEP OUT OF REACH OF CHILDREN

See reference sheet for a complete list of pathogenic organisms, additional features, claims, and directions of use

FOR BARN AND FARM USE

Fragrance Free Formula



US PATENT:
7,632,523
& 7,354,604

Net Contents:
1pound 14 ounces
(850 grams)

160 Sheets
6 in. x 6.75 in.

For Barn And Farm Use

Claims

See reference sheet for a complete list of organisms.

Bactericidal in 1 Minute:

Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Bordetella bronchiseptica
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Pasteurella multocida
Serratia marcescens
Shigella dysenteriae

*Virucidal in 30 Seconds:

Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)

Animal premise *Virucidal Activity in 1 Minute, unless otherwise noted:

Bovine Viral Diarrhea Virus – 30 seconds
Duck Hepatitis B virus – 30 seconds
Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

This product has demonstrated effectiveness against Influenza Type A₂ (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Fungicidal in 30 Seconds, unless otherwise noted:

Trichophyton mentagrophytes
Microsporium canis – 1 minute

Tuberculocidal in 1 Minute:

Mycobacterium bovis (BCG)

Broad-Spectrum Non-Food Contact Sanitizer in 10 seconds against:

Staphylococcus aureus
Enterobacter aerogenes
Acinetobacter baumannii

Listeria monocytogenes
Campylobacter jejuni
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Pseudomonas aeruginosa
Escherichia coli O157:H7
Vancomycin Resistant
Enterococcus faecalis (VRE)
Methicillin Resistant
Staphylococcus aureus (MRSA)

Storage and Disposal

Store this product in a cool dry area, away from direct sunlight and heat.

For canisters that are **not intended to be refilled** by consumer.

CONTAINER HANDLING:

Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture. Non-refillable container. Do not reuse or refill this container. Offer empty container for recycling. If recycling is not available, discard container in trash.

Animal Premise *Virucidal
1 Minute

*Virucidal
30 Seconds

Bactericidal
1 Minute

Fungicidal
30 Seconds, unless otherwise noted

Broad-Spectrum Non-Food Contact Sanitizer
10 Seconds



Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com

UPC CODE
0-00000-00000-0

VAH042 (16/034)

For Barn And Farm Use

For Use in Barns and Farms

This product is a one-step germicidal disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, and stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, automatic feeders, buckets, brushes, exterior surfaces of watering appliances, farm vehicles (hard, non-porous surface of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs and any hard non-porous washable (non-food contact) surface where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces
3. Allow surface to remain visibly wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

For Use as a Cleaner and/or Deodorizer

Pull towelette from canister and wipe hard, non-porous environmental surfaces. Allow to air dry.

READ S.D.S. BEFORE USING PRODUCT

SDS available online at www.virox.com

INTERVENTION®

INTERVENTION®

Farm Animal Care Disinfectant Cleaner & Deodorizer

Ready To Use Wipes

ACTIVE INGREDIENT

Hydrogen Peroxide 0.5%

OTHER INGREDIENTS 99.5%

Total 100.0%

EPA Reg. No. 74559-10
EPA Est. No. 74559-CAN-1

Part No. 24224

KEEP OUT OF REACH OF CHILDREN

See reference sheet for a complete list of pathogenic organisms additional features, claims, and directions of use

FOR BARN AND FARM USE

Fragrance Free Formula



Net Contents:
3 pounds 3 ounces
(1,451 grams)
160 Sheets
8 in. x 9 in.
(20.32 cm x 22.86 cm)

For Barn And Farm Use

Claims

See reference sheet for a complete list of organisms.

Bactericidal in 1 Minute:

Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1(NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Bordetella bronchiseptica
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Pasteurella multocida

Serratia marcescens
Shigella dysenteriae

*Virucidal in 30 Seconds:

Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)
Animal premise *Virucidal Activity in 1 Minute, unless otherwise noted:
Bovine Viral Diarrhea Virus – 30 seconds
Duck Hepatitis B virus – 30 seconds
Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

This product has demonstrated effectiveness against Influenza Type A₂ (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Fungicidal in 30 Seconds, unless otherwise noted:

**Trichophyton mentagrophytes*
**Microsporum canis* – 1 minute

Tuberculocidal in 1 Minute

Mycobacterium bovis (BCG)

Broad-Spectrum Non-Food Contact Sanitizer in 10 seconds against:

Staphylococcus aureus

Enterobacter aerogenes
Acinetobacter baumannii
Listeria monocytogenes
Campylobacter jejuni
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Pseudomonas aeruginosa
Escherichia coli O157:H7
Vancomycin Resistant *Enterococcus faecalis* (VRE)
Methicillin Resistant *Staphylococcus aureus* (MRSA)

Storage and Disposal

Store this product in a cool dry area, away from direct sunlight and heat.

For canisters that are **not intended to be refilled** by consumer.

CONTAINER HANDLING

Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture. Non-refillable container. Do not reuse or refill this container. Offer empty container for recycling. If recycling is not available, discard container in trash.

Animal Premise
*Virucidal
1 Minute

*Virucidal
30 Seconds

Bactericidal
1 Minute

Fungicidal
30 Seconds, unless otherwise noted

Broad-Spectrum Non-Food Contact Sanitizer
10 Seconds



Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578 | virox.com

UPC CODE
0-68460-50394-3

INTERvention®

Farm Animal Care
Disinfectant Cleaner & Deodorizer

Ready to Use Wipes

KEEP OUT OF REACH
OF CHILDREN

**FOR BARN
& FARM USE**

EPA Reg. No. 74559-10
EPA Est. No. 74559-CAN-1

This product is a one-step germicidal disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, and stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, automatic feeders, buckets, brushes, exterior surfaces of watering appliances, farm vehicles (hard, non-porous surface of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs and any hard non-porous washable (non-food contact) surface where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Claims

BACTERICIDAL IN 1 MINUTE:

Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Bordetella bronchiseptica
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Pasteurella multocida
Serratia marcescens
Shigella dysenteriae

FUNGICIDAL IN 30 SECONDS, UNLESS OTHERWISE NOTED:

Trichophyton mentagrophytes
Microsporum canis – 1 minute
Candida Albicans – 1 minute

Reference Sheet

Manufactured by:
Virox Technologies Inc.
2770 Coventry Road
Oakville, ON Canada L6H 6R1
1-800-387-7578
virox.com | viroxanimalhealth.com



TUBERCULOCIDAL IN 1 MINUTE:

Mycobacterium bovis (BCG)

*VIRUCIDAL IN 30 SECONDS:

Adenovirus Type 8
Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)
Cytomegalovirus
Feline Calicivirus
Hepatitis B Virus (HBV)
Hepatitis C Virus (HCV)
Herpes Simplex Virus Type 1
Herpes Simplex Virus Type 2
HIV Type 1 Strain HTLV III_B
Human Coronavirus
Influenza A Virus (Hong Kong Strain)
Influenza B Virus
Norovirus (Feline Calicivirus as the surrogate)
Pandemic Influenza A Virus H1N1
Parainfluenza Virus Type 3
Respiratory Syncytial Virus (RSV)
Rhinovirus R37
Rotavirus (WA strain)
Vaccinia Virus

ANIMAL PREMISE *VIRUCIDAL ACTIVITY IN 1 MINUTE, UNLESS OTHERWISE NOTED

Bovine Viral Diarrhea Virus – 30 seconds
Duck Hepatitis B virus – 30 seconds
Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER IN 10 SECONDS AGAINST:

Staphylococcus aureus
Enterobacter aerogenes
Acinetobacter baumannii
Listeria monocytogenes
Campylobacter jejuni
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Pseudomonas aeruginosa
Escherichia coli O157:H7
Vancomycin Resistant *Enterococcus faecalis* (VRE)
Methicillin Resistant *Staphylococcus aureus* (MRSA)

Mildewcidal Activity – kills the growth of mold and mildew a 4 minute contact time *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mildewstatic Activity – controls and prevents the growth of mold and mildew: at a 4 minutes contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Malodor Activity – eliminates odors and odor-causing bacteria on hard, non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

Animal Housing Facilities in Veterinary Clinics, Hospitals, Farm Premise including, Equine production Farm, Bovine production Farm.

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities including pens, chutes, and barns occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with potable water. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
5. All surfaces must remain visibly wet for 1 minute.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
9. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surface to remain visibly wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

For Use as a Cleaner and/or Deodorizer

Pull towelette from canister and wipe hard, non-porous environmental surfaces. Allow to air dry.

***KILLS HIV AND HCV AND HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/ body fluids, and in which the surfaces/objects likely to be soiled with blood/ body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV and all other organisms listed on the label.

Disposal of Infectious Material: Blood/body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

For Use as a Tuberculocide:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

To Kill Fungi:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time.
4. Wipe surfaces dry or allow to air dry.

To Kill *Trichophyton mentagrophytes*:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 30 seconds. Use additional towelettes if needed to ensure the correct wet contact time.
4. Wipe surfaces dry or allow to air dry.

To Kill Mold and Mildew: Pre-clean surfaces. Wipe product onto non-porous surfaces until completely wet. Allow the surface to remain visibly wet for 4 minutes. Allow to air dry.

To Control Mold and Mildew: Wipe product onto non-porous surfaces until completely wet. Allow to air dry. Repeat application weekly or when growth reappears.

For use as a Broad Spectrum Non-Food Contact Sanitizer:

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 10 seconds.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

*VIRUCIDAL | BACTERICIDAL | TUBERCULOCIDAL | FUNGICIDAL | SANITIZER

INTERvention[®]

Manufactured for

VIROX TECHNOLOGIES INC. 1-800-387-7578 | VIROX.COM

For Barn and Farm Use

INTERvention[®]

Farm Animal Care

Disinfectant Cleaner
& Deodorizer

Ready to Use Wipes

See reference sheet for
additional features, claims,
and directions for use.

**KEEP OUT
OF REACH
OF CHILDREN**

10 Net Contents/Sheets
2.6 ounces (75 grams)
7.4in. x 10in. (19.0cm x 25.4cm)

VAH038 (16/011)
Part No. 24223

EPA REG. NO. 74559-10
EPA EST. NO. 075272-WI-001

Fragrance Free Formula



US PATENT:
7,632,523
& 7,354,604

ACTIVE INGREDIENT

Hydrogen Peroxide 0.5%

OTHER INGREDIENTS ... 99.5%

Total 100.0%

PEEL & SEAL / PULL HERE

INTERvention[®]

Farm Animal Care

Disinfectant Cleaner & Deodorizer

Ready To Use Wipes - Soft Pack

KEEP OUT OF REACH
OF CHILDREN

**FOR BARN
& FARM USE**

EPA Reg. No. 74559-10
EPA Est. No. 075272-WI-001

This product is a one-step germicidal disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, fecal matter and other common soils found in barns, farms, and animal transportation vehicles.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, and stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, automatic feeders, buckets, brushes, exterior surfaces of watering appliances, farm vehicles (hard, non-porous surface of), feed racks, foot baths, grooming equipment, handling equipment, pens, transportation crates, troughs and any hard non-porous washable (non-food contact) surface where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

Claims

BACTERICIDAL IN 1 MINUTE:

Staphylococcus aureus

Staphylococcus aureus (Methicillin Resistant (MRSA))

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Klebsiella pneumoniae

Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))

Klebsiella pneumoniae (New Delhi metallo- β -lactamase 1 (NDM-1))

Klebsiella pneumoniae (Carbapenem Resistant (CRE))

Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))

Staphylococcus aureus (Vancomycin Resistant (VRSA))

Staphylococcus epidermidis (Methicillin Resistant (MRSE))

Enterococcus faecalis (Vancomycin Resistant (VRE))

Enterococcus faecium

Bordetella bronchiseptica

Streptococcus pneumoniae (Penicillin Resistant (PRSP))

Pasteurella multocida

Serratia marcescens

Shigella dysenteriae

FUNGICIDAL IN 30 SECONDS, UNLESS OTHERWISE NOTED:

Trichophyton mentagrophytes

Candida Albicans – 1 minute

Microsporum canis – 1 minute

TUBERCULOCIDAL IN 1 MINUTE:

Mycobacterium bovis (BCG)

Reference Sheet

Manufactured for:

Virox Technologies Inc.

2770 Coventry Road

Oakville, ON Canada L6H 6R1

1-800-387-7578

virox.com | viroxanimalhealth.com



*VIRUCIDAL IN 30 SECONDS:

Adenovirus Type 8

Avian Influenza A Virus (H7N9)

Avian Influenza A Virus (H3N2)

Cytomegalovirus

Feline Calicivirus

Hepatitis B Virus (HBV)

Hepatitis C Virus (HCV)

Herpes Simplex Virus Type 1

Herpes Simplex Virus Type 2

HIV Type 1 Strain HTLV III_B

Human Coronavirus

Influenza A Virus (Hong Kong Strain)

Influenza B Virus

Norovirus (Feline Calicivirus as the surrogate)

Pandemic Influenza A Virus H1N1

Parainfluenza Virus Type 3

Respiratory Syncytial Virus (RSV)

Rhinovirus R37

Rotavirus (WA strain)

Vaccinia Virus

ANIMAL PREMISE *VIRUCIDAL ACTIVITY IN 1 MINUTE, UNLESS OTHERWISE NOTED

Bovine Viral Diarrhea Virus – 30 seconds

Duck Hepatitis B virus – 30 seconds

Equine Influenza A Virus (H3N8) (EIV) – 30 seconds

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER IN 10 SECONDS AGAINST:

Staphylococcus aureus

Enterobacter aerogenes

Acinetobacter baumannii

Listeria monocytogenes

Campylobacter jejuni

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Pseudomonas aeruginosa

Escherichia coli O157:H7

Vancomycin Resistant *Enterococcus faecalis* (VRE)

Methicillin Resistant *Staphylococcus aureus* (MRSA)

Mildewcidal Activity - kills the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mildewstatic Activity - controls and prevents the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Malodor Activity – eliminates odors and odor-causing bacteria on hard, non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

Animal Housing Facilities in Veterinary Clinics, Hospitals, Farm Premise including, Equine production Farm, Bovine production Farm.

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities including pens, chutes, and barns occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with potable water. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
5. All surfaces must remain visibly wet for 1 minute.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
9. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
3. Allow surface to remain visibly wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

For Use as a Cleaner and/or Deodorizer

Pull towelette from dispenser and wipe hard, non-porous environmental surfaces. Allow to air dry.

***KILLS HIV AND HCV AND HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS**

in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV and all other organisms listed on the reference sheet.

Disposal of Infectious Material: Blood/body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

For Use as a Tuberculocide:

1. Pre-clean heavily soiled areas.
2. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

To Kill Fungi:

1. Pre-clean heavily soiled areas.
2. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time.
4. Wipe surfaces dry or allow to air dry.

To Kill *Trichophyton mentagrophytes*:

1. Pre-clean heavily soiled areas.
2. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 30 seconds. Use additional towelettes if needed to ensure the correct wet contact time.
4. Wipe surfaces dry or allow to air dry.

To Kill Mold and Mildew: Pre-clean surfaces. Wipe product onto non-porous surfaces until completely wet. Allow the surface to remain visibly wet for 4 minutes. Allow to air dry.

To Control Mold and Mildew: Wipe product onto non-porous surfaces until completely wet. Allow to air dry. Repeat application weekly or when growth reappears.

For use as a Broad Spectrum Non-Food Contact Sanitizer:

1. Pre-clean heavily soiled areas.
2. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 10 seconds.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

For Use as a Cleaner and/or Deodorizer: Pull towelette from dispenser and wipe hard, non-porous environmental surfaces. Allow to air dry.

Re-sealable soft pack:

1. Remove the seal.
2. Remove one sheet at a time.
3. Close the seal to prevent drying between uses.

STORAGE & DISPOSAL

Store this product in a cool dry area, away from direct sunlight and heat.

For soft-packs that are not intended to be refilled by consumer.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do not flush. Tightly close flap between uses to retain moisture. Nonrefillable container.

Do not reuse or refill this container. Offer empty container for recycling. If recycling is not available, discard container in trash.