

FOR USE IN HEALTH CARE, LABORATORY, MEDICAL FACILITIES AND HOSPITALS

Accel TB is a one-step disinfectant cleaner and odor neutralizer designed for general cleaning, and disinfecting, of hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, mold, mildew, and other common soils found in hospitals, nursing homes, and medical offices.

Accel TB's non-dulling formula eliminates the time and labor normally required for rinsing. It is designed for use on hard, non-porous environmental surfaces listed on this label made from: plastic, chrome, stainless steel, aluminum associated with walls, tables, chairs, countertops, telephones, glazed tile, sinks.

Accel TB is an effective cleaner disinfectant sanitizer on non-food contact surfaces.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

FOR USE AS A ONE-STEP CLEANER/ DISINFECTANT PRODUCT

1. Pre-clean heavily soiled areas.
2. Apply solution by spray, cloth, disposable wipe or mop to hard non-porous environmental surfaces.
3. All surfaces must remain wet for 1 minute. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.
4. Wipe surfaces dry.

***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS Virus), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/ BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces / objects before application of Accel TB.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HCV and HBV. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.

Disposal of Infectious Material: Blood / body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

BROAD SPECTRUM NON-FOOD CONTACT SANITIZER

Apply to hard non-porous surface; allow to remain wet for 30 seconds. Wipe dry. For heavily soiled areas a pre-cleaning is required.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container in areas inaccessible to small children. Keep securely closed.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill empty container. Rinse thoroughly. Offer for recycling or discard in trash.

PART NO.
ACCDISR-TB32
EPA REG. NO.
74559-1
EPA EST. NO.
74559-CAN-1

Accel

Net Contents:
1 U.S. quart



Ready to Use One-Step
Disinfectant Cleaner

Bactericide, *Virucide,
Tuberculocide,
Fungicide, Sanitizer

ACTIVE INGREDIENT:

Hydrogen Peroxide 0.5%
Other Ingredients 99.5%
Total 100.0%



USA PAT#: 7,632,523

**KEEP OUT OF REACH
OF CHILDREN**

See reference sheet for additional features
and claims.

READY TO USE

CLAIMS

See reference sheet for a complete list of pathogenic organisms claimed for this product.

BACTERICIDAL: 1 Min.

In the presence of 5% serum load and 1 minute contact time at 20°C on hard, non-porous environmental surfaces.

Acinetobacter baumannii
Enterococcus faecalis, VRE
Escherichia coli O157:H7
Klebsiella pneumoniae
Pseudomonas aeruginosa
Salmonella enterica (formerly known
as *Salmonella choleraesuis*)
Staphylococcus aureus
Staphylococcus aureus MRSA

Bactericidal activity was determined by the AOAC Use Dilution Test Method.

*VIRUCIDAL: 1 Min.

In the presence of 5% serum load and 1 minute contact time at 20°C on hard, non-porous environmental surfaces.

*Hepatitis B Virus (HBV)
*Hepatitis C Virus (HCV)
*HIV-1 (Aids Virus)
*Norovirus (Feline Calicivirus as the surrogate)
*Poliovirus Type 1

*This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

Virucidal activity was determined by the efficacy test methods for virucidal agents intended for inanimate environmental surfaces; ASTM 1053-97 Standard Test Method for Efficacy of Virucidal Agents Intended for Inanimate Environmental Surfaces and EPA protocols for surrogate viral testing.

NON VARNISH
AREA



Manufactured by:
Virox Technologies Inc.
2770 Coventry Rd.
Oakville, ON Canada L6H 6R1
1-800-387-7578 | www.virox.com

TUBERCULOCIDAL: 5 Min.

In the presence of 5% serum load and 5 minute contact time at 20°C on hard, non-porous environmental surfaces.

Mycobacterium bovis
Tuberculocidal activity was determined by the EPA Quantitative Tuberculocidal Activity Test Method.

FUNGICIDAL: 10 Min

In the presence of 5% serum load and 10 minute contact time at 20°C on hard, non-porous environmental surfaces.

Trichophyton mentagrophytes (ATCC 9533)
Fungicidal activity was determined by the AOAC Fungicidal Activity of Disinfectants Method.

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZING: 30 Sec.

In the presence of 5% serum load and 30 second contact time at 20°C on hard, non-porous environmental surfaces.

Escherichia coli O157:H7
Klebsiella pneumoniae
Pseudomonas aeruginosa
Salmonella enterica, (formerly known
as *Salmonella choleraesuis*)
Staphylococcus aureus

Sanitizing activity was determined by the EPA Sanitizer Test for Inanimate, Non-Food Contact Surfaces and the ASTM E1153 Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces.

READ M.S.D.S. BEFORE USING PRODUCT
MSDS available online at www.virox.com

AL121 (13/225)



68460 50383 7

BROAD-SPECTRUM NON-FOOD
CONTACT SANITIZER

FUNGICIDAL

TUBERCULOCIDAL

*VIRUCIDAL

BACTERICIDAL

BACTERICIDAL

*VIRUCIDAL

TUBERCULOCIDAL

FUNGICIDAL

BROAD-SPECTRUM NON-FOOD
CONTACT SANITIZER



LICENSED

PERIOD 2012-2014 LIC. NO.



LICENSED

PERIOD 2015-2017 LIC. NO.

8477.1

FOR USE IN HEALTHCARE, LABORATORY, MEDICAL FACILITIES AND HOSPITALS

Accel TB is a one-step disinfectant cleaner and odor neutralizer designed for general cleaning, and disinfecting, of hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, mold, mildew, and other common soils found in hospitals, nursing homes, and medical offices.

Accel TB's non-dulling formula eliminates the time and labor normally required for rinsing. It is designed for use on hard, non-porous environmental surfaces listed on this label made from: plastic, chrome, stainless steel, aluminum associated with walls, tables, chairs, countertops, telephones, glazed tile, sinks.

Accel TB is an effective cleaner disinfectant sanitizer on non-food contact surfaces.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

BROAD SPECTRUM NON-FOOD CONTACT SANITIZER

Apply to hard non-porous surface; allow to remain wet for 30 seconds. Wipe dry. For heavily soiled areas a pre-cleaning is required.

FOR USE AS A ONE-STEP CLEANER/DISINFECTANT PRODUCT

1. Pre-clean heavily soiled areas
2. Apply solution by spray, cloth, disposable wipe or mop to hard non-porous environmental surfaces.
3. All surfaces must remain wet for 1 minute. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.
4. Wipe surfaces dry.

***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS Virus), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces / objects before application of Accel TB.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HCV and HBV. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.

Disposal of Infectious Material: Blood / body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in dry, well ventilated area away from chemicals, direct light, heat or open flame. Do not mix with other cleaning or disinfecting products.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. For more information on the disposal of unused, unwanted product and the clean up of spills, contact the EPA Regional Office or the Manufacturer for guidance.

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill; or incinerate, or, if allowed by state and local authorities, burn. If burned, stay out of smoke.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting agency has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PART NO. ACCDISR-TB5G
EPA REG. NO. 74559-1
EPA EST. NO. 74559-CAN-1

Accel

Net Contents:
5 Gallon



Ready to Use One-Step Disinfectant Cleaner

Bactericide, *Virucide, Fungicide, Tuberculocide, Sanitizer

ACTIVE INGREDIENT:

Hydrogen Peroxide	0.5%
Other Ingredients	99.5%
Total	100.0%



KEEP OUT OF REACH OF CHILDREN

See reference sheet for additional features and claims.

READY TO USE

CLAIMS

See reference sheet for a complete list of pathogenic organisms claimed for this product.

BACTERICIDAL: 1 Min.

In the presence of 5% serum load and 1 minute contact time at 20°C on hard, non-porous environmental surfaces.

- Acinetobacter baumannii*
- Enterococcus faecalis*, VRE
- Escherichia coli* O157:H7
- Klebsiella pneumoniae*
- Pseudomonas aeruginosa*
- Salmonella enterica* (formerly known as *Salmonella choleraesuis*)
- Staphylococcus aureus*
- Staphylococcus aureus* MRSA

Bactericidal activity was determined by the AOAC Use Dilution Test Method.

***VIRUCIDAL: 1 Min.**

In the presence of 5% serum load and 1 minute contact time at 20°C on hard, non-porous environmental surfaces.

- *Hepatitis B Virus (HBV)
- *Hepatitis C Virus (HCV)
- *HIV-1 (Aids Virus)
- *Norovirus (Feline Calicivirus as the surrogate)
- *Poliovirus Type 1

*This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

Virucidal activity was determined by the efficacy test methods for virucidal agents intended for inanimate environmental surfaces; ASTM 1053-97 Standard Test Method for Efficacy of Virucidal Agents Intended for Inanimate Environmental Surfaces and EPA protocols for surrogate viral testing.



Manufactured by:
Virox Technologies Inc.
2770 Coventry Rd.
Oakville, ON Canada L6H 6R1
1-800-387-7578 | www.virox.com

TUBERCULOCIDAL: 5 Min.

In the presence of 5% serum load and 5 minute contact time at 20°C on hard, non-porous environmental surfaces.

Mycobacterium bovis
Tuberculocidal activity was determined by the EPA Quantitative Tuberculocidal Activity Test Method.

FUNGICIDAL: 10 Min.

In the presence of 5% serum load and 10 minute contact time at 20°C on hard, non-porous environmental surfaces.

Trichophyton mentagrophytes (ATCC 9533)
Fungicidal activity was determined by the AOAC Fungicidal Activity of Disinfectants Method.

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZING: 30 Sec.

In the presence of 5% serum load and 30 second contact time at 20°C on hard, non-porous environmental surfaces.

- Escherichia coli* O157:H7
- Klebsiella pneumoniae*
- Pseudomonas aeruginosa*
- Salmonella enterica*, (formerly known as *Salmonella choleraesuis*)
- Staphylococcus aureus*

Sanitizing activity was determined by the EPA Sanitizer Test for Inanimate, Non-Food Contact Surfaces and the ASTM E1153 Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces.

READ M.S.D.S. BEFORE USING PRODUCT
MSDS available online at www.virox.com

NON VARNISH AREA





Accel[®] TB

Ready To Use One-Step Disinfectant Cleaner

Reference Sheet

BACTERICIDAL: 1 MINUTE

In the presence of 5% serum load and 1 minute contact time on hard, non-porous environmental surfaces against

Acinetobacter baumannii (ATCC 19606)

Enterococcus faecalis VRE (ATCC 51575)

Escherichia coli O157:H7 (ATCC 35150)

Escherichia coli with extended beta-lactamase resistance (ESBL) (ATCC BAA-196)

Klebsiella pneumoniae (ATCC 4352)

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica (formerly known as *Salmonella choleraesuis*) (ATCC 10708)

Shigella dysenteriae (ATCC 11835)

Staphylococcus aureus (ATCC 6538)

Staphylococcus aureus (CA-MRSA) (NARSA NRS 384) (Genotype US300)

Staphylococcus aureus MRSA (ATCC 33592)

Community Associated Methicillin Resistant *Staphylococcus aureus* (CA-MRSA) (NARSA NRS 123) (Genotype US400) Community Associated Methicillin Resistant

Bactericidal activity was determined by the AOAC Use Dilution Test Method.

*VIRUCIDAL: 1 MINUTE

In the presence of 5% serum load and 1 minute contact time on hard, non-porous environmental surfaces

*Adenovirus type 8 (ATCC VR-1368)

*Avian Influenza A (ATCC VR-2072)

*Feline Calicivirus, Strain F9 (ATCC VR-782)

*Hepatitis B Virus (HBV)

*Hepatitis C Virus (HCV)

*Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)

*Herpes Simplex Virus, Type 2 (HSV-2) (ATCC VR-734)

*HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1), (Acquired from Adv. Biotechnologies, Inc.)

*Human Coronavirus (ATCC VR-740)

*Influenza A/Hong Kong (ATCC VR-544)

*Norovirus (Feline Calicivirus, as the surrogate)

*Poliovirus Type 1, Strain Brunhilde (ATCC VR-1000)

*Rhinovirus Type 37, Strain 151-1 (ATCC VR-1147)

*Rotavirus WA (Acquired from University of Ottawa)

*This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.



Manufactured by:
Virox Technologies Inc.
2770 Coventry Rd., Oakville, ON L6H 6R1
1-800-387-7578 | www.virox.com



Virucidal activity was determined by the efficacy test methods for virucidal agents intended for inanimate environmental surfaces; ASTM 1053-97 Standard Test Method for Efficacy of Virucidal Agents Intended for Inanimate Environmental Surfaces and EPA protocols for surrogate viral testing.

TUBERCULOCIDAL: 5 MINUTES

In the presence of 5% serum load and 5 minute contact time on hard, non-porous environmental surfaces

Mycobacterium bovis (BCG) (OT 451C150)

Tuberculocidal activity was determined by the EPA Quantitative Tuberculocidal Activity Test Method.

FUNGICIDAL: 10 MINUTES

In the presence of 5% serum load and 10 minute contact time on hard, non-porous environmental surfaces

Tricophyton mentagrophytes (ATCC 9533)

Fungicidal activity was determined by the AOAC Fungicidal Activity of Disinfectants Method

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZING: 30 SEC

In the presence of 5% serum load and 30 second contact time on hard, non-porous environmental surfaces

Enterococcus faecalis VRE (ATCC 51575)

Escherichia coli O157:H7 (ATCC 35150)

Klebsiella pneumoniae (ATCC 4352)

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica (formerly known as *Salmonella choleraesuis*) (ATCC 10708)

Staphylococcus aureus (ATCC 6538)

Staphylococcus aureus MRSA (ATCC 33592)

Sanitizing activity was determined by the EPA Sanitizer Test for Inanimate, Non-Food Contact Surfaces and ASTM E1153 Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces.

SOFT SURFACE SANITIZATION

For use on soft surfaces** found in hospitals, nursing homes, healthcare facilities, and medical offices.

SOFT SURFACE SANITIZATION AGAINST THE FOLLOWING ORGANISMS IN 1 MINUTE:

Enterobacter aerogenes (ATCC 13048)

Staphylococcus aureus (ATCC 6538)

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

To Sanitize Soft Surfaces:**

1. Hold bottle upright 6" - 8" from surfaces.
2. Spray fabric until wet. DO NOT SATURATE.
3. Let stand for 1 minute. Allow to air dry.

** composed of cotton or polyester



EPA Reg. No. 75559-1

Ready to Use One-Step Disinfectant Cleaner & Deodorizer

Animal Housing Facilities

See Reference Sheet for complete list of pathogenic organisms

This product cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, poultry facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It also eliminates odor leaving surfaces smelling clean and fresh.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For cleaning and disinfecting the following hard nonporous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including pens, chutes, and barns) occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Saturate surfaces by wiping, [or] spraying [or mopping] with product for a period of 1 minute. Use a 5 minute contact time for TB and a 10 minute contact for fungi.
5. Immerse handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure.
6. Ventilate buildings and other closed spaces.
7. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

Environmental Hazards

This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

