



Manufactured by:
Virox Technologies Inc.
 2770 Coventry Rd.
 Oakville, ON Canada L6H 6R1
 1-800-387-7578 | www.virox.com

**NO VARNISH
 AREA**

* See reference sheet for additional features, claims and directions for use.

FOR USE AS A ONE-STEP CLEANER/ DISINFECTANT

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 1 minute. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.
4. Allow to air dry.

FOR USE AS NON-FOOD CONTACT SANITIZER

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surface to remain wet for 30 seconds.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

PART NO. ACCWP20-TBUS
 EPA REG. NO. 74559-3
 EPA EST. NO. 74559-CAN-1

KEEP OUT OF REACH OF CHILDREN

READ M.S.D.S. BEFORE USING PRODUCT
 M.S.D.S. is available online at www.virox.com

Net Contents:
4 ounces (113 grams)
20 Sheets: 6 in. x 6.75 in.
 (15.24 cm x 17.14 cm)

Accel



One-Step Disinfectant Cleaner Wipes

Bactericidal, *Virucidal, Tuberculocidal, Fungicidal, Sanitizer

ACTIVE INGREDIENT:
 Hydrogen Peroxide0.5%
 Inert Ingredients99.5%
 Total100.0%



USA PAT# 7,632,523

FOR USE IN HEALTH CARE, LABORATORY, MEDICAL FACILITIES AND HOSPITALS.

USE ON NON-FOOD CONTACT SURFACES.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for personal cleansing. **THIS IS NOT A BABY WIPE.** Do not flush in toilet. This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. For Canisters that are not intended to be refilled by consumer.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do NOT flush. Tightly close lid between uses to retain moisture. Nonrefillable container. Do not reuse or refill this container. Offer empty container for recycling. If recycling is not available, discard container in trash.

AL136 (13/206)

WIPES

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER

BACTERICIDAL

***VIRUCIDAL**

TUBERCULOCIDAL

FUNGICIDAL



STATE OF HAWAII
 Department of Agriculture

ACCEPTED

LICENSE NO.

8477.2

DIELINE DOES NOT PRINT

Part No: AL136

Rev No: 13/206

Replaces: AL136 (13/138)

DECO 28 Greensboro Dr.
 Etobicoke, ON
 M9W 1E1

LABELS & TAGS

Proof Date: JULY 29, 2013
CLIENT: VIROX TECHNOLOGIES
CLIENT P/N: AL136 ACCEL TB WIPES ONE STEP DISINFECT CLEANER WIPES
DECO P/N: VIR003-DPN022930A
Size: 2" x 10.118"
Production Artist: CR

ARTWORK SPECIFICATIONS

PANTONE® 185	PANTONE® REF BLUE	PANTONE® Cool Gray 2	PANTONE® BLACK	SPOT UV

This proof is for CONTENT ONLY. It is not to be used for color match. Colors will be matched using the Pantone Matching System (PMS).

COPY POSITION

1	2	3	4
COPY POSITION: 1-4 LABELS WOUND OUT			
5	6	7	8
COPY POSITION: 5-8 LABELS WOUND IN			

9 ANY COPY POSITION

CLIENT ARTWORK APPROVAL AREA

PLEASE COMPLETE APPROVAL AREA IN FULL
Client assumes sole responsibility for printed artwork and copy accuracy.

<input checked="" type="checkbox"/> All Copy Correct <input checked="" type="checkbox"/> Layout Correct <input checked="" type="checkbox"/> Approved as shown	<input type="checkbox"/> Changes required
	<input type="checkbox"/> Not approved

All 3 must be checked to proceed with production!

CLIENT SIGNATURE _____

DATE _____

NOTE TO CLIENT: This block with signature must be faxed or returned to Deco Labels & Tags before production can begin. **Client assumes sole responsibility for printed artwork and copy accuracy.**

PROOF REV. 002 AUG 2011 / Approved by General Manager

FOR USE IN HEALTH CARE, LABORATORY, MEDICAL FACILITIES AND HOSPITALS

This product is a non-woven, disposable, pre-saturated wipe with Hydrogen Peroxide disinfectant cleaner.

Accel TB Wipe is a one-step disinfectant cleaner and odor neutralizer designed for general cleaning, and disinfecting, of hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, dead skin, blood, body soils and other common soils found in hospitals, nursing homes and medical offices.

Accel TB Wipe's non-dulling formula eliminates the time and labor normally required for rinsing. It is designed for use on hard, non-porous environmental surfaces: tables, chairs, countertops, telephones, glazed tile and sinks.

Use on non-foam contact surfaces.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

FOR USE AS A CLEANER AND/OR DEODORIZER

Pull towlette from canister and wipe hard, non-porous environmental surfaces. Allow to air dry.

READ M.S.D.S. BEFORE USING PRODUCT
MSDS available online at www.virox.com

FOR USE AS ONE-STEP CLEANER/ DISINFECTANT:

1. Pre-clean heavily soiled areas.
2. Pull towlette from canister and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 1 minute. Use a 5 minute contact time for Tuberculosis (TB) and a 10 minute contact time for fungi.
4. Allow to air dry.

***KILLS HIV AND HCV AND HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/ OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV. Use a 5 minute contact time for Tuberculosis (TB) and a 10 minute contact time for fungi.

Disposal of Infectious Material: Blood/body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

PART NO.
ACCWIPB-TBUS
EPA REG. NO.
74559-3
EPA EST. NO.
74559-CAN-1

Accel

Net Contents:
1.1 pounds
(493 grams)
60 Sheets:
7 in. x 8 in.
(17.8 cm x 20.3 cm)

Accel[®] TB WIPES

One-Step Disinfectant Cleaner Wipes

Bactericidal, *Virucidal, Tuberculocidal, Fungicidal, Sanitizer

ACTIVE INGREDIENT:

Hydrogen Peroxide 0.5%
Inert Ingredients 99.5%
Total 100.0%



KEEP OUT OF REACH OF CHILDREN

See reference sheet for additional features, claims and directions for use.

WIPES

FOR USE AS A NON-FOOD CONTACT SANITIZER:

1. Pre-clean heavily soiled areas.
2. Pull towlette from dispenser and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 30 seconds.
4. Wipe surfaces dry.
5. Change towlette after each use.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage and disposal.

For canisters that are **not intended to be refilled** by consumer.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture. Nonrefillable container. Do not reuse or refill this container. Offer empty container for recycling. If recycling is not available, discard container in trash.

CLAIMS

See reference sheet for a complete list of organisms.

BACTERICIDAL: 1 Min.

In the presence of 5% serum load and 1 minute contact time at 20°C on hard, non-porous environmental surfaces.

Acinetobacter baumannii,
Enterococcus faecalis, VRE
Escherichia coli O157:H7
Pseudomonas aeruginosa
Salmonella enterica, (formerly known as *Salmonella choleraesuis*)
Staphylococcus aureus
Staphylococcus aureus MRSA

*VIRUCIDAL: 1 Min.

In the presence of 5% serum load and 1 minute contact time at 20°C on hard, non-porous environmental surfaces.

*Poliovirus Type 1, Strain Brunhilde

Kills HIV-1, (AIDS virus) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

Kills HBV & HCV when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

Kills Norovirus (Feline Calicivirus as the surrogate) when used as directed on hard non-porous inanimate surfaces with a 1 minute contact time.

*This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

TUBERCULOCIDAL: 5 Min.

In the presence of 5% serum load and 5 minute contact time at 20°C on hard, non-porous environmental surfaces.

Mycobacterium bovis

FUNGICIDAL: 10 Min.

Trichophyton mentagrophytes

NON-FOOD CONTACT SANITIZER FOR HARD NON-POROUS SURFACES : 30 Sec.

In the presence of 5% serum load and 30 second contact time at 20°C on hard, non-porous environmental surfaces.

Escherichia coli O157:H7

Klebsiella pneumoniae

Pseudomonas aeruginosa

Salmonella enterica, (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus

Notice to User: Accel TB may not be suitable for all items or surfaces. Prolonged exposure to copper, brass, anodized aluminum and some coatings should be avoided. If you are uncertain as to the material composition of your item, confirm with the manufacturer before proceeding or visit the compatibility section on www.virox.com for a listing of compatible materials.

Manufactured by:
Virox Technologies Inc.
2770 Coventry Rd.
Oakville, ON Canada L6H 6R1
1-800-387-7578 www.virox.com



NON VARNISH AREA

AL127 (13/221)

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER

STATE OF HAWAII
Department of Agriculture

FUNGICIDAL • TUBERCULOCIDAL • *VIRUCIDAL • BACTERICIDAL

BACTERICIDAL • *VIRUCIDAL • TUBERCULOCIDAL • FUNGICIDAL • BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER

ACCEPTED

LICENSE NO.

8477.2


DIELINE DOES NOT PRINT

DECO 28 Greensboro Dr.
Etobicoke, ON
M9W 1E1
LABELS & TAGS

ARTWORK SPECIFICATIONS



This proof is for CONTENT ONLY. It is not to be used for color match. Colors will be matched using the Pantone Matching System (PMS).



9 ANY COPY POSITION

CLIENT ARTWORK APPROVAL AREA

PLEASE COMPLETE APPROVAL AREA IN FULL

Client assumes sole responsibility for printed artwork and copy accuracy.

- all 3
- All Copy Correct
 - Layout Correct
 - Approved as shown

- Changes required
- Not approved

All 3 must be checked to proceed with production!

CLIENT SIGNATURE

DATE

NOTE TO CLIENT: This block with signature must be faxed or returned to Deco Labels & Tags before production can begin. Client assumes sole responsibility for printed artwork and copy accuracy.

PROOF REV. 002 AUG 2011 / Approved by General Manager

Part No: AL127
Rev: 13/221
Replaces: AL130
Latest revision
(Replace only after instructed by Virox)

FOR USE AS A ONE-STEP CLEANER/DISINFECTANT

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 1 minute. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.
4. Allow to air dry.

FOR USE AS NON-FOOD CONTACT SANITIZER

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow surface to remain wet for 30 seconds.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

READ M.S.D.S. BEFORE USING PRODUCT
M.S.D.S. is available online at www.virox.com



Manufactured by:
Virox Technologies Inc.
2770 Coventry Rd.
Oakville, ON Canada L6H 6R1
1-800-387-7578
www.virox.com

PART NO.
ACCWIPTBUS-12X12RP

EPA REG. NO.
74559-3

EPA EST. NO.
74559-CAN-1

Accel

Net Contents:
4 lbs. 2.6 oz.
(1.89 kg)
110 sheet
12 in. x 12 in.
(30.48 cm x 30.48 cm)

Accel[®] TB WIPES

One-Step Disinfectant Cleaner Wipes

Bactericidal, *Virucidal, Tuberculocidal,
Fungicidal, Sanitizer

ACTIVE INGREDIENT:

Hydrogen Peroxide	0.5%
Inert Ingredients	99.5%
Total	100.0%



USA PAT#: 7,632,523

KEEP OUT OF REACH OF CHILDREN

WIPES

FOR USE IN HEALTH CARE, LABORATORY, MEDICAL FACILITIES, AND HOSPITALS

Use on non-food contact surfaces.

DIRECTIONS FOR USE: STORAGE AND DISPOSAL:


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for personal cleansing. **THIS IS NOT A BABY WIPE.** Do not flush in toilet. This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

REFILL POUCH: Look for product expiration date on back of refill pouch. Remove lid from empty bucket. Place refill pouch into bucket. Tear open refill pouch. DO NOT remove wipe roll from refill pouch. Pull out corner of first wipe from centre of roll and push ½" to ¾" (1 to 2 cm) through "X" opening on lid. DO NOT push finger through opening. Replace lid and pull wipe at an angle. The next wipe pops up automatically. When finished, close lid to retain moisture. If bucket becomes contaminated or damaged replace immediately. Replace bucket immediately if tabs below lid become damaged.

For buckets, lids, or pouches intended to be refilled by the consumer.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do not flush in toilet. Tightly reseal bucket flap between uses to retain moisture. Do not reuse or refill except as described in the directions for use. Refill only with this product. Offer empty container for recycling. If recycling is not available, wrap and discard in trash.

* See reference sheet for additional features, claims and directions for use.



AL151(14/010)

BACTERICIDAL

*VIRUCIDAL

TUBERCULOCIDAL

FUNGICIDAL

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER



STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO.
8477.2

PART NO.
ACCWIP11x12-TBUS
15/075

One-Step Disinfectant Cleaner Wipes
Bactericidal, *Virucidal, Tuberculocidal,
Fungicidal, Sanitizer

ACTIVE INGREDIENT:
Hydrogen Peroxide 0.5%
Inert Ingredients..... 99.5%
Total 100.0%

EPA REG. NO.
74559-3
EPA EST. NO.
74559-CAN-1



USA PAT#: 7,632,523

KEEP OUT OF REACH OF CHILDREN

Net Contents: 6.2 lbs. (2.8 kg)
160 Wipes, 11 in. x 12 in. (28 cm x 30 cm)

BACTERICIDAL: 1 MIN.
*VIRUCIDAL: 1 MIN.
TUBERCULOCIDAL: 5 MIN.

FUNGICIDAL: 10 MIN.
BROAD-SPECTRUM NON-FOOD
CONTACT SANITIZER: 30 SEC.

BACTERICIDAL
*VIRUCIDAL
TUBERCULOCIDAL
FUNGICIDAL
BROAD-SPECTRUM NON-FOOD
CONTACT SANITIZER

BACTERICIDAL: 1 MIN.
*VIRUCIDAL: 1 MIN.
TUBERCULOCIDAL: 5 MIN.

FUNGICIDAL: 10 MIN.
BROAD-SPECTRUM NON-FOOD
CONTACT SANITIZER: 30 SEC.

FOR USE AS ONE-STEP CLEANER/DISINFECTANT

1. Pre-clean heavily soiled areas.
2. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 1 minute. Use a 5 minute contact time for Tuberculosis (Tb) and a 10 minute contact time for fungi.
4. Allow to air dry.

FOR USE AS A NON-FOOD CONTACT SANITIZER

1. Pre-clean heavily soiled areas.
2. Pull towelette from dispenser and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 30 seconds.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

READ S.D.S. BEFORE USING PRODUCT
S.D.S. is available online at www.virox.com



Manufactured by:
Virox Technologies Inc.
2770 Coventry Rd.
Oakville, ON Canada L6H 6R1
1-800-387-7578 | www.virox.com

**FOR USE IN HEALTH CARE, LABORATORY,
MEDICAL FACILITIES, AND HOSPITALS**

Use on non-food contact surfaces.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for personal cleansing. **THIS IS NOT A BABY WIPE.** Do not flush in toilet. This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

STORAGE AND DISPOSAL:

Store this product in a cool dry area.

For buckets intended to be refilled by the consumer.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do not flush in toilet. Tightly reseal bucket flap between uses to retain moisture. Do not reuse or refill except as described in the directions for use. Refill only with this product. Offer empty container for recycling. If recycling is not available, wrap and discard in trash.

See reference sheet for additional features,
claims and directions for use.



BACTERICIDAL: 1 MIN.
*VIRUCIDAL: 1 MIN.
TUBERCULOCIDAL: 5 MIN.

FUNGICIDAL: 10 MIN.
BROAD-SPECTRUM NON-FOOD
CONTACT SANITIZER: 30 SEC.

BACTERICIDAL
*VIRUCIDAL
TUBERCULOCIDAL
FUNGICIDAL
BROAD-SPECTRUM NON-FOOD
CONTACT SANITIZER

BACTERICIDAL: 1 MIN.
*VIRUCIDAL: 1 MIN.
TUBERCULOCIDAL: 5 MIN.

FUNGICIDAL: 10 MIN.
BROAD-SPECTRUM NON-FOOD
CONTACT SANITIZER: 30 SEC.



**FOR USE AS ONE-STEP
CLEANER/DISINFECTANT**

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. All surfaces must remain visibly wet for 1 minute. Use a 5 minute contact time for Tuberculosis (Tb) and a 10 minute contact time for fungi.
4. Allow to air dry.

**FOR USE AS A NON-FOOD
CONTACT SANITIZER**

1. Pre-clean heavily soiled areas.
2. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 30 seconds.
4. Wipe surfaces dry or allow to air dry.
5. Change towelette after each use.

READ S.D.S. BEFORE USING PRODUCT
S.D.S. is available online at www.virox.com



Manufactured by:
Virox Technologies Inc.
2770 Coventry Rd.
Oakville, ON Canada L6H 6R1
1-800-387-7578
www.virox.com

PART NO.
ACCWIP11x12RP-TBUS
15/075

EPA REG. NO.
74559-3

EPA EST. NO.
74559-CAN-1

Accel

Net Contents:
6.2 lbs.
(2.8 kg)

160 Wipes,
11 in. x 12 in.
(28 cm x 30 cm)



One-Step Disinfectant Cleaner Wipes

Bactericidal, *Virucidal, Tuberculocidal,
Fungicidal, Sanitizer

ACTIVE INGREDIENT:

Hydrogen Peroxide	0.5%
Inert Ingredients.....	99.5%
Total	100.0%



USA PAT#: 7,632,523

KEEP OUT OF REACH OF CHILDREN

WIPES

**FOR USE IN HEALTH CARE, LABORATORY,
MEDICAL FACILITIES, AND HOSPITALS**

Use on non-food contact surfaces.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for personal cleansing. **THIS IS NOT A BABY WIPE.** Do not flush in toilet. This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semicritical medical devices prior to sterilization or high level disinfection.

REFILL POUCH: Look for product expiration date on back of refill pouch. Remove lid from empty bucket. Place refill pouch into bucket. Tear open refill pouch. DO NOT remove wipe roll from refill pouch. Pull out corner of first wipe from centre of roll and push 1/2" to 3/4" (1 to 2 cm) through "X" opening on lid. DO NOT push finger through opening. Replace lid and pull wipe at an angle. The next wipe pops up automatically. When finished, close lid to retain moisture. If bucket becomes contaminated or damaged replace immediately. Replace bucket immediately if tabs below lid become damaged.

STORAGE AND DISPOSAL:

For buckets, lids, or pouches intended to be refilled by the consumer.

CONTAINER HANDLING: Dispose of wipe in trash after use. Do not flush in toilet. Tightly reseal bucket flap between uses to retain moisture. Do not reuse or refill except as described in the directions for use. Refill only with this product. Offer empty container for recycling. If recycling is not available, wrap and discard in trash.

See reference sheet for additional features, claims and directions for use.



AL174 (15/079)

BACTERICIDAL

*VIRUCIDAL

TUBERCULOCIDAL

FUNGICIDAL

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER



Manufactured by:
Virox Technologies Inc.
2770 Coventry Rd., Oakville, ON L6H 6R1
1-800-387-7578 | www.viroxaccel.com



Reference Sheet

FOR USE IN HEALTH CARE, LABORATORY, MEDICAL FACILITIES AND HOSPITALS

Accel TB is a one-step disinfectant cleaner and odor neutralizer designed for general cleaning, and disinfecting, of hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, mold, mildew, and other common soils found in hospitals, nursing homes, and medical offices.

Accel TB's non-dulling formula eliminates the time and labor normally required for rinsing. It is designed for use on hard, non-porous environmental surfaces listed on this label made from: plastic, chrome, stainless steel, aluminum associated with walls, tables, chairs, countertops, telephones, glazed tile, sinks.

BACTERICIDAL: 1 MINUTE

HOSPITAL DISINFECTANT: BACTERICIDAL in the presence of 5% serum load and 1 minute contact time on hard, non-porous environmental surfaces.

BACTERICIDAL ACTIVITY:

Acinetobacter baumannii [ATCC 19606]

Escherichia coli O157:H7 [ATCC 35150]

Pseudomonas aeruginosa [ATCC 15442]

Salmonella enterica [ATCC 10708] formerly known as *Salmonella choleraesuis*

Staphylococcus aureus [ATCC 6538]

ANTIBIOTIC-RESISTANT BACTERIA:

Enterococcus faecalis VRE [ATCC 51575]

Staphylococcus aureus MRSA [ATCC 33592]

*VIRUCIDAL: 1 MINUTE

***VIRUCIDAL** in the presence of 5% serum load and 1 minute contact time on hard, non-porous environmental surfaces.

*VIRUSES:

*Herpes Simplex Virus, Type 1 [HSV-1] [ATCC VR-733]

*Herpes Simplex Virus, Type 2 [HSV-2] [ATCC VR-734]

*Human Coronavirus [ATCC VR-740]

*Influenza A/Hong Kong [ATCC VR-544]

*Poliovirus Type 1, Strain Brunhilde [ATCC VR-1000]

*Rhinovirus Type 14 [ATCC VR-284]

*Rotavirus WA

*Kills HIV-1 [AIDS virus] [HTLV-III_g] when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

*Kills HBV & HCV when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

*Kills Norovirus [Feline Calicivirus as the surrogate] [VR-782] when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

*This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A Viruses including Pandemic 2009 H1N1 Influenza A virus.

VETERINARY *VIRUSES:

*Avian Influenza A, H3N2 [ATCC VR-2072]

*Feline Calicivirus, Strain F9 [ATCC VR-782]

TUBERCULOCIDAL: 5 MINUTES

TUBERCULOCIDAL in the presence of 5% serum load and 5 minute contact time on hard, non-porous environmental surfaces.

Mycobacterium bovis [Tb] [OT 451C150]

FUNGICIDAL: 10 MINUTES

FUNGICIDAL This product is fungicidal against the pathogenic fungi, *Trichophyton mentagrophytes* [ATCC 9533] [the fungus which causes Athlete's Foot] in the presence of 5% serum load and 10 minute contact time when used as directed on hard surfaces found in bathroom, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

Trichophyton mentagrophytes [ATCC 9533]

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 30 SECONDS

NON-FOOD CONTACT SURFACE SANITIZER FOR HARD NON-POROUS SURFACES – in the presence of 5% serum load and 30 second contact time on hard, non-porous environmental surfaces.

Enterococcus faecalis VRE [ATCC 51575]

Escherichia coli O157:H7 [ATCC 35150]

Klebsiella pneumoniae [ATCC 4352]

Pseudomonas aeruginosa [ATCC 15442]

Staphylococcus aureus [ATCC 6538]

Staphylococcus aureus MRSA [ATCC 33592]

Salmonella enterica [ATCC 10708] formerly known as *Salmonella choleraesuis*

***KILLS HIV AND HCV AND HBV ON pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings [Hospitals, Nursing Homes] and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/ body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV. Use a 5 minute contact time for Tuberculosis [Tb] and 10 minute contact for fungi.

Disposal of Infectious Material: Blood/body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

The logo for Accel TB Wipes features a grid of blue dots above the word "Accel" in red, "TB" in blue, and "Wipes" in blue. Below this, the text "One-Step Disinfectant Wipes" is written in blue.

Accel[®] TB Wipes

One-Step Disinfectant Wipes

EPA Reg. No. 74559-3

Animal Housing Facilities

See Reference Sheet for complete list of organisms

This product cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It also eliminates odor leaving surfaces smelling clean and fresh.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For cleaning and disinfecting the following hard nonporous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including pens, chutes, and barns) occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with potable water. Pull towelette from canister and wipe hard, non-porous environmental surfaces.
5. All surfaces must remain visibly wet for 1 minute.
6. Allow to air dry
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
9. Thoroughly scrub all treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

NOTE: Not for use on handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure.

Environmental Hazards

This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

