

PEROXcide Concentrate Disinfectant Cleaner

Animal Housing Facilities

This product is designed for use as a one-step cleaner, disinfectant, sanitizer on hard, non-porous surfaces of vehicles, trucks, trailers, carts, animal crates, grooming equipment, handling equipment, buckets, brushes, tables, dog runs, kennels, cages, transportation crates, foot baths, automatic feeders, troughs, feed racks, wash stalls, non-porous fencing, gates, gathering, guiding / controlling equipment, and non-porous harnesses & leashes; commonly found in animal housing facilities, poultry, stall, barn, pen, agricultural & livestock buildings, greenhouses, zoos, boarding kennels, veterinary clinics, animal rescue and shelter facilities, fish facilities, fish tanks, fish farms, hatcheries, tack shops, pet stores, breeding facilities, grooming facilities and catteries.

For use as a *Virucide: Dilute at 2 oz. – 8 oz. of product per gallon of water (1:16 -1:64)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 5 minutes.
4. Wipe surfaces and let air dry.

For Treatment of Animal Housing Facilities (including Poultry barns): Dilute at 8.0 oz. of product per gallon of water (1:16)

1. Remove all animals and feeds from areas being treated.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
3. Empty or cover all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use Dilution Made Easy

Ratio	Per Gallon of Water	(Per Liter of Water)
1:16	8 oz.	64 mL
1:64	2 oz.	16 mL
1:128	1 oz.	8 mL

For Use as a One-Step Cleaner/Disinfectant: Dilute at 8.0 oz. of product per gallon of water (1:16)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution by coarse trigger sprayer to hard, non-porous environmental surfaces. Spray 6-8 inches from surface; making sure to wet surfaces thoroughly.
3. All surfaces must remain wet for 5 minutes.
4. Wipe surfaces and let air dry.

To Sanitize Non-Food Contact Surfaces: Dilute at 1.0 oz. of product per gallon of water (1:128)

1. Pre-clean heavily soiled hard non-porous surfaces.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 3 minutes.
4. Wipe surfaces and let air dry.
5. Not for use on food contact surfaces or on food preparation areas.

To Kill *Minute virus of mice: Dilute at 8.0 oz. of product per gallon of water (1:16)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 5 minutes.
4. Wipe surfaces and let air dry.

To Kill *Canine Parvovirus: Dilute at 4.0 oz. of product per gallon of water (1:32)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 10 minutes.
4. Wipe surfaces and let air dry.

To Kill Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16). Pre-clean surfaces. Apply Use Solution to hard, non-porous environmental surfaces. Allow surfaces to remain wet for 10 minutes. Wipe surfaces and let air dry.

For Use as a Deodorizer: Dilute at 8 oz. of product per gallon of water (1:16). Apply to hard, non-porous surfaces. Let stand for 30 seconds.

To Kill Fungi: Dilute at 8.0 oz. of product per gallon of water (1:16)

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for 5 minutes. Wipe surfaces and let air dry.

To Control Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16).

Apply Use Solution to pre-cleaned hard, non-porous environmental surfaces. Wipe surfaces and let air dry. Repeat application weekly or when growth reappears.

For Use as a Cleaner/Degreaser: Dilute product at 0.5 - 1.0 oz. of product per gallon of water (1:128 - 1:256). Apply to surfaces. Wipe surfaces and let air dry.

PEROXcide

Concentrate Disinfectant Cleaner

Multi-Surface One Step Disinfectant Cleaner and Deodorizer

ACTIVE INGREDIENT
Hydrogen Peroxide 4.25%
INERT INGREDIENTS 95.75%
Total 100.00%

EPA Reg. No. 74559-4
EPA Est. No. 74559-CAN-1



See Additional Precautionary Statements on Side Panel
See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use.

UNSCENTED

KEEP OUT OF REACH OF CHILDREN
CAUTION

Net Contents:
1 Gallon (3.78L) **1G**
Part No. PXC1G-US

PEROXcide Concentrate Disinfectant Cleaner

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not reuse empty container. Keep from freezing.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear chemical splash-proof goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Have the person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE. Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.

Note to Physician: Probable mucosal damage may contraindicate gastric lavage.

CLAIMS

See reference sheet for a complete list of organisms.

When used as directed, this product is highly effective against a wide variety of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, viruses, fungi, mold and mildew. At a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

Bactericidal Activity:

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica (ATCC 10708)

(formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus (ATCC 6538)

Antibiotic-Resistant Bactericidal Activity:

Escherichia coli (Extended Spectrum Beta-lactamase resistance [ESBL]), (ATCC BAA-196)

Staphylococcus aureus (MRSA) (ATCC 33591)

Staphylococcus epidermidis (MRSE) (ATCC 51625)

Fungicidal Activity:

This product is fungicidal when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F).

Microsporum canis (ATCC 10214)

Trichophyton mentagrophytes (ATCC 9533)

*Virucidal Activity:

At 1:16-1:64 dilution (2-8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

*Norovirus (Feline Calicivirus as the surrogate) (VR-782)

*Poliovirus Type 1 (ATCC VR-1000)

Kills *Influenza A Virus (H1N1) (ATCC VR-1469, Strain A/PR/8/34) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

Animal Premise *Virucidal activity

*Avian Influenza A (H3N2) (ATCC VR-2072)

*Canine Distemper virus (ATCC VR-128) (Strain Lederle)

*Equine Herpes virus type 1 (strain EHV1)

*Equine Influenza A (H3N8) virus (strain Alaska/A/EQ/AK/29759/91)

*Porcine Respiratory & Reproductive Syndrome (PRRS) virus, (Strain NVSL)

***Canine Parvovirus:** Kills *Canine Parvovirus (ATCC VR-2017) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 (five) minute contact time.

***Canine parvovirus:** Kills *Canine parvovirus (ATCC VR-2017, Strain comell) when used as directed on hard, non-porous inanimate surfaces at a 1:32 dilution (4 oz. per gallon of water), and in the presence of a 5% serum load and 10 (ten) minute contact time.

***Minute virus of mice:** Kills *Minute virus of mice (ATCC VR-1346, Strain Prototype (p) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 (five) minute contact time.

Bactericidal Stability of Use-Solution – Tests show this product when diluted at 1:16 dilution (8 oz. of product per gallon of water) in 200 ppm hard water, in the presence of 5% serum, remains effective against *Staphylococcus aureus*, *Salmonella enterica* (formerly known as *Salmonella choleraesuis*), and *Pseudomonas aeruginosa* for up to 90 days in storage as long as the bottle remains sealed. If the use-solution becomes visibly dirty or contaminated, it must be discarded and a fresh product prepared. Always use clean, dry containers when diluting this product. Treated surfaces must remain wet for 5 minutes.

Sanitizing: When used as a non-food contact sanitizer at a 1:28 dilution (1 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 3 minute contact time on hard non-porous inanimate surfaces:

Klebsiella pneumoniae (ATCC 4352)

Staphylococcus aureus (ATCC 6538)

Malodor(s) Activity – eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas (and other places where bacterial growth can cause malodors), when used as directed at a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water in 30 seconds.

Mildewcidal Activity – kills the growth of mold and mildew when diluted at 1:16 in 200 ppm hard water at a 10 minute contact time: *Aspergillus niger* (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.



PERIOD 2015-2017 LIC. NO.

8477.4

*Virucidal: 5 Min. @ 1:64

Bactericidal: 5 Min. @ 1:16

Fungicidal: 5 Min. @ 1:16

READ M.S.D.S. BEFORE USING PRODUCT
MSDS available online at www.virox.com



Manufactured by:
Virox Technologies Inc.
Oakville, ON Canada L6H 6R1
1-800-387-7578 | www.virox.com

UPC CODE
XXXXX XXXXX

PEROXcide Concentrate Disinfectant Cleaner

Animal Housing Facilities

This product is designed for use as a one-step cleaner, disinfectant, sanitizer on hard, non-porous surfaces of vehicles, trucks, trailers, carts, animal crates, grooming equipment, handling equipment, buckets, brushes, tables, dog runs, kennels, cages, transportation crates, foot baths, automatic feeders, troughs, feed racks, wash stalls, non-porous fencing, gates, gathering, guiding / controlling equipment, and non-porous harnesses & leashes; commonly found in animal housing facilities, poultry, stall, barn, pen, agricultural & livestock buildings, greenhouses, zoos, boarding kennels, veterinary clinics, animal rescue and shelter facilities, fish facilities, fish tanks, fish farms, hatcheries, tack shops, pet stores, breeding facilities, grooming facilities and catteries.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use Dilution Made Easy

Ratio	Per Gallon of Water	(Per Liter of Water)
1:16	8 oz.	64 mL
1:64	2 oz.	16 mL
1:128	1 oz.	8 mL

For Use as a One-Step Cleaner/Disinfectant: Dilute at 8.0 oz. of product per gallon of water (1:16)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution by coarse trigger sprayer to hard, non-porous environmental surfaces. Spray 6-8 inches from surface; making sure to wet surfaces thoroughly.
3. All surfaces must remain wet for 5 minutes.
4. Wipe surfaces and let air dry.

For use as a *Virucide: Dilute at 2 oz. – 8 oz. of product per gallon of water (1:16 -1:64)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 5 minutes.
4. Wipe surfaces and let air dry.

For Treatment of Animal Housing Facilities (including Poultry barns): Dilute at 8.0 oz. of product per gallon of water (1:16)

1. Remove all animals and feeds from areas being treated.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
3. Empty or cover all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Apply fresh Use Solution to floors, walls, cages and other washable hard, non-porous environmental surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain wet for 5 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or re-employ equipment until product has dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and watering appliances scrub with use-solution, let stand 5 minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.

To Sanitize Non-Food Contact Surfaces: Dilute at 1.0 oz. of product per gallon of water (1:128)

1. Pre-clean heavily soiled hard non-porous surfaces.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 3 minutes.
4. Wipe surfaces and let air dry.
5. Not for use on food contact surfaces or on food preparation areas.

To Kill *Minute virus of mice: Dilute at 8.0 oz. of product per gallon of water (1:16)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 5 minutes.
4. Wipe surfaces and let air dry.

To Kill *Canine Parvovirus: Dilute at 4.0 oz. of product per gallon of water (1:32)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 10 minutes.
4. Wipe surfaces and let air dry.

To Kill Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16). Pre-clean surfaces. Apply Use Solution to hard, non-porous environmental surfaces. Allow surfaces to remain wet for 10 minutes. Wipe surfaces and let air dry.

For Use as a Deodorizer: Dilute at 8 oz. of product per gallon of water (1:16). Apply to hard, nonporous surfaces. Let stand for 30 seconds.

To Kill Fungi: Dilute at 8.0 oz. of product per gallon of water (1:16)

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for 5 minutes. Wipe surfaces and let air dry. Repeat application weekly or when growth reappears.

For Use as a Cleaner/Degreaser: Dilute product at 0.5 - 1.0 oz. of product per gallon of water (1:128 – 1:256). Apply to surfaces. Wipe surfaces and let air dry.

PEROXcide

Concentrate Disinfectant Cleaner

Multi-Surface One Step Disinfectant Cleaner and Deodorizer

ACTIVE INGREDIENT
Hydrogen Peroxide 4.25%
INERT INGREDIENTS 95.75%
Total 100.00%

EPA Reg. No. 74559-4
EPA Est. No. 74559-CAN-1

UNSCENTED

KEEP OUT OF REACH OF CHILDREN
CAUTION



See Additional Precautionary Statements on Side Panel
See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use.

Net Contents: **5G**
5 Gallon (18.9L)
Part No. PX05G-US

PEROXcide Concentrate Disinfectant Cleaner

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not reuse empty container. Keep from freezing.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting agency has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear chemical splash-proof goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Have the person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE. Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.

Note to Physician: Probable mucosal damage may contraindicate gastric lavage.

CLAIMS

See reference sheet for a complete list of organisms.

When used as directed, this product is highly effective against a wide variety of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, viruses, fungi, mold and mildew. At a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

Bactericidal Activity:

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica (ATCC 10708)

(formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus (ATCC 6538)

Antibiotic-Resistant Bactericidal Activity:

Escherichia coli (Extended Spectrum Beta-lactamase resistance [ESBL]), (ATCC BAA-196)

Staphylococcus aureus (MRSA) (ATCC 33591)

Staphylococcus epidermidis (MRSE) (ATCC 51625)

Fungicidal Activity:

This product is fungicidal when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F).

Microsporum canis (ATCC 10214)

Trichophyton mentagrophytes (ATCC 9533)

*Virucidal Activity:

At 1:16-1:64 dilution (2-8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

*Norovirus (Feline Calicivirus as the surrogate) (VR-782)

*Poliovirus Type 1 (ATCC VR-1000)

Kills *Influenza A Virus (H1N1) (ATCC VR-1469, Strain A/PR/8/34) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

Animal Premise *Virucidal activity

*Avian Influenza A (H3N2) (ATCC VR-2072)

*Canine Distemper virus (ATCC VR-128), (Strain Lederle)

*Equine Herpes virus type 1 (Strain EHV1)

*Equine Influenza A (H3N8) virus (Strain Alaska/A/EQ/AK/29759/91)

*Porcine Respiratory & Reproductive Syndrome (PRRS) virus, (Strain NVSL)

***Canine Parvovirus:** Kills *Canine Parvovirus (ATCC VR-2017) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 (five) minute contact time.

***Canine parvovirus:** Kills *Canine parvovirus (ATCC VR-2017, Strain cornell) when used as directed on hard, non-porous inanimate surfaces at a 1:32 dilution (4 oz. per gallon of water), and in the presence of a 5% serum load and 10 (ten) minute contact time.

***Minute virus of mice:** Kills *Minute virus of mice (ATCC VR-1346, Strain Prototype (p)) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 (five) minute contact time.

Bactericidal Stability of Use-Solution – Tests show this product when diluted at 1:16 dilution (8 oz. of product per gallon of water) in 200 ppm hard water, in the presence of 5% serum, remains effective against *Staphylococcus aureus*, *Salmonella enterica* (formerly known as *Salmonella choleraesuis*) and *Pseudomonas aeruginosa* for up to 90 days in storage as long as the bottle remains sealed. If the use-solution becomes visibly dirty or contaminated, it must be discarded and a fresh product prepared. Always use clean, dry containers when diluting this product. Treated surfaces must remain wet for 5 minutes.

Sanitizing: When used as a non-food contact sanitizer at a 1:128 dilution (1 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 3 minute contact time on hard non-porous inanimate surfaces:

Klebsiella pneumoniae (ATCC 4352)

Staphylococcus aureus (ATCC 6538)

Malodor(s) Activity – eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas (and other places where bacterial growth can cause malodors), when used as directed at a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water in 30 seconds.

Mildewicidal Activity – kills the growth of mold and mildew: when diluted at 1:16 in 200 ppm hard water at a 10 minute contact time: *Aspergillus niger* (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.



PERIOD **2015-2017** LIC. NO.

8477.4

*Virucidal: 5 Min. @ 1:64

Bactericidal: 5 Min. @ 1:16

Fungicidal: 5 Min. @ 1:16

READ M.S.D.S. BEFORE USING PRODUCT
MSDS available online at www.virox.com



Manufactured by:
Virox Technologies Inc.
Oakville, ON Canada L6H 6R1
1-800-387-7578 | www.virox.com

UPC CODE
XXXXX XXXXX

PEROXcide Concentrate Disinfectant Cleaner

Animal Housing Facilities

This product is designed for use as a one-step cleaner, disinfectant, sanitizer on hard, non-porous surfaces of vehicles, trucks, trailers, carts, animal crates, grooming equipment, handling equipment, buckets, brushes, tables, dog runs, kennels, cages, transportation crates, foot baths, automatic feeders, troughs, feed racks, wash stalls, non-porous fencing, gates, gathering, guiding / controlling equipment, and non-porous harnesses & leashes; commonly found in animal housing facilities, poultry, stall, barn, pen, agricultural & livestock buildings, greenhouses, zoos, boarding kennels, veterinary clinics, animal rescue and shelter facilities, fish facilities, fish tanks, fish farms, hatcheries, tack shops, pet stores, breeding facilities, grooming facilities and catteries.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use Dilution Made Easy

Ratio	Per Gallon of Water	(Per Liter of Water)
1:16	8 oz.	64 mL
1:64	2 oz.	16 mL
1:128	1 oz.	8 mL

For Use as a One-Step Cleaner/Disinfectant: Dilute at 8.0 oz. of product per gallon of water (1:16)

- Pre-clean heavily soiled areas.
- Apply Use Solution by coarse trigger sprayer to hard, non-porous environmental surfaces. Spray 6-8 inches from surface; making sure to wet surfaces thoroughly.
- All surfaces must remain wet for 5 minutes.
- Wipe surfaces and let air dry.

For use as a *Virucide: Dilute at 2 oz. – 8 oz. of product per gallon of water (1:16 -1:64)

- Pre-clean heavily soiled areas.
- Apply Use Solution until thoroughly wet.
- Let stand for 5 minutes.
- Wipe surfaces and let air dry.

For Treatment of Animal Housing Facilities (including Poultry barns): Dilute at 8.0 oz. of product per gallon of water (1:16)

- Remove all animals and feeds from areas being treated.
- Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
- Empty or cover all troughs, racks and other feeding and watering appliances.
- Thoroughly clean all surfaces with soap or detergent and rinse with water.
- Apply fresh Use Solution to floors, walls, cages and other washable hard, non-porous environmental surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain wet for 5 minutes.
- Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
- Ventilate buildings and other closed spaces. Do not house animals or re-employ equipment until product has dried.
- For disinfection of feed racks, troughs, automatic feeders, fountains and watering appliances scrub with use-solution, let stand 5 minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.

To Sanitize Non-Food Contact Surfaces: Dilute at 1.0 oz. of product per gallon of water (1:128)

- Pre-clean heavily soiled hard non-porous surfaces.
- Apply Use Solution until thoroughly wet.
- Let stand for 3 minutes.
- Wipe surfaces and let air dry.
- Not for use on food contact surfaces or on food preparation areas.

To Kill *Minute virus of mice: Dilute at 8.0 oz. of product per gallon of water (1:16)

- Pre-clean heavily soiled areas.
- Apply Use Solution until thoroughly wet.
- Let stand for 5 minutes.
- Wipe surfaces and let air dry.

To Kill *Canine Parvovirus: Dilute at 4.0 oz. of product per gallon of water (1:32)

- Pre-clean heavily soiled areas.
- Apply Use Solution until thoroughly wet.
- Let stand for 10 minutes.
- Wipe surfaces and let air dry.

To Kill Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16). Pre-clean surfaces. Apply Use Solution to hard, non-porous environmental surfaces. Allow surfaces to remain wet for 10 minutes. Wipe surfaces and let air dry.

For Use as a Deodorizer: Dilute at 8 oz. of product per gallon of water (1:16). Apply to hard, nonporous surfaces. Let stand for 30 seconds.

To Kill Fungi: Dilute at 8.0 oz. of product per gallon of water (1:16)

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for 5 minutes. Wipe surfaces and let air dry.

To Control Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16).

Apply Use Solution to pre-cleaned hard, non-porous environmental surfaces. Wipe surfaces and let air dry. Repeat application weekly or when growth reappears.

For Use as a Cleaner/Degreaser: Dilute product at 0.5 - 1.0 oz. of product per gallon of water (1:128 – 1:256). Apply to surfaces. Wipe surfaces.

PEROXcide

Concentrate Disinfectant Cleaner

Multi-Surface One Step Disinfectant Cleaner and Deodorizer

ACTIVE INGREDIENT
Hydrogen Peroxide 4.25%
INERT INGREDIENTS 95.75%
Total 100.00%

EPA Reg. No. 74559-4
EPA Est. No. 74559-CAN-1

UNSCENTED

KEEP OUT OF REACH OF CHILDREN
CAUTION



See Additional Precautionary Statements on Side Panel
See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use.

Net Contents:
30 Gallon (113.5L) **30G**
Part No. PXC30G-US

PEROXcide Concentrate Disinfectant Cleaner

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not reuse empty container. Keep from freezing.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for at least 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for reconditioning, if appropriate.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting agency has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear chemical splash-proof goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Have the person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE. Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.

Note to Physician: Probable mucosal damage may contraindicate gastric lavage.

CLAIMS

See reference sheet for a complete list of organisms.

When used as directed, this product is highly effective against a wide variety of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, viruses, fungi, mold and mildew. At a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

Bactericidal Activity:

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica (ATCC 10708)

(formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus (ATCC 6538)

Antibiotic-Resistant Bactericidal Activity:

Escherichia coli (Extended Spectrum Beta-lactamase resistance [ESBL]), (ATCC BAA-196)

Staphylococcus aureus (MRSA) (ATCC 33591)

Staphylococcus epidermidis (MRSE) (ATCC 51625)

Fungicidal Activity:

This product is fungicidal when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F).

Microsporium canis (ATCC 10214)

Trichophyton mentagrophytes (ATCC 9533)

*Virucidal Activity:

At 1:16-1:64 dilution (2-8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

*Norovirus (Feline Calicivirus as the surrogate) (VR-782)

*Poliovirus Type 1 (ATCC VR-1000)

Kills *Influenza A Virus (H1N1) (ATCC VR-1469, Strain A/PR/8/34) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

Animal Premise *Virucidal activity

*Avian Influenza A (H3N2) (ATCC VR-2072)

*Canine Distemper virus (ATCC VR-128), (Strain Lederle)

*Equine Herpes virus type 1 (Strain EHV1)

*Equine Influenza A (H3N8) virus (Strain Alaska/A/EQ/AK/29759/91)

*Porcine Respiratory & Reproductive Syndrome (PRRS) virus, (Strain NVSL)

***Canine Parvovirus:** Kills *Canine Parvovirus (ATCC VR-2017) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 (five) minute contact time.

***Canine parvovirus:** Kills *Canine parvovirus (ATCC VR-2017, Strain cornell) when used as directed on hard, non-porous inanimate surfaces at a 1:32 dilution (4 oz. per gallon of water), and in the presence of a 5% serum load and 10 (ten) minute contact time.

***Minute virus of mice:** Kills *Minute virus of mice (ATCC VR-1346, Strain Prototype (p)) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 (five) minute contact time.

Bactericidal Stability of Use-Solution – Tests show this product when diluted at 1:16 dilution (8 oz. of product per gallon of water) in 200 ppm hard water, in the presence of 5% serum, remains effective against *Staphylococcus aureus*, *Salmonella enterica* (formerly known as *Salmonella choleraesuis*) and *Pseudomonas aeruginosa* for up to 90 days in storage as long as the bottle remains sealed. If the use-solution becomes visibly dirty or contaminated, it must be discarded and a fresh product prepared. Always use clean, dry containers when diluting this product. Treated surfaces must remain wet for 5 minutes.

Sanitizing: When used as a non-food contact sanitizer at a 1:128 dilution (1 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 3 minute contact time on hard non-porous inanimate surfaces:

Klebsiella pneumoniae (ATCC 4352)

Staphylococcus aureus (ATCC 6538)

Malodor(s) Activity – eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas (and other places where bacterial growth can cause malodors), when used as directed at a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water in 30 seconds.

Mildewicidal Activity – kills the growth of mold and mildew: when diluted at 1:16 in 200 ppm hard water at a 10 minute contact time: *Aspergillus niger* (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.



PERIOD **2015-2017** LIC. NO.

8477.4

*Virucidal: 5 Min. @ 1:64

Bactericidal: 5 Min. @ 1:16

Fungicidal: 5 Min. @ 1:16

READ M.S.D.S. BEFORE USING PRODUCT
MSDS available online at www.virox.com



Manufactured by:

Virox Technologies Inc.
Oakville, ON Canada L6H 6R1
1-800-387-7578 | www.virox.com

UPC CODE
XXXXX XXXXX

XXXXXX (XX/XXX)

PEROXcide Concentrate Disinfectant Cleaner

Animal Housing Facilities

This product is designed for use as a one-step cleaner, disinfectant, sanitizer on hard, non-porous surfaces of vehicles, trucks, trailers, carts, animal crates, grooming equipment, handling equipment, buckets, brushes, tables, dog runs, kennels, cages, transportation crates, foot baths, automatic feeders, troughs, feed racks, wash stalls, non-porous fencing, gates, gathering, guiding / controlling equipment, and non-porous harnesses & leashes; commonly found in animal housing facilities, poultry, stall, barn, pen, agricultural & livestock buildings, greenhouses, zoos, boarding kennels, veterinary clinics, animal rescue and shelter facilities, fish facilities, fish tanks, fish farms, hatcheries, tack shops, pet stores, breeding facilities, grooming facilities and catteries.

For use as a *Virucide: Dilute at 2 oz. – 8 oz. of product per gallon of water (1:16 -1:64)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 5 minutes.
4. Wipe surfaces and let air dry.

For Treatment of Animal Housing Facilities (including Poultry barns): Dilute at 8.0 oz. of product per gallon of water (1:16)

1. Remove all animals and feeds from areas being treated.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
3. Empty or cover all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Apply fresh Use Solution to floors, walls, cages and other washable hard, non-porous environmental surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain wet for 5 minutes.

6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

7. Ventilate buildings and other closed spaces. Do not house animals or re-employ equipment until product has dried.

8. For disinfection of feed racks, troughs, automatic feeders, fountains and watering appliances scrub with use-solution, let stand 5 minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.

To Sanitize Non-Food Contact Surfaces: Dilute at 1.0 oz. of product per gallon of water (1:128)

1. Pre-clean heavily soiled hard non-porous surfaces.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 3 minutes.
4. Wipe surfaces and let air dry.
5. Not for use on food contact surfaces or on food preparation areas.

To Kill *Minute virus of mice: Dilute at 8.0 oz. of product per gallon of water (1:16)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 5 minutes.
4. Wipe surfaces and let air dry.

To Kill *Canine Parvovirus: Dilute at 4.0 oz. of product per gallon of water (1:32)

1. Pre-clean heavily soiled areas.
2. Apply Use Solution until thoroughly wet.
3. Let stand for 10 minutes.
4. Wipe surfaces and let air dry.

To Kill Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16). Pre-clean surfaces. Apply Use Solution to hard, non-porous environmental surfaces. Allow surfaces to remain wet for 10 minutes. Wipe surfaces and let air dry.

For Use as a Deodorizer: Dilute at 8 oz. of product per gallon of water (1:16). Apply to hard, nonporous surfaces. Let stand for 30 seconds.

To Kill Fungi: Dilute at 8.0 oz. of product per gallon of water (1:16)

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for 5 minutes. Wipe surfaces and let air dry.

To Control Mold and Mildew: Dilute at 8.0 oz. of product per gallon of water (1:16).

Apply Use Solution to pre-cleaned hard, non-porous environmental surfaces. Wipe surfaces and let air dry. Repeat application weekly or when growth reappears.

For Use as a Cleaner/Degreaser: Dilute product at 0.5 - 1.0 oz. of product per gallon of water (1:128 – 1:256). Apply to surfaces. Wipe surfaces.

PEROXcide

Concentrate Disinfectant Cleaner

Multi-Surface One Step Disinfectant Cleaner and Deodorizer

ACTIVE INGREDIENT
Hydrogen Peroxide 4.25%
INERT INGREDIENTS 95.75%
Total 100.00%

EPA Reg. No. 74559-4
EPA Est. No. 74559-CAN-1

UNSCENTED

KEEP OUT OF REACH OF CHILDREN
CAUTION



See Additional Precautionary Statements on Side Panel
See reference sheet for a complete list of pathogenic organisms, additional features, claims and directions for use.

Net Contents:
55 Gallon (208.1L) **55G**
Part No. PXC55G-US

PEROXcide Concentrate Disinfectant Cleaner

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not reuse empty container. Keep from freezing.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for at least 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for reconditioning, if appropriate.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting agency has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear chemical splash-proof goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Have the person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE. Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.

Note to Physician: Probable mucosal damage may contraindicate gastric lavage.

CLAIMS

See reference sheet for a complete list of organisms.

When used as directed, this product is highly effective against a wide variety of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, viruses, fungi, mold and mildew. At a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

Bactericidal Activity:

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica (ATCC 10708)

(formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus (ATCC 6538)

Antibiotic-Resistant Bactericidal Activity:

Escherichia coli (Extended Spectrum Beta-lactamase resistance [ESBL]), (ATCC BAA-196)

Staphylococcus aureus (MRSA) (ATCC 33591)

Staphylococcus epidermidis (MRSE) (ATCC 51625)

Fungicidal Activity:

This product is fungicidal when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F).

Microsporium canis (ATCC 10214)

Trichophyton mentagrophytes (ATCC 9533)

*Virucidal Activity:

At 1:16-1:64 dilution (2-8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

*Norovirus (Feline Calicivirus as the surrogate) (VR-782)

*Poliovirus Type 1 (ATCC VR-1000)

Kills *Influenza A Virus (H1N1) (ATCC VR-1469, Strain A/PR/8/34) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

Animal Premise *Virucidal activity

*Avian Influenza A (H3N2) (ATCC VR-2072)

*Canine Distemper virus (ATCC VR-128), (Strain Lederle)

*Equine Herpes virus type 1 (strain EHV1)

*Equine Influenza A (H3N8) virus (strain Alaska/A/EQ/AK/29759/91)

*Porcine Respiratory & Reproductive Syndrome (PRRS) virus, (Strain NVSL)

***Canine Parvovirus:** Kills *Canine Parvovirus (ATCC VR-2017) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 (five) minute contact time.

***Canine parvovirus:** Kills *Canine parvovirus (ATCC VR-2017, Strain cornell) when used as directed on hard, non-porous inanimate surfaces at a 1:32 dilution (4 oz. per gallon of water), and in the presence of a 5% serum load and 10 (ten) minute contact time.

***Minute virus of mice:** Kills *Minute virus of mice (ATCC VR-1346, Strain Prototype (p) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 (five) minute contact time.

Bactericidal Stability of Use-Solution – Tests show this product when diluted at 1:16 dilution (8 oz. of product per gallon of water) in 200 ppm hard water, in the presence of 5% serum, remains effective against *Staphylococcus aureus*, *Salmonella enterica* (formerly known as *Salmonella choleraesuis*) and *Pseudomonas aeruginosa* for up to 90 days in storage as long as the bottle remains sealed. If the use-solution becomes visibly dirty or contaminated, it must be discarded and a fresh product prepared. Always use clean, dry containers when diluting this product. Treated surfaces must remain wet for 5 minutes.

Sanitizing: When used as a non-food contact sanitizer at a 1:128 dilution (1 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 3 minute contact time on hard non-porous inanimate surfaces:

Klebsiella pneumoniae (ATCC 4352)

Staphylococcus aureus (ATCC 6538)

Malodor(s) Activity – eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas (and other places where bacterial growth can cause malodors), when used as directed at a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water in 30 seconds.

Mildewicidal Activity – kills the growth of mold and mildew: when diluted at 1:16 in 200 ppm hard water at a 10 minute contact time: *Aspergillus niger* (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.



PERIOD **2015-2017** LIC. NO.

8477.4

*Virucidal: 5 Min. @ 1:64

Bactericidal: 5 Min. @ 1:16

Fungicidal: 5 Min. @ 1:16

READ M.S.D.S. BEFORE USING PRODUCT
MSDS available online at www.virox.com



Manufactured by:

Virox Technologies Inc.
Oakville, ON Canada L6H 6R1
1-800-387-7578 | www.virox.com

UPC CODE
XXXXXX XXXXX

XXXXXX (XX/XXX)

PEROXCide

CONCENTRATE DISINFECTANT CLEANER

Manufactured by:
Virox Technologies Inc.
 Oakville, ON Canada L6H 6R1
 1-800-387-7578 | www.virox.com



Reference Sheet

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms (including bacteria, antibiotic-resistant bacteria, *viruses, fungi, mold and mildew.) At a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard non-porous inanimate surfaces:

USE DILUTION MADE EASY

Ratio	Per Gallon of Water	(Per Liter of Water)
1:16	8 oz.	64 mL
1:64	2 oz	16 mL
1:128	1 oz.	8 mL
1:256	0.5 oz.	4 mL

BACTERICIDAL: 5 MINUTES @ 1:16

- Acinetobacter baumannii*, (ATCC 19606)
- Escherichia coli* O157:H7 (ATCC 35150)
- Klebsiella pneumoniae* (ATCC 4352)
- Listeria monocytogenes* (ATCC 19111)
- Pseudomonas aeruginosa* (ATCC 15442)
- Salmonella enterica*, (ATCC 10708)
(formerly known as *Salmonella choleraesuis*)
- Shigella dysenteriae* (ATCC 13313)
- Staphylococcus aureus* (ATCC 6538)
- Streptococcus pyogenes* (ATCC 12204)

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY

- Enterococcus faecium*, (Resistant to Vancomycin [VRE]) (ATCC 51559)
- Escherichia coli*, (Extended Spectrum Beta-lactamase resistance [ESBL]), (ATCC BAA-196)
- Klebsiella pneumoniae*, (Carbapenem-resistant [KPC]), (ATCC BAA-1705)
- Staphylococcus aureus*, (CA-MRSA) (clinical Isolate 08001)
(Genotype USA300)
- Staphylococcus aureus*, (CA-MRSA) (clinical Isolate 08005)
(Genotype USA400)
- Staphylococcus aureus* (MRSA) (ATCC 33591)
- Staphylococcus epidermidis* (MRSE) (ATCC 51625)
- Streptococcus pneumoniae*, Penicillin Resistant (PRSP), (ATCC 700671)

***VIRUCIDAL: 5 MINUTES @ 1:64**

At 1:16 - 1:64 dilution (2 - 8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load and 5 minute contact time, unless otherwise noted, this product kills the following on hard, non-porous inanimate surfaces:

- *Adenovirus type 8, (ATCC VR-1368, Strain Trim)
- *Herpes simplex Type 2 (VR-734)
- *Human Coronavirus (VR-740)
- *Influenza Type A₂ (Hong Kong) (VR-544)
- *Norovirus (Feline Calicivirus as the surrogate) (VR-782)
- *Parainfluenza Type 3 (VR-93)
- *Poliovirus Type 1 (ATCC VR-1000)
- *Respiratory Syncytial Virus (VR-26)
- *Rhinovirus Type 37, Strain 151-1, (ATCC VR-1147)
- *Rotavirus (Strain WA)
- *Vaccinia Virus (smallpox vaccine virus) (ATCC VR-119)

Kills *Hepatitis B Virus (HBV) & *Hepatitis C Virus (HCV) when used as directed on hard, non-porous inanimate surfaces with a 5 minute contact time.

Kills *HIV-1 (AIDS virus) (HTLV-III_B) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

*This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

***VETERINARY VIRUSES**

- *Avian Influenza A, (H3N2) (ATCC VR-2072)
- *Canine Distemper virus, (ATCC VR-128), (Strain Lederle)
- *Equine Herpes virus type 1, (strain EHV1)
- *Equine Influenza A (H3N8) virus, (strain Alaska/A/EQ/AK/29759/91)
- *Porcine Respiratory & Reproductive Syndrome (PRRS) virus, (Strain NVSL)
- *Canine parvovirus: Kills *Canine parvovirus (ATCC VR-2017) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time.
- *Canine parvovirus: Kills *Canine parvovirus (ATCC VR-2017, Strain cornell) when used as directed on hard, non-porous inanimate surfaces at a 1:32 dilution (4 oz. per gallon of water), and in the presence of a 5% serum load and 10 minute contact time.
- *Minute virus of mice: Kills *Minute virus of mice (ATCC VR-1346, Strain Prototype (p) when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time.

FUNGICIDAL: 5 MINUTES @ 1:16

This product is fungicidal when used as directed on hard, non-porous inanimate surfaces at a 1:16 dilution (8 oz. per gallon of water), and in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F).

Trichophyton mentagrophytes (ATCC 9533)

Microsporium canis (ATCC 10214)

MALODORS

Malodors – reduces odors and odor-causing bacteria on hard non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors, when used as directed at a 1:16 dilution (8 oz. of product per gallon of water, in the presence of 200 ppm hard water and a 30 seconds contact time.

MOLD/MILDEW

Mildewcidal Activity: Kills the growth of mold and mildew when diluted at 1:16 in 200 ppm hard water at a 10 minute contact time: *Aspergillus niger* (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mildewstatic Activity: Controls and prevents the growth of mold and mildew when diluted at 1:16 in 200 ppm hard water at a 10 minute contact time: *Aspergillus niger* (ATCC 6275) and the odors caused by them when applied to hard, non-porous environmental surfaces.

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 30 SECONDS @ 1:16

Sanitizing: When used as a non-food contact sanitizer at a 1:16 dilution (8 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 30 second contact time on hard non-porous inanimate surfaces:

Enterobacter aerogenes (ATCC 13048)

Enterococcus faecalis, *Vancomycin-Resistant* (VRE), (ATCC 51575)

Escherichia coli, *Extended Spectrum Beta-lactamase resistance* (ESBL), (ATCC BAA-196)

Listeria monocytogenes, (ATCC 19111)

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica (ATCC 10708) (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus (ATCC 6538)

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 3 MINUTES @ 1:128

Sanitizing: When used as a non-food contact sanitizer at a 1:128 dilution (1 oz. of product per gallon of water), in the presence of 200 ppm hard water, 5% serum load, this product reduces the following bacteria by 99.9% with a 3 minute contact time on hard non-porous inanimate surfaces:

Escherichia coli (ATCC 11229)

Enterococcus faecalis (ATCC 29212)

Klebsiella pneumoniae (ATCC 4352)

Listeria monocytogenes (ATCC 19111)

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica, (ATCC 10708) (formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus (ATCC 6538)

BACTERICIDAL STABILITY OF USE-SOLUTION

Tests show this product as a disinfectant, when diluted at 1:16 dilution (8 oz. of product per gallon of water) in 200 ppm hard water, in the presence of 5% serum, remains effective against *Staphylococcus aureus*, *Salmonella enterica* (formerly known as *Salmonella choleraesuis*) and *Pseudomonas aeruginosa* for up to 90 days in storage as long as the bottle remains sealed.

If the use-solution becomes visibly dirty or contaminated, it must be discarded and a fresh product prepared. Always use clean, dry containers when diluting this product. Treated surfaces must remain wet for 5 minutes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

*This product kills HBV, HCV and HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Hepatitis B Virus (HBV), Hepatitis C Virus (HCV) and Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HBV, HCV and HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, and for 5 minutes to kill HBV, HCV, fungi and all other organisms cited on the label.

Disposal of Infectious Material: Blood and other body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.