



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD **2006-2008** LIC. NO.

———— BLEED LINE

———— DIE LINE



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD **2009-2011** LIC. NO.

**DISCONTINUED**



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD **2012-2014** LIC. NO.



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD **2015-2016** LIC. NO.

**8485.13**

**Do it  
Best**

**LAWN WEED  
CONTROL**

**Concentrate**  
**Control para Césped**  
**y Mala Hierba**  
**Concentrado**



kills: chickweed, clover,  
dandelion, ground ivy,  
plantain, poison ivy,  
poison oak, and other  
broadleaf weeds

- kills weeds **NOT** the lawn
- starts working **overnight**
- controls **over 200** weeds
- rain-proof in **hours!**

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid .....	5.56%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid .....	1.34%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid .....	0.62%

INERT INGREDIENTS .....	92.48%
TOTAL .....	100.00%

THIS PRODUCT CONTAINS:

4.62% 2,4-dichlorophenoxyacetic acid equivalent.  
1.11% (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent.  
0.51% 3,6-dichloro-o-anisic acid equivalent.

Isomer Specific by AOAC Method.  
Contains the single isomer  
form of Mecoprop-p.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION** See booklet back panel for first aid and  
additional precautionary statements

728034 O/M 1-12 NET CONTENTS 1QT (32 FL OZ) 946ml

# 32 OZ LAWN WEED CONTROL CONCENTRATE Front Label

EPA Reg No. 2217-540-75111

**PRINTS 4 COLOR PROCESS**

RED = PMS 485 (C-0, M-100, Y-90, K-0)

YELLOW = PMS 115 (C-0, M-9, Y-80, K-0)

**OTHER WEEDS CONTROLLED (continued):**

Orange hawkweed	Purslane, common	Spotted catsear	Wild buckwheat
Oriental cocklebur	Ragweed	Spotted spurge	Wild carrot
Oxalis	Red sorrel	Spurweed	Wild four-o'clock
Parsley-piert	Redroot pigweed	Stinging nettle	Wild garlic
Parsnip	Redstem filaree	Strawberry, India mock	Wild geranium
Pearwort	Rough cinquefoil	Tall nettle	Wild lettuce
Pennycress	Rough fleabane	Tall vervain	Wild marigold
Pepperweed	Russian pigweed	Tansy ragwort	Wild mustard
Pigweed	Russian thistle	Tansy mustard	Wild onion
Pineywoods bedstraw	Scarlet pimpernel	Tanweed	Wild parsnip
Plains coreopsis	Scotch thistle	Thistle	Wild radish
Plantain	Sheep sorrel	Trailing crownvetch	Wild rape
Poison ivy	Shepherdspurse	Tumble mustard	Wild strawberry
Poison oak	Slender plantain	Tumble pigweed	Wild sweet potato
Pokeweed	Smallflower galinsoga	Velvetleaf	Wild vetch
Poorjoe	Smooth dock	Venice mallow	Woodsorrel
Prairie sunflower	Smooth pigweed	Virginia buttonweed	Woolly croton
Prickly lettuce	Sorrel	Virginia creeper	Woolly morningglory
Prickly sida	Sowthistle	Virginia pepperweed	Woolly plantain
Prostrate knotweed	Spanishneedles	Wavyleaf bulthistle	Wormseed
Prostrate pigweed	Speedwell	Western clematis	Yarrow
Prostrate spurge	Spiny amaranth	Western salsify	Yellow rocket
Prostrate vervain	Spiny cocklebur	White mustard	Yellowflower pepperweed
Puncturevine	Spiny sowthistle	Wild aster	and other broadleaf weeds

**LIMITED WARRANTY AND DISCLAIMER.**

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



**OPEN HERE**

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

**WHERE TO USE:**

This product may be used on residential and ornamental lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. Established turf of bahiagrass, bermudagrass, Kentucky bluegrass, centipede-grass, fescues, perennial ryegrass, St. Augustinegrass, and zoysiagrass may be treated. This product is especially effective on all hard-to-kill weeds such as poison ivy, poison oak, sheep sorrel and spurge. In addition, this product controls most other broadleaf weeds, including sensitive weeds such as chickweed, clover, dandelion, knotweed and plantain. A partial list is given on a following panel.

EPA Reg. No. 2217-540-75111  
EPA Est. No. 58996-M0-1

Distributed by:  
Maid Brands, Inc.  
Hazelwood, MO 63042

Do it Best® is a trademark of Do it Best Corp., Ft. Wayne, IN 46801.



**OTHER USE PRECAUTIONS:** Do not allow drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not pour spray solutions near these plants.

Whether using a hose end sprayer or pressure sprayer, do not use sprays that are fine because they can drift and cause damage to nontarget plants. Coarse droplet sprays are less likely to drift. Use only lawn-type sprayers.

Do not spray on carpetgrass, dichondra, Lippia, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals.

Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species.

This product may be injurious to St. Augustinegrass and centipede-grass if misapplications occur. Do not apply this product to 'Floratam' St. Augustinegrass in Florida.

The maximum application rate to turf is 2.0 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

**WHEN TO USE:**

1. Spray at any time during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks after treatment before seeding grass seed or planting sprigs or stolons.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow before treatment.
5. Do not water lawn within 24 hours after treatment.

**HOSE END SPRAYERS**

**HOW TO APPLY:** Use a hose end sprayer designed for home lawn care. Refer to Charts 1 and 2 for the proper amounts of this product. Determine the area of turf to be treated. Pour the recommended amount of product (Column 1) into the bottle, add water to the indicated level (Column 2), and spray uniformly on turf area shown in Column 3.

CHART 1: Amounts of this product for bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns applied with hose end sprayers.			
Column 1		Column 2	Column 3
Amount of this product to Use		Fill Bottle to, Gallons	Area to Treat, Square Feet
(Tablespoons)	(Fluid Ounces)		
10 to 12	5 to 6	5	950
20 to 24	10 to 12	10	1,900
30 to 36	15 to 18	15	2,850
40 to 48	20 to 24	20	3,800

**Note:** For these grasses apply the entire contents of this container (32 fluid ounces) on 5,000 to 6,000 square feet of turf.

CHART 2: Amounts of this product for centipede-grass and St. Augustinegrass lawns applied with hose end sprayers			
Column 1		Column 2	Column 3
Amount of this product to Use		Fill Bottle to, Gallons	Area to Treat, Square Feet
(Tablespoons)	(Fluid Ounces)		
6	3	5	950
12	6	10	1,900
18	9	15	2,850
24	12	20	3,800

**Note:** For St. Augustinegrass and centipede-grass apply the entire contents of this container (32 fluid ounces) on 10,000 square feet of turf.

**PRESSURE SPRAYERS, BACKPACK SPRAYERS, OR OTHER MANUALLY OPERATED SPRAYERS:** Prepare the appropriate amounts of spray mixtures according to Chart 3. Measure the recommended amount of this product into the spray tank and dilute with the proper amount of water. Use a spray pattern with small droplets and treat the area indicated in Chart 3.

CHART 3A: For bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns.			
Amount of this product to Use		Amount of Water in Sprayer,	Area to Treat,
(Tablespoons)	(Fluid Ounces)	Gallons	Square Feet
2 to 2.4	1 to 1.2	¾	190
4 to 4.8	2 to 2.4	1½	380
8 to 9.6	4 to 4.8	3	760

CHART 3B: For centipede-grass and St. Augustinegrass lawns.			
Amount of this product to Use		Amount of Water in Sprayer,	Area to Treat,
(Tablespoons)	(Fluid Ounces)	Gallons	Square Feet
2 to 2.4	1 to 1.2	¾	380
4 to 4.8	2 to 2.4	1½	760
8 to 9.6	4 to 4.8	3	1,520

**SPOT SPRAYING:** To spray individual weeds, do not use a hose end sprayer. Mix 1 fluid ounce (2 Tablespoons) of this product in ½ gallon of water in a pressure sprayer. Spray to wet individual weeds.

**GENERAL BRUSH CONTROL:**

For control of brush along roads and other similar noncrop areas, mix 10 to 14 fluid ounces of this product in 3 gallons of water and apply over 1,000 square feet. Use the higher rate of product for larger plants or more dense brush stands. Refer to Table 1 for additional instructions for spray preparations with water. Best results are obtained when brush is young and actively growing. Do not cut brush until the herbicide has translocated throughout the plant causing root death. Keep out of lakes, streams and ponds.

Table 1. Mixing instructions for preparing 1 to 9 gallons of spray solution with water for foliar application.		
Spray Solution Desired, Gallons	Amount of this product Required	
	Normal Rate	High Rate (For larger plants or more dense brush)
0.25 (32 fl. oz.)	0.83 fl. oz.	1.16 fl. oz.
1	3 1/3 fl. oz.	4 2/3 fl. oz.
2	6 2/3 fl. oz.	9 1/3 fl. oz.
3	10 fl. oz.	14 fl. oz.
4	13 1/3 fl. oz.	18 2/3 fl. oz.
5	16 2/3 fl. oz.	23 1/3 fl. oz.
6	20 fl. oz.	28 fl. oz.
7	23 1/3 fl. oz.	32 2/3 fl. oz.
8	26 2/3 fl. oz.	37 1/3 fl. oz.
9	30 fl. oz.	42 fl. oz.

Equal measures: 1.0 fl. oz. = 2 tablespoons of product  
1.0 quart = 32 fl. oz. = 2 pints = 4 cups

**CONTROLS:**

**BRUSH AND VINES:**

Alder	Cucumber tree	Moonsseed vine	Snowberry
Ash	Dogwood	Mountain ash	Spruce
Aspen	Elderberry	Multiflora rose	Sumac
Bindweed	Elm	Oak	Sumac, dwarf black
Birch	Giant ragweed	Olive, Russian	Sycamore
Black cherry	Gooseberry	Peppervine	Tamarisk
Black locust	Hackberry	Persimmon	Thistle
Blackberry	Hazel	Pine	Trumpet Vine
Brambles	Hemp	Pine, shortleaf	Trumpet creeper
Buckbrush	Himalayan blackberry	Poison ivy	Virginia creeper
Buffalo berry	Honeylocust	Poison oak	Wild plum
Bush morningglory	Honeysuckle	Poplar	Willow
Buttonbush	Honeysuckle, Japanese	Raspberry	Winterberry & other brush, trees & vines.
Cedar	Kudzu	Redvine	
Chamise	Lilac, common	Sagebrush	
Cherry, wild	Marijuana	Salibush	
Coastal sage	Manzanita	Serviceberry	
Cottonwood	Matrimony vine	Skunkbrush	

**HARDY WEEDS:**

For hardy weeds such as clover, ground ivy, oxalis, sheep sorrel, dock, thistles, and morningglory, and Virginia buttonweed in lawns and other areas, repeat application in two to three weeks if necessary.

**CONTROLS:**

**SENSITIVE WEEDS:**

Bedstraw	Chickweed	Knotweed	Plantain	Shepherdspurse
Black medic	Clover	Lambsquarters	Purslane	Wild lettuce
Buckhorn	Dandelion	Peppergrass	Ragweed	

**HARD-TO-KILL WEEDS:**

Chicory	Oxalis	Virginia buttonweed
Dock	Pigweed	Wild carrot
Ground ivy	Poison ivy	Wild onion
Healall	Poison oak	Yarrow & many other broadleaf weeds.
Lespedeza	Sheep sorrel	
Mallow	Speedwell	
Morningglory	Spurge	

**OTHER WEEDS CONTROLLED:**

Annual yellow sweetclover	Bulbous buttercup	Cocklebur	Heath aster
Aster	Bull thistle	Common mullein	Hedge bindweed
Austrian fieldcress	Bullnettle	Creeping jenny	Hedge mustard
Bedstraw	Burclover	Cudweed	Henbit
Beggarticks	Burdock	Curly dock	Hoary cress
Betony, Florida	Burning nettle	Daisy, English	Hoary plantain
Bindweed, field	Burweed	Daisy fleabane	Hoary vervain
Bird vetch	Buttercup	Daisy, oxeye	Horse-nettle
Bitter wintercress	Buttonweed	Dichondra	Jimsonweed
Bittercress, hairy	Canada thistle	Dogfennel	Knawel
Bitterweed	Carolina geranium	Dollarweed	Knotweed
Black-eyed Susan	Carpetweed	False dandelion	Kochia
Black medic	Catchweed bedstraw	False flax	Lambsquarters
Black mustard	Catnip	Fiddleneck	Mallow
Blackseed plantain	Catsear	Florida pusley	Lespedeza
Blessed thistle	Chickweed, common	Frenchweed	Matchweed
Bloodflower milkweed	Chickweed, mouseear	Galinsoga	Mexicanweed
Blue lettuce	Chicory	Goldenrod	Milk vetch
Blue vervain	Cinquefoil	Ground ivy	Morningglory
Bracted plantain	Clover, crimson	Gumweed	Mouseear hawkweed
Bristly ox-tongue	Clover, hop	Hairy fleabane	Mugwort
Broadleaf dock	Clover, red	Hawkweed	Musk thistle
Broadleaf plantain	Clover, strawberry	Heartleaf drymary	Mustard
Broomweed	Clover, sweet		Narrowleaf plantain
Buckhorn	Clover, white		Narrowleaf vetch
Buckhorn plantain	Cockle		Nettle

# 32 OZ LAWN WEED CONTROL CONCENTRATE BACK BOOKLET

EPA Reg No. 2217-540-75111

BLUE LINES DO NOT PRINT

**STORAGE AND DISPOSAL**

**STORAGE:** Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets. Do not store near food or feedstuffs.

**DISPOSAL: If Empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

**PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear face shield or goggles when mixing, loading or applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

**First Aid**

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

You may also call toll-free 1-877-800-5556 for emergency medical treatment information.

**Environmental Hazards:**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Distributed by:  
Maid Brands, Inc.  
Hazelwood, MO 63042

EPA Reg. No. 2217-540-75111  
EPA Est. No. 58996-M0-1

Do it Best® is a trademark of Do it Best Corp., Ft. Wayne, IN 46801.

PRESS TO RESEAL