

LICENSED

PERIOD 2015-2017 LIC. NO.

PERIOD 2009-2011 LIC. NO.

**8488.1**

**RESTRICTED USE PESTICIDE**

# BROMACIL 80

<b>ACTIVE INGREDIENT:</b>	<b>By Weight</b>
Bromacil: (5-bromo-3-sec-butyl-6-methyluracil) .....	80.0%
<b>OTHER INGREDIENTS:</b> .....	20.0%
<b>TOTAL:</b> .....	100.0%

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

FIRST AID	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

See inside label booklet for additional PRECAUTIONARY STATEMENTS

**PRECAUTIONARY STATEMENTS  
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
 CAUTION**

Harmful if absorbed through skin or swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

EPA Reg. No. 81927-4

EPA Est. No. 37429-GA-001

**Net Weight: 25 lbs.**

**Manufactured for:  
 ALLIGARE, LLC  
 13 North 8th Street, Opelika, AL 36801  
 Ph: 888-255-4427**

So  
ch  
Ap  
• L  
• C  
• S  
Mi:  
A r  
Die  
ma  
ara  
  
Wt  
agi  
U  
U  
•  
•  
  
Do  
wh  
Br  
soi  
pei  
  
It  
it  
Oth  
Trit  
  
U  
tt  
tr  
tr  
c  
D  
P  
tr  
•  
•  
  
T  
C  
D  
  
D  
P  
P  
C  
b  
  
Br  
tro  
Br  
As  
rai  
ha  
tex

RECEIVED  
 FEB 27 2006  
 Pesticides Branch



### USE PRECAUTIONS AND RESTRICTIONS

To avoid injury to or loss of desirable trees or other plants, observe the following use guidelines:

- Do not apply this product through any type of irrigation system.
- Except as recommended, do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use in any recreational areas or in or around homes, on lawns, walks, tennis courts, driveways, or other similar areas.
- Do not allow dry powder or spray to drift to desirable plants.
- Keep from contact with fertilizers, insecticides, fungicides, and seeds.
- Do not graze cattle in treated areas.
- Thoroughly clean all traces of Bromacil 80 from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens; clean these parts separately.
- Treated areas may be planted to citrus or pineapple one year after last application. Do not replant to other crops within two years after last application as injury may result.

#### When Preparing for Use:

- Calibrate sprayers only with clean water away from well sites.
- Regularly inspect spray equipment.
- Mix only enough Bromacil 80 for the specific application.
- Do not discharge excess material on the soil at a single spot in the field/grove or mixing/loading station.
- Ensure accurate measurement of pesticides.
- Avoid overfilling of spray tank.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.

#### Tank Mixture Specific Guidelines:

- Bromacil 80 may be tank mixed with other suitable herbicides registered for agricultural and non-agricultural use. Refer to the label(s) of the other products being added to the tank mix for any additional use information or restrictions. *Follow the most restrictive label guidelines.*
- When tank mixing with Bromacil 80, completely mix the product in the spray tank carrier (water, liquid fertilizer) before adding any other herbicide or spray adjuvant. A small compatibility test (see below) should be performed prior to adding the products into the spray tank using a combination of products not previously used. Refer to the "ADDITIONAL INSTRUCTIONS FOR AG AND NON-AG USES" section of this label for further information.
- The spray tank contents must be thoroughly re-agitated if they are allowed to settle for any period of time.

### APPLICATION INFORMATION

Follow the application guidelines below:

- Apply using a properly calibrated fixed-boom power sprayer.
- Bromacil 80 use rates listed on this label are expressed for broadcast treatments. For band treatments, use proportionately less.
- Use sufficient spray volume (minimum of 10 gallons per acre) to provide uniform coverage of the treated areas and to allow proper dispersion and suspension of the product in the spray tank.
- Continuous agitation in the spray tank is required to keep the product in suspension. Agitate spray tank contents by mechanical or hydraulic means. If a by-pass or return line is used, it should terminate at the bottom of the tank to minimize foaming. *Do not use air agitation.*
- Spray booms must be shut off while starting, turning, slowing or stopping. Overapplication of the herbicide may result in injury to the crop or successive crops.
- Nozzle screens should be 50 mesh or larger.

### SPRAY PREPARATION

**Mixing in Water** – Fill tank half full with water. Start agitation system and while continuing to add water, add Bromacil 80 and each additional component of any tank-mix separately. Be sure to agitate the entire time.

**Mixing with Other Herbicides** – Determine the tank mixture partner(s) compatibility with Bromacil 80 by following the directions below. If the testing procedure shows the mixture to be compatible, Bromacil 80 may be used in the tank mixture.

- 1 Put 1 pint of water into a quart jar with a tightly sealing lid.
- 2 In a separate container, combine 2 teaspoons of Bromacil 80 with 2 tablespoonfuls of water; mix thoroughly and add to the water.
- 3 Close the jar and shake well.
- 4 If additional herbicides are to be used in the mixture, follow steps two and three above for each additional herbicide.
- 5 Once all components of the tank mix are combined in the test jar, watch the mixture for several seconds and then check again in 30 minutes. If mixture does not separate, foam, gel or become lumpy, it may be used.

**Mixing in Liquid Fertilizer (citrus only)** – A fertilizer solution may be used in the spray mixture *only when applying to citrus*. Use the procedure above to test for compatibility before full-scale mixing, but in Step 1, in lieu of water, use the liquid fertilizer.

If the above procedure indicates the desired mixture will be compatible, prepare the tank mixture as follows:

- 1 Add the fertilizer solution to the spray tank first.
- 2 In a separate container, mix the required amount of Bromacil 80 with water to form a slurry that can be poured.
- 3 With the agitator running, slowly add the slurry to the tank and mix thoroughly.

### SPRAY TANK CLEAN-OUT

Thoroughly clean all traces of Bromacil 80 from application equipment immediately after use. Flush the tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Dispose of the equipment wash water by applying it to a use-site listed on this label.

### VERIFICATION OF SAFE ROTATIONAL USE IN ARID CLIMATES

In arid climates (areas that experience 10 inches of rainfall or less in a year) or areas that have experienced drought conditions for one or more years, a field bioassay should be conducted prior to planting any desired crop(s). The bioassay may consist of a test strip of the crop and should cross the entire field, including high- and low-lying portions. If a test strip of the crop(s) intended for production is not successfully grown to maturity, it may be necessary for the two-year crop rotation interval to be extended.

### WEED RESISTANCE TO HERBICIDES

Weeds may become resistant to any herbicide if an herbicide is used in the same field repeatedly over several years. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product with a different mode of action.

The following suggestions will assist in managing herbicide resistance:

- It may be necessary to change cropping practices within and between crop seasons. For example, using a combination of tillage, retreatment, tank-mixtures and/or sequential herbicide applications that have different modes of action.
- Preventing weeds from going to seed (by mowing, tilling, etc.) will prevent the spread of resistant plants.
- Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

Thi  
ecc  
itor  
col

Thi  
col  
Thi  
rec  
1.  
2. I  
(

IM  
Thi  
dro  
ho  
IF /  
"St

Co  
• V  
• F  
V  
• N  
C

Co  
• I  
• N  
ti  
• N  
t

BO  
• E  
u  
• E  
v

WI  
Dri  
inc  
DIT

No  
TE  
Wf

SU  
Dri  
ren  
are  
enc  
an  
sm

SE  
Thi  
enc

SH  
Shi  
ver



**Perennial Weeds Controlled**

Bahiagrass	<i>Paspalum notatum</i>
Bermudagrass	<i>Cynodon dactylon</i>
Johnsongrass	<i>Sorghum halepense</i>
Nutsedge	<i>Cyperus spp.</i>
Pangolagrass	<i>Digitaria decumbens</i>
Paragrass	<i>Panicum purpurascens</i>
Torpedograss	<i>Panicum repens</i>

**Note:** Partial control of perennials usually occurs with a single treatment; repeat applications are required to control perennials. Control of perennials may be improved by cultivation prior to treatment; otherwise, avoid working the soil as long as weed control continues otherwise effectiveness of the treatment may be reduced. Multiple applications may improve control of hard-to-kill weeds.

**AGRICULTURAL USES**

**NOTE:** Bromacil 80 use rates listed on this label are for broadcast treatments. For band treatments, use proportionately less.

**CITRUS**

Bromacil 80 may be applied as a broadcast or band treatment beneath and/or between trees. Avoid contact of foliage and fruit with spray or mist.

Important considerations for use of Bromacil 80 on Citrus:

- Following treatment, temporary yellowing of citrus leaves may occur.
- Do not use on soils with less than 1% organic matter as injury to citrus trees may result.
- Do not use on poorly drained soils, gravelly soils or thinly covered or exposed subsoils.
- Do not use on diseased or stressed trees.
- Do not use on trees planted in irrigation furrows.
- Do not use in citrus groves interplanted with other desirable trees or plants or in areas where roots of desirable trees or plants may extend as injury to desirable trees or plants may result.
- Do not use in home citrus plantings.
- Do not apply more than 8 pounds product (6.4 pounds active ingredient), inclusive of all bromacil formulations, per treated acre per year.
- When making multiple applications in a single growing season, do not apply at less than 60-day intervals.
- Bromacil 80 may be applied at any time of the year provided rainfall or overhead irrigation is available to activate the herbicide, preferably just before or just after weeds have germinated.

**U.S. (EXCEPT FLORIDA and TEXAS, see below for state specific instructions)**

**NOTE: THE FOLLOWING APPLIES TO TREES THAT ARE FOUR OR MORE YEARS OLD**

**For Control of Annual Weeds** – Apply 2 to 4 pounds of Bromacil 80 per acre. Apply anytime of the year, preferably shortly before or after weed growth begins when adequate moisture is available.

**For Control of Perennial Weeds** – Best results are obtained if application is made shortly before or shortly after weed growth begins; if dense growth is present, remove tops and spray the ground. Effects on perennial weeds are slow to appear, usually taking place over a period of several months.

Make a single application per year using the rates listed below during the period from winter to early summer:

Soil Texture	Pounds of BROMACIL 80 per Acre
Sand, Loamy Sand	4 to 5
Sandy Loam	5 to 6
Silt Loam, Clay Loam	6 to 8

Alternatively, except for Texas, make two applications of 3 to 4 pounds Bromacil 80 per acre per year. In Texas and Louisiana, apply in spring and summer; in California and Arizona, apply in fall and spring.

**Note:** Partial control usually occurs with a single treatment; repeat applications are required to control perennials. Control of perennials may be improved by cultivation prior to treatment; otherwise, avoid working the soil as long as weed control continues or else effectiveness of the treatment may be reduced.

Bromacil 80 controls the following:

Bermudagrass	All areas of the U.S.
Johnsongrass	Texas
Nutsedge	Texas, California

**Texas**

In Texas, do not apply more than 3 pounds Bromacil 80 per acre per year.

For control of the annual weeds listed above, apply 2-3 pounds of Bromacil 80 per acre. Use the lower rate for 1- to 2-year-old trees and the higher rate for trees more than 2 years old.

For best results in the control of perennial weeds make two applications, the first in spring and the second in the summer.

**Florida**

The use of Bromacil 80 is prohibited for weed control in non-bedded citrus groves located on any permeable, better drained soil identified in the intended site of application. Permeable, better drained soils which occur in citrus-producing areas of the state including unnamed soils and soils with characteristics of quartzipsamments, and the following soil series classifications:

Adamsville	Broward	Dade	Lake	Orsino	St. Augustine
Archbold	Canaveral	Florahome	Lakewood	Palm Beach	St. Lucie
Astatula	Candler	Fort Meade	Neilhurst	Paola	Tavares
Bahia Honda	Cocoa	Gainesville	Orlando	Satellite	

Treated areas may be planted to citrus trees one year after the last Bromacil 80 application. Do not replant to other crops within two years after the last Bromacil 80 application as plant injury may result.

**Application Instructions**

Apply Bromacil 80 as a band treatment **only** using a properly calibrated fixed-boom power sprayer. **Do not use Trunk to Trunk.**

Use sufficient spray volume (minimum of 10 gallons per acre) to provide uniform coverage of the treated area and to allow proper dispersion and suspension of the product in the spray tank. Continuous agitation in the spray tank is required to keep the product in suspension.

Avoid contact of foliage and fruit with spray or mist.

**NOTE:** All use rates of Bromacil 80 are expressed for broadcast treatments. For band treatments as required in Florida, use proportionately less.

**Trees Established Up to Three Years:** For control of annual weeds, apply 2 to 3 pounds of Bromacil 80 per treated acre as needed to maintain weed control. Do not apply more than 6 pounds of Bromacil 80 per treated acre within any 6-month period, or more than 8 pounds per treated acre per year.

**Trees Established Three or More Years:** For control of annual weeds, apply 2 to 4 pounds of Bromacil 80 per treated acre as needed to maintain weed control. Do not apply more than 4 pounds per treated acre in any one application. Do not use more than 8 pounds of Bromacil 80 per treated acre per year.

Annual Weeds Controlled		
Barnyardgrass	Lambsquarter	Sandspur
Crabgrass	Mustard	Sedge (annual)
Crowfootgrass	Natalgrass (red top)	Sprangletop
Florida pusley	Puncturevine	Texas panicum
Henbit	Purslane	Turkey mullein
Perennial Weeds Controlled		
Bahiagrass	Nutsedge	Paragrass
Bermudagrass	Pangolagrass	Torpedograss
Johnsongrass		

**Note:** Use the highest rates allowed by this label for best control of perennial weeds listed on this label. Partial control of perennial weeds can result with only a single treatment of Bromacil 80. Repeat applications are required (in season and/or annually) for best control of the perennial weeds on this label. Control of perennials may be improved by cultivation prior to treatment, otherwise, avoid working the soil as long as weed control continues or else effectiveness of the treatment may be reduced.

**Application Timing**

Allow a minimum interval of 60 days between applications of Bromacil 80.

**PINEAPPLE**

Treated areas may be planted to citrus or pineapple one year after last application. Do not replant to other crops within two years after last application as injury may result.

When making multiple applications, do not apply at less than 60-day intervals.

**Hawaii and Florida** – Apply 2 to 6 pounds per acre broadcast before the planting material begins to grow for control of seedling weeds such as amaranthus, balsamapple, chloris, crabgrass, Floras paintbrush, foxtail, goosegrass (wiregrass), and Hialoa. Use the lower rates in low rainfall areas (5 to 10 inches annually) and on clean-culture fields; use the higher rates in high rainfall areas (above 10 inches annually) and for transmulch fields. Follow with either 2 to 4 pounds per acre as a directed interline spray as needed prior to differentiation or apply 2 pounds per acre broadcast as needed up to differentiation. Do not apply more than 10 pounds of Bromacil 80 total per acre per plant crop.

For ratoon crop, apply 1 to 4 pounds per acre broadcast after harvesting plant crop but before differentiation. Do not apply more than 4 pounds total per acre per ratoon crop.

**Puerto Rico** – For control of seedling weeds such as crabgrass, goosegrass, jungle rice, pigweed, and purslane, apply 2 to 4 pounds Bromacil 80 per acre broadcast immediately after planting and before planting material begins to grow.

**NON-AGRICULTURAL USES**

**Use Restrictions – State of Florida**

**In Florida, the use of Bromacil 80 is prohibited in Hardee, Highland, Polk, Orange and Lake counties.** For Non-Agricultural Usage in all other areas of the state, do not apply more than 8 pounds of Bromacil 80 (or 6.4 pounds of bromacil) per acre per year, including all formulations of bromacil.

**Recommendations for Non-Agricultural Uses for Bromacil 80**

Bromacil 80 is recommended for control of undesirable vegetation for extended periods of time in non-crop areas such as railroads, highway and pipeline rights-of-way, petroleum tank farms, lumberyards, storage areas and industrial plant sites.

Do NOT apply this product to:

- Open water (such as creeks, estuaries, lakes, reservoirs, rivers, streams or saltwater bays);
- When water is present in fresh water wetlands (such as bogs, marshes, potholes or swamps);
- Saltwater marshes within tidal areas;
- Ditches, banks along waterways or impervious substrates; or,
- Areas near desirable plants where roots of these plants may extend.

**Application Information**

Apply Bromacil 80 using a properly calibrated fixed-boom power sprayer with sufficient spray volume to provide uniform coverage of the treated area and to allow proper dispersion and suspension of the product in the spray tank. All use rates of Bromacil 80 are expressed for broadcast treatments. For band treatments, use proportionately less.

**Notes for Non-Agricultural Uses:**

For small areas, a hand sprayer or sprinkling can be used. When preparing to spray a small area, 1/4 cup of Bromacil 80 per 200 sq. ft. is approximately 15 pounds per acre.

Use a spray volume of at least 40 gallons per acre to ensure uniform coverage.

**Application Timing**

Apply Bromacil 80 as a preemergence spray prior to or during the rainy season when weeds are actively germinating or growing. Moisture is required to move Bromacil 80 into the root zone of weeds for preemergence control. For best preemergence weed control, apply prior to rainfall and weed germination.

**Application Rates**

Apply Bromacil 80 at the rates indicated by weed type in the tables below. When applied at lower rates, Bromacil 80 provides short-term control of the weeds listed; when applied at higher rates, weed control is extended.

**Note:** Use the higher levels of the dosage ranges listed when applying on adsorptive soils (for example, those high in organic matter or carbon).

**Annual Broadleaf Weeds and Grasses – 3 to 6 pounds per acre**

Brome, downy	<i>Bromus tectorum</i>
Cheat	<i>Bromus secalinus</i>
Crabgrass	<i>Digitaria spp.</i>
Foxtails	<i>Setaria spp.</i>
Lambsquarter, common	<i>Chenopodium album</i>
Mullein, turkey	<i>Eremocarpus setigerus</i>
Oat, wild	<i>Avena fatua</i>
Puncturevine, common	<i>Tribulus terrestris</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ryegrass, Italian	<i>Lolium multiflorum</i>

When applied just prior to or just after emergence of annuals, rates as low as 2 pounds per acre control many annual weeds and grass in low rainfall areas and gives short term control in higher rainfall areas.

Applc  
Where  
satisf  
Retre  
not a  
**NOTE**

To coi  
spring  
cent t  
**Broad**  
soils (  
**Basal**  
to run

**UNDE**  
**Impoi**  
• Do r  
or g  
• Des

**Appli**  
Brom:  
media  
Brom:  
erage

**Appli**  
Brom:  
ical m

**Appli**  
Apply  
**Tank**  
To coi  
or Ou

If tem  
the se  
claim


Alliga  
used i  
**RANT**

It is ir  
becal  
soil cc  
tion, c

The e.  
ories),  
(1) Re  
(2) Re

Alliga  
or dar  
The te  
emplc  
in any

Oust\*



**Perennial Broadleaf Weeds and Grasses – 7 to 15 pounds per acre**

Bahiagrass	<i>Paspalum notatum</i>
Bermudagrass	<i>Cynodon dactylon</i>
Bluegrass	<i>Poa spp.</i>
Brome, smooth	<i>Bromus inermis</i>
Broomsedge	<i>Andropogon virginicus</i>
Carrot, wild	<i>Caudus carota</i>
Dandelion, common	<i>Taraxacum officinale</i>
Dogfennel	<i>Eupatorium capaillifolium</i>
Goldenrod	<i>Solidago spp.</i>
Johnsongrass	<i>Sorghum halepense</i>
Natalgrass (redtop)	<i>Rhynchelytrum repens</i>
Nutsedge	<i>Cyperus spp.</i>
Plantain	<i>Plantago spp.</i>
Purpletop	<i>Verbena bonariensis</i>
Quackgrass	<i>Agropyron repens</i>

Application rates as low as 5 pounds per acre will control many perennial weeds and grasses in areas with low or seasonal rainfall.

Where limited rainfall (usually less than 4 inches) occurs during the active growth period, such as some areas of the West, Bromacil 80 may not provide satisfactory control of hard-to-kill, deep-rooted perennial weeds such as johnsongrass.

**Retreating** – Apply 2 to 6 pounds of Bromacil 80 per acre when annual weeds and grasses reappear on sites where weed growth has been controlled. Do not apply Bromacil 80 at more than 15 pounds per acre per year per site.

**NOTE:** Do not apply at less than 60-day intervals when making multiple applications.

**BRUSH CONTROL**

To control undesirable woody plants on non-cropland areas such as railroad rights-of-way, storage areas, and industrial plant sites, apply Bromacil 80 in spring or summer as a broadcast or basal (spot) treatment. Do not use where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated area.

**Broadcast Treatment** – Apply 7 to 15 pounds per acre to control brush such as oak, pine, sweet gum, and willow. Use the higher rates on adsorptive soils (those high in organic matter or carbon).

**Basal (Spot) Treatment** – Mix 2-1/2 pounds in 5 gallons of water and apply at the rate of 1 to 2 fl. oz. per stem 2" to 4" in basal diameter; wet base of stem to runoff. Treatment controls woody plants such as cottonwood, hackberry, maple, oak, poplar, red bud, sweet gum, wild cherry, willow and winged elm.

**SPECIAL USES**

**UNDER ASPHALT AND CONCRETE PAVEMENT**

**Important Precautions when Applying Under Asphalt**

- Do not use Bromacil 80 under pavement in residential properties such as driveways, or in recreational areas, including jogging or bike paths, tennis courts, or golf cart paths.
- Desirable plants may be injured if their roots extend into treated areas or if planted in treated areas.

**Application Information**

Bromacil 80 may be used in combination with Oust® XP to control weeds under asphalt and concrete pavement such as parking lots, highway shoulders, median strips, roadways and other industrial sites.

Bromacil 80 should only be used in an area that has been prepared according to good construction practices. Use sufficient water to ensure uniform coverage, generally 100 gallons per acre. Agitate the tank continuously to keep Bromacil 80 in suspension.

**Application Timing**

Bromacil 80 should be applied immediately before paving to avoid lateral movement of the herbicide as a result of soil movement due to rainfall or mechanical means.

**Application Rates**

Apply Bromacil 80 at 6 to 15 pounds per acre. Use a higher rate on hard to control weeds and/or for longer term weed control.

**Tank Mixtures**

To control a broader spectrum of weeds, or for an extended period of weed control, a tank mixture of Bromacil 80 at 6 to 15 pounds per acre plus Oust® or Oust® XP at 4 to 8 ounces per acre may be used.

**TERMS AND CONDITIONS OF USE**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under "Warranty Disclaimer", "Inherent Risks of Use" and "Limitations of Remedies".

**WARRANTY DISCLAIMER**

Alligare, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Alligare, LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

**INHERENT RISKS OF USE**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Alligare, LLC or the seller. All such risks shall be assumed by buyer.

**LIMITATION OF REMEDIES**

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Alligare, LLC's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used

Alligare, LLC shall not be liable for losses or damages resulting from handling or use of this product unless Alligare, LLC is promptly notified of such loss or damage in writing. In no case shall Alligare, LLC be liable for consequential, incidental or special damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Alligare, LLC or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner. Buyer accepts this product subject to the foregoing limitations of warranty and of liability.

Oust® is a registered trademark of E.I. du Pont de Nemours and Company.

Some materials chemical-resistant

- Applicators and**
- Long-sleeved s
  - Chemical-resis
  - Shoes plus soc

**Mixers and loa**  
A non-powered

Discard clothing manufacturer's ir arately from othe

When handlers u agricultural pesti

**User Safety**  
Users should:

- Wash hands t
- Remove cloth
- Remove PPE
- Remove PPE clean clothing

Do not apply dire when cleaning ex

Bromacil is know soils are permea permeability and

It is a violation o other persons, e Tribe, consult the

Use this produc the protection c training, decont this label about covered by the

Do not enter or

PPE required fo treated, such as:

- Coveralls
- Chemical-resi
- Shoes plus sc

The requiremen CFR Part 170).  
Do not enter or

Do not contami

**Pesticide Sto**

**Pesticide Dis**

**Container Dis**  
by State and lo

Bromacil 80 is tc trol in certain cr Bromacil 80 also

As this product n rainfall or sprinkk has been carried texture, and othe

