



STATE OF HAWAII  
Department of Agriculture  
**ACCEPTED**  
LICENSE NO.

8581.32

**Combiva® II**  
FOR DOGS AND PUPPIES

**Over 55 lbs**  
7 weeks or older



Contains the same  
active ingredients  
as in Advantage® II  
for Dogs\*



**Combiva® II**  
FOR DOGS AND PUPPIES

**Over 55 lbs**  
7 weeks or older

**ACTIVE INGREDIENTS:**

Imidacloprid ..... 9.10 %

Pyriproxyfen ..... 0.46 %

**OTHER INGREDIENTS:** ..... 90.44 %

**TOTAL** ..... 100.00 %

Net Contents: 0.405 fl. oz. (3 doses, each 0.135 fl. oz.)

EPA Est. No. 74720-DEU-01

EPA Reg. No. 83399-17

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Back Panel / Package Insert for additional  
Precautionary Statements, First Aid  
and Directions for Use.

Use only on dogs and puppies weighing  
Over 55 lbs. and 7 weeks of age or older

Combiva® II for Dogs and Puppies  
treats and prevents further flea infestation  
on dogs and puppies 7 weeks and older



x3

\*Combiva® II for Dogs and Puppies is not  
manufactured or distributed by Bayer. Advantage is a  
registered trademark of Bayer.




**DO NOT USE ON CATS**



# Combiva® II

## FOR DOGS AND PUPPIES



**READ ENTIRE LABEL BEFORE EACH USE**  
**Use only on dogs and puppies 7 weeks**  
**or older weighing at least 3 lbs**

- Effectively kills adult fleas and prevents further re-infestation on dogs over 7 weeks of age weighing at least 3 lbs
- Kills re-infesting fleas within 2 hours
- Breaks the flea life cycle and prevents flea eggs and larvae from developing into adult fleas
- Effective once a month flea protection
- Combiva® II for Dogs and Puppies is waterproof after application

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Keep out of reach of children.

#### HAZARDS TO DOMESTIC ANIMALS

##### FOR EXTERNAL USE ON DOGS ONLY. DO NOT USE ON CATS.

Do not use on puppies under 7 weeks of age or puppies weighing less than 3 lbs. As with any product, consult your veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing dogs. If your dog is exhibiting signs or is being treated for skin dermatitis, talk to your vet before applying any topical flea and tick control product.

#### SIDE EFFECTS

Monitor your dog after application. Side effects, although rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects occur, consult your veterinarian or call 1-800-999-0297. Have the product container or label with you when calling your veterinarian for advice.

FIRST AID	
If swallowed:	Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything to an unconscious person.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	Wash with plenty of soap and water. Remove and wash contaminated clothing before reuse.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may contact 1-800-999-0297 weekdays between 9am and 6pm EST or 1-888-426-4435 for emergency medical treatment information.	

Manufactured by:  
 Ceva Animal Health, LLC  
 8735 Rosehill Road, Lenexa, KS 66215  
 D93160B-US / 17-B1-v1  
 Combiva® is a registered  
 trademark of Ceva Santé Animale S.A.  
 © 2016 Ceva Animal Health, LLC

#### DO NOT USE ON CATS

Keep cats away from treated dog for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian.



Made in Germany

**Combiva® II**  
**FOR DOGS AND PUPPIES**

**Over 55 lbs**  
**7 weeks or older**





# Combiva<sup>®</sup> FOR DOGS AND PUPPIES



**1** Tear through perforation



**2** Fold back the safety tab



**3** Cut with scissors to open applicator



# Combiva II

## FOR DOGS AND PUPPIES

3-10 lbs, 7 weeks or older  
 11-20 lbs, 7 weeks or older  
 21-55 lbs, 7 weeks or older  
 Over 55 lbs, 7 weeks or older

<b>ACTIVE INGREDIENTS:</b>	
Imidacloprid .....	9.10%
Pyriproxyfen .....	0.46%
<b>OTHER INGREDIENTS</b> .....	<b>90.44%</b>
<b>TOTAL</b>	<b>100.00%</b>

**Combiva II For Dogs and Puppies treats and prevents further flea infestation on dogs and puppies 7 weeks and older**

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

**READ ENTIRE LABEL BEFORE EACH USE**  
**Use only on dogs and puppies 7 weeks or older weighing at least 3 lbs**

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Keep out of reach of children.

#### HAZARDS TO DOMESTIC ANIMALS

##### FOR EXTERNAL USE ON DOGS ONLY. DO NOT USE ON CATS.

Do not use on puppies under 7 weeks of age or puppies weighing less than 3 lbs. As with any product, consult your veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing dogs. If your dog is exhibiting signs or is being treated for skin dermatitis, talk to your vet before applying any topical flea and tick control product.

FIRST AID	
If swallowed:	Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything to an unconscious person.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present after the first 5 minutes then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	Wash with plenty of soap and water. Remove and wash contaminated clothing before reuse.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may contact 1-800-999-0297 weekdays between 9am and 6pm EST or 1-888-426-4435 for emergency medical treatment information.	

#### SIDE EFFECTS:

Monitor your dog after application. Side effects, although rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects occur, consult your veterinarian or call 1-800-999-0297. Have the product container or label with you when calling your veterinarian for advice.

#### DO NOT USE ON CATS.

Keep cats away from treated dog for 24 hours.  
 If applied to a cat or ingested by a cat, contact your veterinarian.



### DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

#### CONSUMER INFORMATION


##### RESTRICTIONS:

- Do not allow children to apply product.
- For use only on dogs and puppies 7 weeks and older.
- Do not use on other animals.

- Do not apply to dogs or puppies weighing less than 3 lbs.
- Weigh your dog to be sure you are using the right size product for your dog.
- Do not apply more than one tube per treatment.
- Do not split one tube between two dogs.
- Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death.
- Do not have contact or allow children to have contact with treated area until completely dry.

**TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. FOR EXTERNAL USE ON DOGS ONLY. DO NOT USE ON CATS. DO NOT USE ON OTHER ANIMALS.**

**OPENING INSTRUCTIONS:**



**HOW TO APPLY:**


1. The dog should be standing or in a comfortable position for easy application. Apply the entire contents of the applicator evenly to three spots along the dog's back, from the shoulder to the base of the tail.

2. At each spot, part the hair down to the level of the skin and gently squeeze the applicator to apply the product. Avoid superficial application to the dog's hair.

3. Do not apply an excessive amount of solution to any one spot. This could cause the product to run off the dog. Do not get the product in the dog's eyes or mouth or allow your dog to ingest the product. Do not use more than one tube on dogs greater than 55 lbs.

4. Discard the empty applicator as outlined in the Storage and Disposal section.

5. Repeat every month, or as recommended by your veterinarian.



**FREQUENCY OF APPLICATION**

Treatment of Combiva® II For Dogs & Puppies will kill fleas on dogs and puppies within 12 hours and re-infesting fleas within 2 hours. It is possible pre-existing pupae in the environment may continue to emerge for 6 weeks or longer depending on climatic conditions. Use Combiva® II For Dogs & Puppies monthly for the control and prevention of flea re-infestations.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place. Keep out of reach of children.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND LIMITATION OF DAMAGES**

Seller warrants that the material conforms to the chemical parameters of the US EPA registration and the label. To the extent consistent with applicable law, seller makes no warranty, express or implied, other than indicated on the label. Buyer and user assume all risk of use and handling of this material. To the extent consistent with applicable law, any damages arising from use of this product or a breach of this warranty shall be limited to direct damages and shall not include consequential or incidental damages such as loss of profit or values.

D93130B-US / 08-A1-v1  
 Ceva Animal Health, LLC, 8735 Rosehill Road, Lenexa, KS 66215  
 Combiva® is a registered trademark of Ceva Santé Animale S.A.  
 EPA Reg. No. 83399-17 EPA Est. no. 74720-DEU-01

Made in Germany

51092

