

4.25"

Special Instructions for Cleaning and Decontamination Against HIV-1, HBV, and HCV of Surfaces or Objects Soiled With Blood or Body Fluids:

Personal Protection: When using this disinfecting towelette, wear appropriate barrier protection such as disposable protective gloves, protective gown, face mask and eye covering.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfecting towelette. A BARBICIDE Wipes towelette can be used for this purpose.

Contact Time: Thoroughly wet surface with a new BARBICIDE Wipes towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes. This contact time will not be sufficient for some of the organisms listed on the label. Refer to label for contact times against other organisms.

Disposal Of Infectious Materials: Cleaning materials used that may contain blood or other body fluids should be autoclaved and disposed of in accordance with Federal, State and local regulations for infectious materials disposal.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for further advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

STORAGE AND DISPOSAL

Towelette - Do not reuse towelette. Dispose of used towelette in trash.

Dispenser - Do not reuse empty container. Wrap container and discard in trash.

Packed For:
King Research, Inc.
2730 West Silver Spring Drive
Milwaukee, WI 53209

For emergency, product or technical information, contact King Research, Inc. at 800-222-8160 or visit our web site at www.barbicide.com.

EPA Reg. No. 46781-8-954
EPA Est. No. 56952-WI-001

ITEM NO. 13-2100

70-2101-3 80805-22B MADE IN USA



BARBICIDE® Wipes

disinfecting towelettes

• TUBERCULOCIDAL • VIRUCIDAL*
• BACTERICIDAL • FUNGICIDAL

BARBICIDE Wipes Disinfecting Towelettes are intended for use in salon settings, veterinary facilities, correctional facilities, schools, health club facilities, tanning salons, daycare centers and police and fire vehicles. BARBICIDE Wipes are effective against the following microorganisms on hard, non-porous surfaces when used as directed:

- *Mycobacterium tuberculosis* var: *bovis* (BCG)
- *Staphylococcus aureus*
- *Pseudomonas aeruginosa*
- *Salmonella choleraesuis*
- *Clostridium difficile* (vegetative cells only)
- *Trichophyton mentagrophytes*
- Methicillin Resistant *Staphylococcus aureus* (MRSA)
- Vancomycin Resistant *Enterococcus faecalis* (VRE)
- *Staphylococcus aureus* with reduced susceptibility to vancomycin
- Hepatitis B Virus (HBV)*
- Hepatitis C Virus (HCV)*
- Herpes Simplex Virus Types 1 and 2*
- Human Immunodeficiency Virus (HIV-1)*
- Influenza A2 Virus*

ACTIVE INGREDIENTS:

Diisobutylphenoxyethoxyethyl dimethyl benzyl ammonium chloride	0.28%
Isopropanol	17.20%
INERT INGREDIENTS†	82.52%
TOTAL	100.00%

† Does not include the weight of the dry towelette.

Net Contents: 1 lb. 13 oz.
160 Pre-Saturated Towelettes (6" x 6.75")



KEEP OUT OF REACH OF CHILDREN CAUTION

See other panel for additional precautionary statements.



LICENSED

LICENSED

PERIOD 2011-2013 LIC. NO.

PERIOD 2014-2016 LIC. NO.

16.75"

8619.1



LICENSED

PERIOD 2008-2010 LIC. NO.

- Critical Text Box
- Bleeds
- Adhesive area
- Die Cut area



Vipes

**TUBERCULOCIDAL • VIRUCIDAL*
BACTERICIDAL • FUNGICIDAL**

Barbicide Wipes are effective against the

Dimethyl benzyl ammonium chloride	0.28%
.....	17.20%
.....	82.52%
.....	100.00%

per dry towelette.

**3 oz.
Towelettes (6" x 6.75")**



**KEEP OUT OF
REACH OF
CHILDREN
CAUTION**

See other panel for additional precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BARBICIDE Wipes Disinfecting Towelettes are intended for use on hard, non-porous surfaces in salon settings, veterinary facilities, correctional facilities, schools, health club facilities, tanning salons, daycare centers, police and fire vehicles and other critical care areas where control of cross contamination between treated surfaces is required.

BARBICIDE Wipes will clean and disinfect, when used as directed, such items as infant/child care equipment surfaces, diaper changing stations, high chairs, SilkPeel™ with Dermalfusion™, workstations, stools, nail/hair care implements, tanning beds, hair dryers, hair clippers, shears, razors, hair cutting implements, clipper blades, salon surfaces, scissors, combs, brushes, manicure implements, washable nail files, vanity tops, tiles, floors, walls, handrails, door knobs, tables, chairs, faucets, bathroom fixtures, exterior toilet surfaces, garbage cans, carts, baskets, weight-lifting surfaces, cabinets, lights and telephones, which are made of plastic (i.e., polycarbonate, polypropylene, polyvinyl-chloride, polystyrene), and use sites listed on this label consisting of non-porous vinyl, stainless steel, painted surfaces, Plexiglas®, Formica® and glass.

DO NOT USE AS A BABY WIPE.

Not For Use On Skin

Dispenser Directions:

To start feed, remove cover and discard seal. From center of towelette roll, pull up a towelette corner, twist it into a point and thread it through the hole in the container cover. Pull through about one inch. Replace cover. Pull out first towelette and tear off at a 90° angle. Remaining towelettes feed automatically, ready for next

use. When not in use, keep center cap closed to prevent solution loss.

Cleaning Instructions:

Use one BARBICIDE Wipes towelette to completely preclean surface of all gross debris.

For use as a virucide*, including HIV-1, HBV and HCV, applications and effectiveness against Clostridium difficile (vegetative cells only), Methicillin Resistant Staphylococcus aureus (MRSA), Vancomycin, Resistant Enterococcus faecalis (VRE) and Staphylococcus aureus with reduced susceptibility to vancomycin:

Use a second BARBICIDE Wipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes at room temperature (69 °F/20 °C).

For use as a disinfectant:

Use a second BARBICIDE Wipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at room temperature (69 °F/20 °C) for Tuberculocidal activity and effectiveness against Staphylococcus aureus, Pseudomonas aeruginosa, Trichophyton mentagrophytes and Salmonella choleraesuis.

BARBICIDE Wipes effectively kills Human Immunodeficiency Virus Type 1 (HIV), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV) on precleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HIV, HBV, and HCV.

.75"

No Print Area • Glue Adhesive



Text Box

ve area

t area

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands before eating, drinking, chewing gum, or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for further advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

STORAGE AND DISPOSAL

Towelette - Do not reuse towelette. Dispose of used towelette and empty packet in trash.

Packed For:

King Research, Inc.
2730 West Silver Spring Drive
Milwaukee, WI 53209

For emergency, product or technical information, contact King Research, Inc. at 800-222-8160 or visit our web site at www.barbicide.com.

ITEM NO. 13-2124

70-2125-2 • 65092-22A • Patent # 7,228,968 B1 • MADE IN USA



KEEP OUT OF
REACH OF
CHILDREN
CAUTION

See other panel for additional
precautionary statements.

Net Contents: 12.6 oz.
45 Pre-Saturated Towelettes (7" x 9")

BARBICIDE® Wipes

disinfecting towelettes

- TUBERCULOCIDAL • VIRUCIDAL*
- BACTERICIDAL • FUNGICIDAL

PULL HERE

BARBICIDE Wipes Disinfecting Towelettes are intended for use in salon settings, veterinary facilities, correctional facilities, schools, health club facilities, tanning salons, daycare centers and police and fire vehicles. BARBICIDE Wipes are effective against the following microorganisms on hard, non-porous surfaces when used as directed:

- *Mycobacterium tuberculosis* var: *bovis* (BCG)
- *Staphylococcus aureus*
- *Pseudomonas aeruginosa*
- *Salmonella choleraesuis*
- *Clostridium difficile* (vegetative cells only)
- *Trichophyton mentagrophytes*
- Methicillin Resistant *Staphylococcus aureus* (MRSA)
- Vancomycin Resistant *Enterococcus faecalis* (VRE)
- *Staphylococcus aureus* with reduced susceptibility to vancomycin
- Hepatitis B Virus (HBV)*
- Hepatitis C Virus (HCV)*
- Herpes Simplex Virus Types 1 and 2*
- Human Immunodeficiency Virus (HIV-1)*
- Influenza A2 Virus*

ACTIVE INGREDIENTS:

Diisobutylphenoxyethoxyethyl dimethyl benzyl ammonium chloride	0.28%
Isopropanol.....	17.20%
INERT INGREDIENTS†:	82.52%
TOTAL	100.00%

† Does not include the weight of the dry towelette.

KING
RESEARCH®

EPA Reg. No. 46781-8-954

EPA Est. No.: 075272-WI-001
56952-WI-001

81043-22A

BARBICIDE® Wipes

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. BARBICIDE Wipes Disinfecting Towelettes are intended for use on hard, non-porous surfaces in salon settings, veterinary facilities, correctional facilities, schools, health club facilities, tanning salons, daycare centers, police and fire vehicles and other critical care areas where control of cross contamination between treated surfaces is required.

BARBICIDE Wipes will clean and disinfect, when used as directed, such items as infant/child care equipment surfaces, diaper changing stations, high chairs, SilkPeel™ with DermalInflusion™, workstations, stools, nail/hair care implements, tanning beds, hair dryers, hair clippers, shears, razors, hair cutting implements, clipper blades, salon surfaces, scissors, combs, brushes, manicure implements, washable nail files, vanity tops, tiles, floors, walls, handrails, door knobs, tables, chairs, faucets, bathroom fixtures, exterior toilet surfaces, garbage cans, carts, baskets, weight-lifting surfaces, cabinets, lights and telephones, which are made of plastic (i.e., polycarbonate, polypropylene, polyvinylchloride, polystyrene), and use sites listed on this label consisting of non-porous vinyl, stainless steel, painted surfaces, Plexiglas®, Formica® and glass.

DO NOT USE AS A BABY WIPE. Not For Use On Skin

Packet Directions:

Open packet, remove towelette and unfold.

Cleaning Instructions:

Use one BARBICIDE Wipes towelette to completely pre-clean surface of all gross debris.

For use as a virucide*, including HIV-1, HBV and HCV applications and effectiveness against Clostridium difficile

(vegetative cells only), Methicillin Resistant *Staphylococcus aureus* (MRSA), Vancomycin Resistant *Enterococcus faecalis* (VRE) and *Staphylococcus aureus* with reduced susceptibility to vancomycin:

Use a second BARBICIDE Wipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes at room temperature (69 °F/20 °C).

For use as a disinfectant:

Use a second BARBICIDE Wipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at room temperature (69 °F/20 °C) for tuberculocidal activity and effectiveness against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Trichophyton mentagrophytes* and *Salmonella choleraesuis*. BARBICIDE Wipes effectively kills Human Immunodeficiency Virus Type 1 (HIV), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HIV, HBV, and HCV.

Special Instructions for Cleaning and Decontamination Against HIV-1, HBV, and HCV of Surfaces or Objects Soiled With Blood or Body Fluids:

Personal Protection: When using this disinfecting towelette, wear appropriate barrier protection such as disposable protective gloves, protective gown, face mask and eye covering.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfecting towelette. A BARBICIDE Wipes towelette can be used for this purpose.

Contact Time: Thoroughly wet surface with a new BARBICIDE Wipes towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes. This contact time will not be sufficient for some of the organisms listed on the label. Refer to label for contact times against other organisms.

Disposal Of Infectious Materials: Cleaning materials used that may contain blood or other body fluids should be autoclaved and disposed of in accordance with Federal, State and local regulations for infectious materials disposal.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for further advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

STORAGE AND DISPOSAL

Towelette - Do not reuse towelette. Dispose of used towelette and empty packet in trash.

Packed For:
King Research, Inc.
2730 West Silver Spring Drive
Milwaukee, WI 53209

For emergency, product or technical information, contact King Research, Inc. at 800-222-8160 or visit our web site at www.barbicide.com.

ITEM NO. 13-2124

70-2125-2 • 65092-22A • Patent # 7,228,968 B1 • MADE IN USA



BARBICIDE® Wipes

disinfecting towelettes

• TUBERCULOCIDAL • VIRUCIDAL*
• BACTERICIDAL • FUNGICIDAL

PULL HERE

BARBICIDE Wipes Disinfecting Towelettes are intended for use in salon settings, veterinary facilities, correctional facilities, schools, health club facilities, tanning salons, daycare centers and police and fire vehicles. BARBICIDE Wipes are effective against the following microorganisms on hard, non-porous surfaces when used as directed:

- *Mycobacterium tuberculosis* var. *bovis* (BCG)
- *Staphylococcus aureus*
- *Pseudomonas aeruginosa*
- *Salmonella choleraesuis*
- *Clostridium difficile* (vegetative cells only)
- *Trichophyton mentagrophytes*
- Methicillin Resistant *Staphylococcus aureus* (MRSA)
- Vancomycin Resistant *Enterococcus faecalis* (VRE)
- *Staphylococcus aureus* with reduced susceptibility to vancomycin
- Hepatitis B Virus (HBV)*
- Hepatitis C Virus (HCV)*
- Herpes Simplex Virus Types 1 and 2*
- Human Immunodeficiency Virus (HIV-1)*
- Influenza A2 Virus*

ACTIVE INGREDIENTS:

Disodiumphenoxethoxyethyl
dimethyl benzyl ammonium chloride..... 0.28%

Isopropanol..... 17.20%

INERT INGREDIENTS†:..... 82.52%

TOTAL..... 100.00%

† Does not include the weight of the dry towelette.

Net Contents: 12.6 oz.
45 Pre-Saturated Towelettes (7" x 9")



KEEP OUT OF
REACH OF
CHILDREN
CAUTION

See other panel for additional
precautionary statements.

KING
RESEARCH®

EPA Reg. No. 46781-8-954
EPA Est. No.: 075272-WI-001
56952-WI-001

81043-22A

BARBICIDE® Wipes

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BARBICIDE Wipes Disinfecting Towelettes are intended for use on hard, non-porous surfaces in salon settings, veterinary facilities, correctional facilities, schools, health club facilities, tanning salons, daycare centers, police and fire vehicles and other critical care areas where control of cross contamination between treated surfaces is required.

BARBICIDE Wipes will clean and disinfect, when used as directed, such items as infant/child care equipment surfaces, diaper changing stations, high chairs, SilkPeel™ with DermalInfusion™, workstations, stools, nail/hair care implements, tanning beds, hair dryers, hair clippers, shears, razors, hair cutting implements, clipper blades, salon surfaces, scissors, combs, brushes, manicure implements, washable nail files, vanity tops, tiles, floors, walls, handrails, door knobs, tables, chairs, faucets, bathroom fixtures, exterior toilet surfaces, garbage cans, carts, baskets, weight-lifting surfaces, cabinets, lights and telephones, which are made of plastic (i.e., polycarbonate, polypropylene, polyvinylchloride, polystyrene), and use sites listed on this label consisting of non-porous vinyl, stainless steel, painted surfaces, Plexiglas®, Formica® and glass.

DO NOT USE AS A BABY WIPE. Not For Use On Skin

Packet Directions:

Open packet, remove towelette and unfold.

Cleaning Instructions:

Use one BARBICIDE Wipes towelette to completely pre-clean surface of all gross debris.

For use as a virucide*, including HIV-1, HBV and HCV applications and effectiveness against *Clostridium difficile*

(vegetative cells only), Methicillin Resistant *Staphylococcus aureus* (MRSA), Vancomycin Resistant *Enterococcus faecalis* (VRE) and *Staphylococcus aureus* with reduced susceptibility to vancomycin:

Use a second BARBICIDE Wipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes at room temperature (69 °F/20 °C).

For use as a disinfectant:

Use a second BARBICIDE Wipes towelette to thoroughly wet the surface. Repeated use of the product may be required to ensure that the surface remains visibly wet for 3 minutes at room temperature (69 °F/20 °C) for Tuberculocidal activity and effectiveness against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Trichophyton mentagrophytes* and *Salmonella choleraesuis*.

BARBICIDE Wipes effectively kills Human Immunodeficiency Virus Type 1 (HIV), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HIV, HBV, and HCV.

Special Instructions for Cleaning and Decontamination Against HIV-1, HBV, and HCV of Surfaces or Objects Soiled With Blood or Body Fluids:

Personal Protection: When using this disinfecting towelette, wear appropriate barrier protection such as disposable protective gloves, protective gown, face mask and eye covering.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfecting towelette. A BARBICIDE Wipes towelette can be used for this purpose.

Contact Time: Thoroughly wet surface with a new BARBICIDE Wipes towelette. Repeated use of the product may be required to ensure that the surface remains visibly wet for 2 minutes. This contact time will not be sufficient for some of the organisms listed on the label. Refer to label for contact times against other organisms.

Disposal Of Infectious Materials: Cleaning materials used that may contain blood or other body fluids should be autoclaved and disposed of in accordance with Federal, State and local regulations for infectious materials disposal.