

DOG MD

MAXIMUM DEFENSE™
Flea & Tick Topical

READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS 3 MONTHS OF AGE OR OLDER.
DO NOT USE ON CATS

PRECAUTIONARY STATEMENTS
HAZARDS TO DOMESTIC ANIMALS
CAUTION

FOR EXTERNAL USE ON DOGS ONLY. Do not use on puppies under 3 months of age. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. DO NOT USE ON CATS or animals other than dogs. Cats that actively groom or engage in close physical contact with treated dogs may be at risk of serious harmful effects. Sensitivities may occur after using ANY pesticide product on pets. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Signs of sensitivity to this product include skin sensitivity, increased itchiness, redness, rash, hair discoloration or hair loss at the application site. Dogs may also show "lethargy." Cats exhibiting these same symptoms from ingestion should be taken to veterinarian immediately.



DO NOT USE ON CATS. May be toxic or potentially fatal if applied to or ingested by cats. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to or near water. Drift and run-off may be hazardous to fish in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate. See Directions for Use for additional precautions and requirements. Treated dogs should not be allowed to go swimming for 12 hours following treatment.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Side Effects: Monitor your dog after application. The most common signs of ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of the product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If signs persist or become more severe within a few days of application, consult a veterinarian immediately by calling 1-877-416-7330. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application.

Do not get this product in your dog's eyes or mouth. Repeat applications may be made if necessary, but do not apply more often than once every 4 weeks.

ONLY for dogs weighing 15 lbs. & Under: Apply one tube (1.0 mL) (.034 fl oz) of this product as a spot or stripe to the dog's back between the shoulder blades.

HOW TO APPLY: Remove product tube from the package. Holding tube with notched end pointing up and away from face and body, cut off narrow end at notches. Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin. Wrap tube and put in trash.



STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 25° C), dry place out of reach of children. Do not refrigerate. Protect from direct sunlight.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Dog MD Maximum Defense™ Flea & Tick Topical is an effective and easy to use product. This product has demonstrated greater than 92% control of fleas within one day of application. It prevents eggs from fleas on treated dogs from developing into biting adults for up to 4 months. As with all flea and tick control products, **Dog MD Maximum Defense™ Flea & Tick Topical** should be used as part of a program aimed at reducing flea populations in the dog's environment (bedding, carpets, kennel, yard). Consult your veterinarian, entomologist, or retailer for program recommendations.

If you have questions or comments about this product, please write:
Consumer Response, P.O. Box 540224, Omaha, NE 68154-0224.

Distributed by: Meridian Animal Health, Omaha, NE 68154

In Case of Emergency, call: 1-877-416-7330, Non-Emergency, call: 1-888-814-4738 (4PET)

EPA Reg. No. 2517-87-84742
EPA Est. No. 2517-NE-02

3 MONTH SUPPLY

For use ONLY 15 lbs & under on dogs

DOG MD
MAXIMUM DEFENSE™

DOG MD
MAXIMUM DEFENSE™

For use ONLY 15 lbs & under on dogs

DOG MD
MAXIMUM DEFENSE™

Flea & Tick Topical

3 MONTH SUPPLY

For use ONLY 15 lbs & under on dogs

Kills fleas, ticks, flea eggs & flea larvae



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO. 8642.19

KEEP OUT OF REACH OF CHILDREN

CAUTION See Back Label Panel for Additional Precautionary Statements

ACTIVE INGREDIENTS:

Permethrin*	45.00%
Pyriproxyfen**	1.90%
OTHER INGREDIENTS:	53.10%
TOTAL	100.00%

* CAS# 52845-53-1
** Cis:trans ratio: Max 55% (+/-) cis and min 45% (+/-) trans
** CAS# 95737-68-1



Net Contents: 0.102 fl oz (3.0 mL) • Three 0.034 fl oz (1.0 mL) applicators



DOG MD
MAXIMUM DEFENSE™

For use ONLY 15 lbs & under on dogs

3 MONTH SUPPLY

P068325 RARH



8 41664 69080 6