

STATE OF HAWAII
Department of Agriculture

ACCEPTED
LICENSE NO.

8642.40

12 month
protection
with **2 collars**
per package

DOGMD
MAXIMUM DEFENSE™

Flea & Tick Collar
LARGE DOG



Kills Ticks



Kills Fleas



Waterproof



Long-Lasting
Protection



Compare to Scalibor®
Protector Band for Dogs,
contains the same
active ingredient.*

DO NOT ALLOW CHILDREN TO PLAY WITH COLLAR

CAUTION Net Weight: 2.02 oz (57.2 g)

Multi-Pack: Two 1.01 oz (28.6 g) Collars

ACTIVE INGREDIENT:

Deltamethrin 4.0%

OTHER INGREDIENTS: 96.0%

Total 100.0%

This product contains the toxic inert ingredient Bis (2-ethylhexyl) adipate (DEHA) at 24.76%.

*Scalibor® is a registered trademark of Intervet International B.V. This product is not manufactured or distributed by the makers of Scalibor® Protector Band for Dogs.

See Back Label Panel for Additional Precautionary Statements

P068471 RARM

FRONT

DogMD Maximum Defense™ Flea & Tick Collar LARGE DOG Fits Neck to 22"

**READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON DOGS • DO NOT USE ON CATS**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Do not open protective package until ready to use. Do not let children play with this collar. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

Do not use on puppies under 12 weeks. Do not use on cats. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated, or nursing animals. Sensitivities may occur after using ANY pesticide products for pets. If signs of sensitivity occur, remove collar and bathe your dog with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

This collar is intended for use only as an insecticide and is not to be taken internally by humans or animals.

If you have questions or comments about this product, please write:

Consumer Response, P.O. Box 540224, Omaha, NE 68154-0224

Distributed by: Meridian Animal Health, Omaha, NE 68154

In case of Emergency, call: 1-877-416-7330

Non-Emergency, call: 1-888-814-4738(4PET)

EPA Reg. No. 2517-166-84742

EPA Est. No. 42836-TX-1^A

EPA Est. No. 2517-NE-02^K

Superscript is the first letter in the lot code



PEEL HERE

P068406 AVR M



INSIDE FRONT

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may contact 1-877-416-7330 for emergency medical treatment information.

BASE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Do not open until ready to use. Dust will form on collar during storage. Do not get dust in mouth or eyes. Take the collar out of the packaging and stretch to full length. The tabs that held the collar in place will fall off when stretching; pull on any remaining tabs to remove them.
2. Bend the buckle in half between the holes in the buckle, so that you can easily pass the end of the collar through both holes.
3. Place the collar around your pet's neck. Feed the collar through both holes facing out. Fit to the neck of the dog. The collar must be worn loosely to allow for growth of the pet and to permit the collar to move about the neck. Generally, a properly fitted collar is one that, when fastened, will snugly slide over pet's head, and two fingers can be placed between the collar and pet's neck. Leave 1 or 2 extra inches for collar adjustment, cut off remaining excess collar and dispose of in the trash.
4. Wash hands after placing collar on pet. Replace collar every 6 months or sooner if needed.

STORAGE AND DISPOSAL

Pesticide Storage: Store this product in a cool, dry place away from children and pets. **Pesticide and Container Disposal:** Do not reuse container or used collar. Dispose of in trash.