

DISCONTINUED



STATE OF HAWAII
Department of Agriculture


ACCEPTED

LICENSE NO.

8793.5



Earth-tone[®] ***Garden Fungicide***



FOR

ORGANIC GARDENING

- For Flowers, Fruits, & Vegetables
- Use for Early & Late Blight on Tomatoes
- Controls Powdery Mildew, Black Spot, & Rust

Net Contents 24 fl oz.(1 pt 8 fl oz/709 ml)

Liquid Copper Fungicide Ready-to-Use

ACTIVE INGREDIENT	
Copper Octanoate (Copper Soap) ...	0.08%
OTHER INGREDIENTS	99.92%
TOTAL	100.00%

Metallic Copper Equivalent 0.017%

KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel for first aid and additional precautionary statements

Top Panel



Earth-tone® Garden Fungicide

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable directions and precautions on this label before using.

APPLICATION DIRECTIONS

Spray all plant surfaces (top and bottom of leaves) thoroughly with EARTH-TONE® GARDEN FUNGICIDE. 22 fluid ounces will treat 7 square yards.

Begin treatment 2 weeks before disease normally appears or when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. Re-apply after rain.

To control powdery mildews, begin treatment 2 weeks before disease normally appears. Alternatively, begin treatment when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. Spray every 7 days when the amount of disease is increasing rapidly. If possible, time applications so that 12 hours of dry weather follow application. On plants that are very susceptible to powdery mildew, such as home-greenhouse-grown cucumber, it is best to spray the plants twice a week during the first 2 weeks after emergence, and weekly thereafter. On outdoor plants, re-apply after rain.

To control downy mildews, leaf and fruit spots, blights, and rust, begin treatment 2 weeks before disease normally appears or when weather forecasts predict a long period of wet weather. Alternatively, begin treatment when disease first appears, and repeat at 7 to 10 day intervals for as long as needed. Spray all plant parts thoroughly. Re-apply following heavy rain. Apply every 7 days when the amount of disease is increasing rapidly. Sprays every 5 to 7 days should be used for preventing late blight on potatoes and tomatoes. If possible, time applications so that at least 12 hours of dry weather follows application.

EPA REG. NO. 67702-1-83598
EPA EST. 4-NY-1



Sold under a license of
W. Neudorff GmbH KG, Germany.

Distributed by The Espoma Company
6 Espoma Road, Millville, NJ 08332

For more information on EARTH-TONE®
GARDEN FUNGICIDE and other Espoma
products call 1-888-ESPOMA-1 or
visit our website www.espoma.com.



For Additional Directions
and Precautions PEEL HERE 

Back Panel

To prevent fruit rots, spray EARTH-TONE® GARDEN FUNGICIDE thoroughly onto flowers and fruit. Apply at the start of flowering and continue every 7 to 10 days until harvest. Fungicidal sprays are especially warranted when weather forecasts predict a long period of wet weather. Re-apply after rain.

GRAPES

Note: Do not mix EARTH-TONE® GARDEN FUNGICIDE with lime. Certain Vinifera and French Hybrid varieties may be sensitive to copper sprays resulting in marginal leaf burn. Before spraying these varieties, consult your State Experiment Station or make test sprays.

Downy mildew, Black rot, Phomopsis Cane and Leaf Spot, Powdery mildew
Spray all plant surfaces thoroughly with EARTH-TONE® GARDEN FUNGICIDE. For best control begin treatment when new growth reaches ½ inch and repeat at 7 to 14 day intervals throughout the growing season.

Gray mold (Fruit rot)

Spray all plant surfaces thoroughly with EARTH-TONE® GARDEN FUNGICIDE. For best control begin treatment at the end of bloom and repeat at 7 to 14 day intervals.

PEACHES

Peach leaf curl

Spray all plant surfaces thoroughly with EARTH-TONE® GARDEN FUNGICIDE. Can be used to control peach leaf curl. Apply as a dormant spray in late fall during a period of dry weather.

ORNAMENTAL PLANTS Spray all plant surfaces thoroughly with Earth-tone® Garden Fungicide. Begin treatment when new growth emerges and repeat every 7 to 10 days for as long as needed to control disease. EARTH-TONE® GARDEN FUNGICIDE may cause some copper toxicity on some plant species. Before spraying a specific plant species, consult your State Experiment Station or make a test spray.

ROSES

Blackspot, Downy mildew, Gray mold, Leafspots, Powdery mildew, Rust Spray all plant surfaces thoroughly with EARTH-TONE® GARDEN FUNGICIDE. Begin treatment when new spring growth emerges and repeat every 7 to 10 days for as long as needed to control disease. EARTH-TONE® GARDEN FUNGICIDE may cause some copper toxicity on some roses. Copper toxicity appears as purple spots.

Base

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Caution: Harmful if swallowed, absorbed through skin or inhaled. Wash thoroughly with soap and water after handling. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS: This product is toxic to fish and aquatic organisms and may contaminate water through runoff. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

FIRST AID

IF IN EYES: -Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. -Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: -Take off contaminated clothing. -Rinse skin immediately with plenty of water for 15-20 minutes. -Call a poison control center or doctor for treatment advice.

IF SWALLOWED: -Call a poison control center or doctor immediately for treatment advice. -Have person sip a glass of water if able to swallow. -Do not induce vomiting unless told to by a poison control center or doctor. -Do not give anything by mouth to an unconscious person.

IF INHALED: -Move person to fresh air. -If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. -Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

STORAGE AND DISPOSAL

Pesticide Storage

Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals. Store in a secure place, away from open fire or flame. Keep container closed and reseal after use. Product may be damaged by freezing. Do not store product below 40°F (4°C). If spilled, use absorbent materials and dispose of in an approved manner.

Pesticide Disposal and Container Handling

Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions.

Never place unused product down any indoor or outdoor drain.