Doc ID: 522624

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

8816.9

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

FIRST SHIELD™

Dog & Puppy

FOR DOGS AND PUPPIES 21-55 LBS OVER 8 WEEKS OF AGE

EFFECTIVE MONTHLY CONTROL OF:

- Adult Fleas
- Flea Larvae
- Flea Eggs
- Flea Pupae

ACTIVE INGREDIENTS:

Dinotefuran	22.00%
Pyriproxyfen	3.00%
OTHER INGREDIENTS	75.00%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN

WARNING: See package insert for directions for use, precautionary statements and first aid and other information.

- Quick onset of activity within 2 hours; kills fleas in 6 hours
- Kills fleas that may transmit diseases, including bartonellosis, tapeworm and tularemia
- Prevents reinfestation through 30 days
- Kills fleas that may cause flea allergy dermatitis and anemia
- Convenient topical treatment with patented applicator
- Quick drying, non-greasy

4 DOSES

Net Contents:
4 doses, each 0.135 fl. oz. (4.0 mL)

RECEIVED
JAN 18 2011
Pesticides Branch

FIRST SHIELD™
Dog & Puppy

FOR DOGS AND PUPPIES OVER 8 WEEKS OF AGE
WEIGHING 21 LBS TO 55 LBS

4 DOSES

FIRST SHIELD™ Dog & Puppy

FOR DOGS AND PUPPIES OVER 8 WEEKS OF AGE WEIGHING 21 TO 55 LBS

Available From Licensed Veterinarians Only

READ ENTIRE LABEL BEFORE EACH USE. See package insert.

USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
WARNING: Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum and using tobacco. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS
For external use on dogs only. Do not use on puppies less than 8 weeks old.
Do not use this product on debilitated, aged, medicated, pregnant or nursing animals or animals known to be sensitive to pesticide products without first consulting a veterinarian. Sensitivity, such as slight transitory redness of the skin at the site of application, may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap or shampoo and rinse with large amounts of water. If signs of individual animal sensitivity occur and persist, contact your veterinarian. Have the product container or label with you when calling your veterinarian for advice.

FIRST AID	**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.	**If swallowed:** Call a poison center doctor for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything to an unconscious person.	**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center, doctor, or **1-877-872-3744** or after 6 pm EST **1-888-426-4435** immediately for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may contact **1-877-872-3744**, or after 6 pm EST **1-888-426-4435** for emergency medical treatment information.			

SATISFACTION *Guaranteed*
Schuyler LLC will provide new product or your money back.

Manufactured in the USA
for Distribution By:
Schuyler LLC
8000 NE Tillamook Street
Portland, Oregon 97213-0998

Schuyler, LLC

Net Contents: 4 doses, each 0.135 fl. oz. (4.0 mL)

EPA Reg No. 83399-10-85581 EPA Est. No. 68669-NC-1

FIRST SHIELD™
Dog & Puppy

• over 8 weeks of age
• 2.5-10 lbs • 11-20 lbs • 21-55 lbs • 56-100 lbs

Effective monthly control of:
• Adult fleas • Flea eggs • Flea larvae • Flea pupae

- Quick onset of activity within 2 hours; kills fleas within 6 hours
- Kills fleas that may transmit diseases including bartonellosis, tapeworm and tularemia
- Quick drying, non-greasy
- Kills fleas that may cause flea allergy dermatitis and anemia
- Convenient topical treatment with patented applicator
- Prevents reinfestation through 30 days

AVAILABLE FROM LICENSED VETERINARIANS ONLY

ACTIVE INGREDIENTS:
Dinotefuran..... 22.00%
Pyriproxyfen.....3.00%
OTHER INGREDIENTS....75.00%
TOTAL.....100.00%

KEEP OUT OF REACH OF CHILDREN

WARNING:

READ ENTIRE LABEL BEFORE EACH USE

USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE

FOR MORE INFORMATION, CALL 1-877-872-3744

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

WARNING: Causes substantial, but temporary eye injury. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum and using tobacco. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS

For external use on dogs only. Do not use on puppies less than 8 weeks old.
Do not use this product on debilitated, aged, medicated, pregnant or nursing animals or animals known to be sensitive to pesticide products without first consulting a veterinarian. Sensitivity, such as slight transitory redness of the skin at the site of application, may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap or shampoo and rinse with large amounts of water. If signs of individual animal sensitivity occur and persist, contact your veterinarian. Have the product container or label with you when calling your veterinarian for advice.

FIRST AID

If in eyes:	Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center, or doctor for treatment advice.
If swallowed:	Call a poison control center, or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
If on skin:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center, doctor, or **1-877-872-3744** or after 6 pm EST call **1-888-426-4435** immediately for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may contact **1-877-872-3744** or after 6 pm EST call **1-888-426-4435** for emergency medical information.	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. [Precaucion al consumidor: si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (To the User: If you cannot speak English, do not use this product until the label has been fully explained to you.)]

HOW TO APPLY:

1. **USE ONLY ON DOGS. Do not use on cats or other animals.**
2. Remove applicator from package.
3. The dog should be standing or in a comfortable position for easy application.
4. Holding the applicator upright, place thumb and index finger around the applicator tip under the large disk. With other hand, grasp the stem of applicator tip above smaller disk. Press down firmly on small disk until both disks meet, piercing the seal. (See illustration.)
5. Using the tip of the applicator, part the hair between the shoulder blades and slowly apply the product (as shown below) directly onto the skin until the applicator tube is completely empty. Avoid superficial application to the animal's hair.







For dogs weighing between 2.5-10 lbs OR 11-20 lbs, apply the product to 1 spot, as shown, onto the skin.



For dogs weighing 21-55 lbs, apply the product evenly to 3 spots along the dog's back, beginning between the shoulder blades and continuing in the order shown in the diagram.

For dogs weighing 56-100 lbs, apply the product evenly to 4 spots on the dog's back, beginning between the shoulder blades and continuing in the order shown in the diagram.

6. Discard empty applicator tube as outlined in the storage and disposal section.
7. For treatment, control and prevention of fleas, flea eggs, flea larvae and flea pupae, year-round treatment is recommended.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Storage: Store in a cool, dry place. Protect from freezing.

Disposal: If empty, do not reuse the container. Place in trash or offer for recycling if available. If partially filled, call your local solid waste agency for disposal instructions. Never place unused products down any indoor or outdoor drain.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that the material conforms to the chemical parameters of the US EPA registration and the label. Seller makes no warranty to the extent consistent with applicable law, express or implied, other than indicated on the label. Buyer and user assume all risk of use and handling of this material when such use and handling are contrary to the label directions. To the extent consistent with applicable law any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profit or values.

NET CONTENTS: each 0.044 fl. oz. (1.3 mL)
each 0.068 fl. oz. (2.0 mL)
each 0.135 fl. oz. (4.0 mL)
each 0.204 fl. oz. (6.0 mL)

EPA Reg No. 83399-10-85581
EPA Est. No. 68669-NC-1

Manufactured in the USA
for Distribution By:

Schuyler, LLC
8000 NE Tillamook Street
Portland, Oregon 97213-0998

SATISFACTION Guaranteed
Schuyler LLC will provide new product or your money back.



OPEN END
CENTER LINE
CENTER LINE
3 mm — 0.118" Eyemark
9.5 mm — 0.374" Free Copy Area
MONTEBELLO LENGTH: 2 9/32"
TUBE LENGTH: 2 1/4"
OVER
Side Seam – Overlap
UNDER
FIRST SHIELD
Dog & Puppy
OVER 8 WEEKS OF AGE
WEIGHING 21–55 LBS
4.0mL
Dinotefuran..... 22.00%
Pyriproxyfen..... 3.00%
WARNING
Read Entire Label Before Use
Keep Out Of Reach Of Children
EPA Reg. No. 83399-10-85581 EPA Est. No. 68669-NC-1
5 mm 0.197"
Shoulder .7925 mm 0.0312"
CAP END
5 mm 0.197"
5 mm 0.197"
Trim 2 mm 0.079"
BACK PANEL
15.859 mm – 0.624"
FRONT PANEL
19.95 mm – 0.785"
Trim 2 mm 0.079"
SLIT WIDTH 45.395 mm – 1.7872"