

FINAL



Blocks Weeds Before They Sprout™

Northern Garden Formula



WeedBlock®

PRE-EMERGENT NORTHERN

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel.

ACTIVE INGREDIENT:

Prodiamine*0.22%

OTHER INGREDIENTS:99.78%

TOTAL.....100.00%

*CAS No. 29091-21-2

NET WT 4.65 LBS (2.11 kg)



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2015-2017** LIC. NO.

8870.2



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2012-2014** LIC. NO.



Easy Gardener Products, Inc.
P.O. Box 21025
Waco, Texas 76702-1025 USA
Phone: 1-800-327-9462
www.easygardener.com

© Copyright 2012 Easy Gardener Products, Inc.

Easy Gardener and WeedBlock are registered trademarks of Easy Gardener Products, Inc.

EPA Reg. No. 74468-6-46260 EPA Est. No. 81245-MN-1
NET CONTENT: 4.65 Lbs.

PROOF OF PURCHASE



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

See text at top of next column.



WeedBlock®

PRE-EMERGENT NORTHERN

Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or physician immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or physician. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1212 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight as deemed appropriate to the situation.

See inside booklet for complete directions for use, plant listings, and precautionary statements

Lift here to open. Press to reseal.




and on established lawns 60 days after overseeding. WeedBlock Pre-Emergent Northern may also be incorporated into the soil or applied after mulching beds. Since it does not kill existing weeds, first remove any that have already begun to grow. Best results are obtained when applied to soil free of weeds, clods and debris such as leaves. Applying WeedBlock Pre-Emergent Northern as early as possible eliminates the need for hand weeding. Rain does not wash this product out of the soil. **To keep ornamental beds, ground covers, trees, shrubs and lawns weed-free all season long, reapply WeedBlock Pre-Emergent Northern every 60 to 90 days; however, do not exceed the maximum use rates listed in the "Maximum Annual Application Rate" table of this label.**

HOW TO APPLY WEEDBLOCK PRE-EMERGENT NORTHERN To avoid plant injury, do not apply WeedBlock Pre-Emergent Northern when foliage is wet. When plants are dry, sprinkle WeedBlock Pre-Emergent Northern on soil surface at rate given for your specific planting situation. Sprinkle over the top of established ground covers and lawns. Watering immediately after application will wash the granules off the foliage and activate the product in the soil. Always wash or brush off any granules from plant foliage to avoid damage or discoloration. Once activated by watering, WeedBlock Pre-Emergent Northern forms a weed control barrier in the top layer of soil, preventing weed seeds from germinating in ornamental gardens, ground covers, ornamental grasses, shrubs, trees and turf grasses listed on this label. Disturbance to the soil surface after activation may result in erratic weed control.

HOW MUCH WEEDBLOCK PRE-EMERGENT NORTHERN TO APPLY

Apply WeedBlock Pre-Emergent Northern evenly at the rate of 1 teaspoon per square foot to the soil surface, ground cover, mulch or lawn as illustrated by the picture below. Immediately water to incorporate WeedBlock Pre-Emergent Northern into the soil.



areas where water is not readily available, lightly rake the surface of the soil to incorporate into the root zone. Wash granules off of sidewalks, driveways, patios and other hard surfaces to avoid discoloration.

APPLYING WEEDBLOCK PRE-EMERGENT NORTHERN WITH MULCH

For best weed control, apply WeedBlock Pre-Emergent Northern after spreading mulch on flower and shrub beds. Sprinkle WeedBlock Pre-Emergent Northern on top of mulch at surface rate noted above (see picture above), being sure to keep granules away from plant foliage. Immediately water WeedBlock Pre-Emergent Northern into mulch. If water is not readily available, lightly rake the product into top layer of mulch. Always wash or brush off stray granules from plant foliage to avoid damage or discoloration.

APPLYING WEEDBLOCK PRE-EMERGENT NORTHERN WITH LANDSCAPE FABRIC

WeedBlock Pre-Emergent Northern may be applied to the soil prior to the installation of landscape fabric, or over the top of landscape fabric that has previously been installed.

USE PRECAUTIONS AND RESTRICTIONS

The following precautions apply to the use of this product in turf grasses and lawns.

- Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
- Do not apply to newly set sod until the following year.
- Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control.
- Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
- Do not apply through any type of irrigation system.
- Do not apply aerially.
- Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION

WeedBlock Pre-Emergent Northern may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates listed for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control. WeedBlock Pre-Emergent Northern is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not apply more than 680 lbs./A (1.5 lb. A/A) of this product per calendar year.

MAXIMUM ANNUAL APPLICATION RATES	
Turf Species	Maximum Annual Rate ¹
Bahiagrass Bermudagrass ² Centpedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysiagrass	15.3 lbs/1,000 sq. ft. per year (680 lbs/A per year) (1.5 lbs A/A)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	10.2 lbs/1,000 sq. ft. per year (455 lbs/A per year) (1.0 lbs A/A)
Creeping Red Fescue	8.0 lbs/1,000 sq. ft. per year (340 lbs/A per year) (0.75 lbs A/A)
Creeping Bentgrass	6.8 lbs/1,000 sq. ft. per year (295 lbs/A per year) (0.65 lbs A/A)

¹These are the maximum rates per calendar year.
²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 235 lbs./A. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

WEEDS CONTROLLED

When used according to label instructions, WeedBlock Pre-Emergent Northern will control the following weeds:

Barnyardgrass Bluegrass, Annual (<i>Poa annua</i>) Carpetweed Chickweed, Common Chickweed, Mouseear (from seed) Crabgrass (Large, Smooth) Crownfootgrass Cupgrass, Woolly Foxtails, Annual	Goosegrass ¹ Henbit Itchgrass Johnsongrass (from seed) Junglerice Knotweed Kochia Lambsquarters, Common Lovegrass Panicum (Texas, Fall, Browntop)	Pigweed Purslane, Common Pusley, Florida Rescuegrass ² Shepard's Purse ² Signalgrass, Broadleaf Speedwell, Persian Sprangletop Spurge, Prostrate Witchgrass Woodsorrel, Yellow (from seed)
--	--	---

¹In many areas a single application of 295 to 680 lbs./A will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 295 to 455 lbs./A, followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate ("Maximum Annual Application Rate" table in the Rates of Application section of this label).

²For this weed, apply in late summer, fall or winter prior to germination.

³Suppression only.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).

WHEN TO OVERSEED AFTER APPLICATION

WeedBlock Pre-Emergent Northern will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results.

USE RATE	MONTHS BEFORE OVERSEEDING			
	AI/A	North	Transition	South
0.50	4	4	4	4
0.66	5	4	4	4
0.74	6	5	5	5
0.80	-	6	6	6
1.00	-	7	7	7
1.14	-	-	9	9
1.30	-	-	10	10
1.50	-	-	12	12

LANDSCAPE ORNAMENTALS (including established perennials and wildflower plantings)

WeedBlock Pre-Emergent Northern may be applied for residual pre-emergence weed control in ornamentals listed on this label.

APPLICATION TIMING AND INFORMATION

WeedBlock Pre-Emergent Northern is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

WeedBlock Pre-Emergent Northern may be applied at any time to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate the product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

WeedBlock Pre-Emergent Northern may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle and



water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

WeedBlock Pre-Emergent Northern will not harm most trees, shrubs, vines and flowers listed on this label. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME	COMMON NAME
Abies spp.	Fir species** (Balsam, Fraser, Noble, etc.)
Abelia grandiflora	Abelia; Sherwood
Acer palmatum	Japanese Maple
A. Platanoides	Norway Maple
Achillea spp.	Yarrow; King Edward
Actinidia chinensis	Kiwifruit
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Agapanthus orientalis	
Akebia quinata	Five-leaf or Chocolate Vine
Allium cernuum	Lady's Leek, Nodding Onion
Anemone hybrida	Japanese Anemone
Aquilegia spp.	Aquilegia; Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemisia spp.	Wormwood; Silver Mound, Castle
Aster spp.	Aster; Bonnie Blue, Purple Dome
Aster X Frikartii	
Athrium Filix-femina	Lady Fern; Fern Lady
Aucuba japonica	Japanese Aucuba
Begonia spp.	Fibrous Begonia; Hardy Grasses
Berberis gladywensis	Barberry
B. julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. thunbergii	Japanese Barberry
B. verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower, (White Clips)
Campsis X Tagliabuona	Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisioides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Chrysanthemum nipponicum	
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Cornus Florida	Flowering Dogwood
C. Stolonifera	American Dogwood
Cortaderia selleana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Barberry Cotoneaster

SCIENTIFIC NAME	COMMON NAME
C. Microphyllus	Rockspray Cotoneaster
Crateagus spp.	Hawthorne
Crocosmia spp.	Luzifer
Cyprusus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma spp.	Cooperi Pink
Delphinium spp.	Larkspur; Blue Elf
Dianthus deltoideus	Dianthus; Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonaea viscosa	Hop Bush
Echinacea purpurea	Coneflower, Purple; Magnus
Elaeagnus pungens	Silverberry
Euonymus fortunei	Winterreeger
E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
Forsythia suspensa	Weeping Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia spp.	Gaillardia; Blanket Flower; "Goblin"
Gardenia jasminoides	Gardenia, Cape-Jasmine
Gaura spp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranebill
Gladiolus spp.	Gladiolus species**
Gypsophila repens	Baby's Breath
Hedera helix	English Ivy
Hellanthenum spp.	Sunrose
Hemerocallis spp.	Daylily; Aztec Gold, Stella de Oro, Tender Love
Hibiscus	Rose of Sharon**
Hibiscus spp.	Mallow; Disco Belle White
Hibiscus rosa-sinensis	Chinese Hibiscus
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. Sieboldiana	Hosta, "Seersucker"
Houttuynia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
Ilex Bernii	Holly
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. vomitoria	Yaupon Holly, Schillings
Inula ensifolia	
Iris ensata	Sword-Leaved Iris; Jodlesong
Iris sibirica	Siberian Iris; Cabernet
Iris spp.	Iris species**
Jasminium nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii
J. conferta	Shore Juniper; Blue Pacific
J. davurica	Parsoni
J. horizontalis	Creeping Juniper
Juglans spp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crape Myrtle
Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora
Lantana montevidensis	Weeping Lantana
Lavender spp.	Lavender; Munstead
Leontopodium alpinum	Edelweiss
Ligustrum amurense	Amur Privet
Ligustrum sinense	Chinese Privet; Variegata
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Lilium spp.	Lily; Jazz
Liriope muscari	Big Blue Lily, Silver Mound, Evergreen Giant
Liriope muscari var. variegata	Liriope, Variegated
Liriope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Lorapetalum chinense	Burgundy
Lythrum spp.	Loosestrife; Modern Pink

SCIENTIFIC NAME	COMMON NAME
Magnolia spp.	Magnolia species**
Malephora luteola	Ice Plant
Malus spp.	Crabapple*
Miscanthus sinensis	Yaku, Jima, Silberfeder
Nandina domestica	Heavenly Bamboo
Narcissus spp.	Narcissus species**
Nerium oleander	Oleander
Olea europaea	Olive*
Ophiopogon japonicus	Mondo Grass
Osmanthus heterophyllus	Osmanthus (False Holly); Gulf Tide
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
Paonia suffruticosa	Tree Peony
Pennisetum setaceum	Fountain Grass (Dwarf)
Perovskia atriplicifolia	
Persea americana	Avocado*
Photinia Fraseri	Frasier's Photinia (Redtip)
Physotegia Virginiana	Dragonhead, False; Vivid
Picea spp.	Spruce species** (Colorado Blue, Norway, etc.)
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine
P. canariensis	Canary Island Pine
P. Elliotti	Slash Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Pine
P. palustris	Longleaf Pine
P. radiata	Monterey Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
P. taeda	Loblolly Pine
P. thunbergiana	Japanese Black Pine
P. virginiana	Virginia Pine
Pistachio spp.	Pistachio
Pittosporum rhombifolium	Queensland Pittosporum
P. tobira	Japanese Pittosporum
Podocarpus macrophyllus	Japanese Yew
Prunus laurocerasus	English Laurel
Prunus spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Pseudotsuga menziesii	Douglas Fir**
Pyraecantha coccinea	Firethorn (Scarlet)
P. fortuneana	Firethorn
P. korizumii	Firethorn
Pyrus spp.	Breadford Pear spp.
Quercus shumardii	Oak, Shumard's Red
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorne
Raphiolepis umbellata	Yedda Hawthorne
Rhododendron	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'P.J.M.,' 'Delaware Valley White,' 'Roseum elegans,' 'Flame Creeper,' 'Girard Crimson,' 'Wakeleibis,' 'White,' 'Gumbo,' 'George L. Tabor'
Rosa banksiae	Lady Bank's Rose
Rosmarinus officinalis	Rosemary*
Rudbeckia spp.	Black-Eyed Susan; Goldsturm
Santolina virens	Santolina
Saxifraga spp.	Saxifrage; Purple Dome
Scabiosa spp.	Pincushion Flower
Sedum album	Stonecrop
Sedum caudicatum	Stonecrop; Lidakense
Sedum dasylphyllum	Stonecrop
Sedum spurtium	Stonecrop; Dragon's Blood
Spiraea bumalda	Spiraea; Anthony Waterer
Syzygium paniculatum	Australian Brushcherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Teucrium spp.	Germander
Thalictrum dipterocarpum	Meadow Rue
Thuja occidentalis	American Arborvitae

SCIENTIFIC NAME	COMMON NAME
Trachelospermum asiaticum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Veronica japonica	Veronica, Speedwell; Sunny Border
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. suspensum	Arrowwood Viburnum
V. tinus	Laurustrinus
V. trilobium	Cranberry Bush
V. wrightii	Leathereaf Viburnum
Vinca major	Greater Periwinkle
Vinca Vinca Minor	Common Periwinkle
Vinca Vitis spp.	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca
Y. filametosa	Adam's Needle, Yucca

*Do not use on food-producing trees, vines or plants.
**Not for use on container grown plants.

STORAGE AND DISPOSAL:
Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry, cool, locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND WARRANTY
The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Easy Gardener Products, Inc. or the Seller. To the extent consistent with applicable law, Buyer and User agree to hold Easy Gardener Products, Inc. harmless for any claims relating to such factors.

Easy Gardener Products, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above. To the extent consistent with applicable law, Buyer and User agree to hold Easy Gardener Products, Inc. harmless for any claims relating to such factors.

EASY GARDENER PRODUCTS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

To the extent consistent with applicable law, in no event shall Easy Gardener Products, Inc. be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY FOR EASY GARDENER PRODUCTS AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES (INCLUDING CLAIMS BASED ON BREACH OR WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF EASY GARDENER PRODUCTS OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Easy Gardener Products, Inc. and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY, which may not be modified except by written agreement signed by a duly authorized representative of Easy Gardener Products, Inc.

© Copyright Easy Gardener Products, Inc. 2012

"Blocks Weeds Before They Start" is a trademark and Easy Gardener and WeedBlock are registered trademarks of Easy Gardener Products, Inc.

EPA 111611/Rev B

WEEDBLOCK® PRE-EMERGENT Northern

For Selective Pre-Emergence Control of Grass and Broadleaf Weeds in Lawns and Ornamental Turf (excluding golf course putting greens), and Landscape Ornamentals (including established perennials and wildflower plantings)

Stops Weed Grasses Before They Grow
Prevents Crabgrass, Annual Bluegrass, Foxtail & More

ACTIVE INGREDIENT:	
Proflamime**	0.22%
OTHER INGREDIENTS:	99.78%
TOTAL:	100.00%
*CAS No. 29091-21-2	

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for precautionary statements

EPA Reg. No. 74468-6-46260

EPA Est. No. 81245-MN-1

NET CONTENT: 4.65 Lbs.

Manufactured for:
Easy Gardener Products, Inc.
Waco, TX 76710

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

WeedBlock Pre-Emergent Northern is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting germination, not root development.

For landscape ornamentals, ground covers, ornamental grasses, shrubs, trees and turf grasses as listed on this label.

WHEN TO APPLY WEEDBLOCK PRE-EMERGENT NORTHERN

Apply WeedBlock Pre-Emergent Northern during the growing season around established plants, transplants and lawns. **Do not apply WeedBlock Pre-Emergent Northern over newly planted flower seeds or over newly overseeded lawns.** It can be used after flowering plants have germinated and are 2-3 inches tall,