

Transferred from LT BioSyn, Inc. EPA Reg. No. 72639-12

**CP Bio, Inc. MegaGro™**  
**Reduces transplant shock and promotes plant growth**  
**For Agricultural Use**

Not for Use, Sale or Resale in California

**ACTIVE INGREDIENTS:** ..... By wt  
 2-Indolebutyric acid (IBA) ..... 0.85%  
 Cytokinin, as Kinetin ..... 0.15%  
**OTHER INGREDIENTS:** ..... 99.00%  
**TOTAL** ..... **100.00%**

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**FIRST AID**  
 If in eyes: • Hold eye open and rinse slowly and gently with water for 15-20 minutes.  
 • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.  
 • Call a poison control center or doctor for treatment advice.

**HOT LINE**  
 Have the product container or label with you when calling a poison control center or doctor or going for treatment. For general product information, call CP Bio, Inc. at (909) 348-5133 between the hours of 9 a.m. - 4 p.m. Pacific Time.

**PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using toilet.

**ENVIRONMENTAL HAZARDS**  
 Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

EPA Reg. No. 88031-14 EPA Est. No. 88031-CA-001  
 CP Bio, Inc. Batch No.  
 4802 Murrieta St.  
 Chino, CA 91710 USA  
 Email: info@cpbio.com

Net Contents: One gallon (3.875 L)

Department of Agriculture  
STATE OF HAWAII  
**LICENSED**  
PERIOD 2010-2012 LIC. NO.

Department of Agriculture  
STATE OF HAWAII  
**LICENSED**  
PERIOD 2013-2015 LIC. NO.  
**8887.2**

**DISC.**

Crop	Dilute	Concentrate
Vegetables, Field Crops	15	5
Small Fruits, Vines, Miscellaneous	150	15
Tree crops	400	20
Citrus	800	20

Depending upon the equipment used, and the specific crop, the spray volume applied per acre will differ. Special considerations: MegaGro™ compatibility with other agricultural products has not been fully investigated. Compatibility of MegaGro™ with other products requires testing for crop safety and performance prior to large-scale use. Products mixed with MegaGro™ must be acidic (pH less than 7). Do not mix MegaGro™ with any application having a pH greater than 7. Repeat application may be necessary if it rains within 2 hours after application. Depending upon the equipment used and specific crop, spray volume applied per acre will differ. Apply sufficient water volume to ensure thorough coverage.

**APPLICATION INSTRUCTIONS**  
**TO REDUCE TRANSPLANT SHOCK AND PROMOTE NEW PLANT GROWTH**

Crop	Amount of MegaGro™ (fl. oz.)	Application Timing	Remarks
Alfalfa, Clover, Hay	2 fl. oz. per acre*	Seedling Alfalfa, Clover, Hay apply at 2-4 trifoliate stage. Established crops at green-up.	Promotes new root development and growth.
Brassica Vegetables such as: Broccoli, Cauliflower, Cabbage & Mustard Greens	Drench application in transplant water at 1.3 fl. oz. per 10 gallons water. Foliar application: 2 fl. oz. per acre*	1st: At time of transplant. 2nd: 10-14 days after first application. 1st: At 2-4 true leaf stage. 2nd: 10-14 days after first application.	Foliar application: Apply to achieve full coverage use a nonionic surfactant for hard to wet crops such as cabbage.
Canola (Oil)	2 fl. oz. per acre*	1st: At 2-4 leaf stage.	Tank mix with Glyphosate products registered for use on Roundup Ready® canola.
Citrus Fruits such as: Sweet orange, Lemon and Grapefruit	13 fl. oz. in 100 gallons water	Apply when fruit are 5 mm in size. Make additional applications if needed.	Thoroughly apply spray.

Crop	Amount of MegaGro™ (fl. oz.)	Application Timing	Remarks
Corn (Sweet, Field and Popcorn)	2 fl. oz. per acre*	1st: At 2-4 leaf stage.	Tank mix with Glyphosate products registered for use on Roundup Ready® canola.
Cotton	2 fl. oz. per acre*	1st: At 2-4 true leaf stage. 2nd: 10-14 days after first application.	Can be tank mixed with Glyphosate products registered for use on Roundup Ready® cotton.
Cucurbit Vegetables such as: Cucumbers, Melon, Watermelon and Honeydew	2 fl. oz. per acre*	1st: If seeded, apply at 2-4 true leaf stage. 2nd: 10-14 days after first application.	Enhances root growth.
Fruiting Vegetables such as: Tomatoes, Pepper and Eggplant	2 fl. oz. per acre*	1st: At 2-4 true leaf stage. 2nd: 10-14 days after first application.	Increases potential to enhance berry size.
Grape	4-6 fl. oz. in 100 gallons water	1st: Apply when grapes are 2-3 mm in size. 2nd: 10-14 days after first application. 3rd: 45 days prior to harvest. 4th: 30 days prior to harvest.	Increases potential to enhance berry size.
Leafy Vegetables such as: Celery, Head Lettuce, Leaf Lettuce and Spinach	2 fl. oz. per acre*	1st: At 2-4 true leaf stage. 2nd: 10-14 days after first application.	Can be tank mixed with Glyphosate products registered for use on Roundup Ready® soybeans.
Legume Vegetables (Succulent or dry) such as: Soybeans, Peas, all types of beans	2 fl. oz. per acre*	At 2-5 trifoliate leaf stage.	Can be tank mixed with Glyphosate products registered for use on Roundup Ready® soybeans.
Peanut	2 fl. oz. per acre*	1st: At 2-4 true leaf stage. 2nd: 10-14 days after first application.	Foliar application: Apply thoroughly until dripping.
Tuber Vegetables such as: Potatoes, Sweet potato, Yam	2 fl. oz. per acre*	1st: At 2-4 true leaf stage. 2nd: 10-14 days after first application.	Foliar application: Apply thoroughly until dripping.
Root Vegetables such as: Carrots, Radish, Turnip, Ginseng, Horseradish, Parsley (tump-rooted) and Sugar beet	2 fl. oz. per acre*	1st: At 2-4 true leaf stage. 2nd: 10-14 days after first application.	Foliar application: Apply thoroughly until dripping.

Crop	Amount of MegaGro™ (fl. oz.)	Application Timing	Remarks
Apple/Stone Fruits such as: Apple, Apricot, Cherry, Plum, Peach and Peach	13 fl. oz. in 100 gallons water	Apply when fruit are 5 mm in size. Make additional applications if needed.	
Rice	2 fl. oz. per acre*	1st: At 4-5 leaf stage.	
Strawberry	1.3 fl. oz. in 10 gallons water	1st: Spray immediately after transplant. 2nd: 10-14 days after first application.	
Soybeans	2 fl. oz. per acre*	At 2-5 trifoliate leaf stage.	Tank mix with Glyphosate products registered for use on Roundup Ready® soybeans.
Sorghum	2 fl. oz. per acre*	At 2-6 leaf stage.	
Tobacco	2 fl. oz. per acre*	At 2-4 leaf stage.	

\*If application spray volume is greater than 15 gallons per acre, use the dilution rate of 13 fl. oz. per 100 gallons water. Thorough spray coverage is necessary.

**STORAGE AND DISPOSAL**  
 Do not contaminate water, food or feed by storage or disposal.  
**PESTICIDE STORAGE:**  
 MegaGro™. Store in its original container in a cool, dry, locked place out of the reach of children and out of direct sunlight.  
**PESTICIDE DISPOSAL:**  
 Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
 Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent). Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**CP Bio, Inc.**  
**4802 Murrieta St.**  
**Chino, CA 91710 USA**  
**Tel: (909) 348-5133 Fax: (909) 348-5135**  
**Email: info@cpbio.com**

MegaGro™ is a trademark of CP Bio, Inc.  
 Roundup Ready® is a registered trademark of the Monsanto Company.

**LIMITED WARRANTY AND DISCLAIMER**  
 NOTICE: CP Bio, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use. Buyer assumes all risks of use and handling which is a variance in any way with the directions herein. CP Bio, Inc. makes no other express or implied warranty of fitness or merchantability. In no case shall CP Bio, Inc. or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. CP Bio, Inc. and Seller offer this product and the Buyer and user accept it, subject to the foregoing Limited Warranty and Disclaimer which may be varied only by agreement in writing signed by a duly authorized representative of CP Bio, Inc.

**CP Bio, Inc. MegaGro™**  
**Reduces transplant shock and promotes plant growth**  
**For Agricultural Use**

Not for Use, Sale or Resale in California

**ACTIVE INGREDIENTS:** ..... By wt  
 2-Indolebutyric acid (IBA) ..... 0.85%  
 Cytokinin, as Kinetin ..... 0.15%  
**OTHER INGREDIENTS:** ..... 99.00%  
**TOTAL** ..... **100.00%**

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**FIRST AID**  
 If in eyes: • Hold eye open and rinse slowly and gently with water for 15-20 minutes.  
 • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.  
 • Call a poison control center or doctor for treatment advice.

**HOT LINE**  
 Have the product container or label with you when calling a poison control center or doctor or going for treatment. For general product information, call CP Bio, Inc. at (909) 348-5133 between the hours of 9 a.m. - 4 p.m. Pacific Time.

**PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using toilet.

**ENVIRONMENTAL HAZARDS**  
 Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

EPA Reg. No. 88031-14 EPA Est. No. 88031-CA-001  
 CP Bio, Inc. Batch No.  
 4802 Murrieta St.  
 Chino, CA 91710 USA  
 Email: info@cpbio.com

Net Contents: One gallon (3.875 L)

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**  
 Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear:  
 • Long sleeved shirt and long pants.  
 • Chemical resistant gloves Category A, such as butyl rubber ≥ 14 mils, or natural rubber ≥ 14 mils, or neoprene rubber ≥ 14 mils or nitrile rubber ≥ 14 mils and shoes plus socks.  
 Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**PHYSICAL OR CHEMICAL HAZARDS**  
 Combustible. Do not use or store near heat or open flame.  
 For chemical spill, leak, fire or exposure, call CP Bio, Inc. at (909) 348-5133

**USER SAFETY RECOMMENDATIONS**  
 Users should:  
 • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.  
 • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**DIRECTIONS FOR USE**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the state or tribal agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**  
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.  
 PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:  
 • Coveralls - Chemical resistant gloves Category A, such as butyl rubber ≥ 14 mils, or natural rubber ≥ 14 mils, or neoprene rubber ≥ 14 mils or nitrile rubber ≥ 14 mils - Shoes plus socks.

**GENERAL INFORMATION**  
 MegaGro™ contains two (2) different plant growth regulators (PGR) that synergistically stimulate early and improved root development. In addition, MegaGro™ contains a mixture of vitamins that aid in successful transplanting and plant growth.

**USE DIRECTIONS FOR CHEMIGATION**  
 MegaGro™ Apply through fixed or standing irrigation systems or thru foliar applications. Foliar applications are preferred. Apply this product only through the following types of irrigation systems:  
 1. Sprinklers including big gun, solid set or hand move irrigation systems.  
 2. Calibrated overhead watering booms  
 3. Drip (or micro sprinkler) irrigation systems

Before applying this product through any type of irrigation system, perform a small-scale trial to determine if product performance and phytotoxicity results are acceptable. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems), used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

**CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS**  
 Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.  
 Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Agitate the pesticide supply tank throughout the application of MegaGro™. Except for turfgrass, apply MegaGro™ at the rate of 20 fl. oz. per acre at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop. Fill the supply tank one-half full with water, add the appropriate amount of MegaGro™ to the tank and finish filling the tank with water.

**DRIP/TRICKLE OR SPRINKLER CHEMIGATION**  
 The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment (this statement only applies to sprinkler chemigation).

Agitate the pesticide supply tank throughout the application of MegaGro™. Except for turfgrass, apply MegaGro™ at the rate of 20 fl. oz. per acre at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop. Fill the supply tank one-half full with water, add the appropriate amount of MegaGro™ to the tank and finish filling the tank with water.