Altosid® SBG II

SINGLE BROOD GRANULE

ZOËCON®

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.
8910.49

PREVENTS ADULT MOSQUITO EMERGENCE WHEN APPLIED PRE OR POST-HATCH

(including those which may transmit West Nile virus)

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACTIVE INGREDIENT: (S)-Methoprene (CAS# 65733-16-6) 0.3%
OTHER INGREDIENTS: 99.7%
TOTAL: 100.0%

NET WT 40 LBS (18.14 kg)

PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS AND DOMESTIC ANIMALS – CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID	
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of rinsate or equipment washwaters.

INTRODUCTION

Altosid® SBG II Single Brood Granule (Altosid® SBG II) slowly releases effective levels of (S)-methoprene insect growth regulator to prevent the emergence of adult mosquitoes from treated sites. Apply **Altosid® SBG II** up to 7 days prior to flooding (i.e. "pre-hatch") in known mosquito breeding areas, or post-hatch to sites containing mosquito larvae.

Treated larvae continue to develop normally to the pupal stage where they die. **Altosid® SBG II** prevents the emergence of adult mosquitoes from treated sites.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

APPLICATION DIRECTIONS

Altosid® SBG II is designed to control single broods of mosquito larvae in water depths up to 15 inches. Under typical environmental conditions, one application will control for up to 7 days. Applications can be made pre-hatch to known mosquito breeding areas, or post-hatch to sites containing mosquito larvae. Pre-hatch treatments should be applied in the low areas of breeding sites to maintain continuous control as the site floods and dries. Repeat application cycle throughout the mosquito season.

NOTE: The active ingredient (S)-methoprene has no effect on mosquitoes that have reached the pupal or adult stage prior to treatment.

Lot No.

APPLICATION RATES

CENTRAL Garden & Pet

Select the target mosquito species and corresponding application rate per acre.

Aedes, Ochlerotatus, Anopheles and Psorophora spp. :

- Apply **Altosid® SBG II** at 3 – 7 lbs per acre

Culex and Culiseta spp. :

- Apply **Altosid® SBG II** at 10 – 20 lbs per acre

For smaller application sites, apply **Altosid® SBG II** at 0.35 oz to 0.70 oz per 100 square feet of water surface.

APPLICATION SITES

Apply **Altosid® SBG II** treatments pre-hatch or post-hatch as directed above to temporary and permanent water sites which support mosquito larval development. Examples of typical sites are: freshwater and salt water marshes, cattail marshes, woodland pools and meadows, grassy swales, floodplains, irrigated croplands, pastures and rangeland (without removal of livestock), vineyards, rice fields (including wild rice), fruit and nut orchards, berry fields and bogs, dredge spoil sites, storm water retention basins, storm drains, retention and detention ponds, drainage areas, roadside ditches, tree holes, tires, water-holding receptacles, ornamental ponds and fountains, animal watering troughs and ponds, fish ponds, flooded crypts, transformer vaults, abandoned swimming pools, artificial water-holding containers, and other natural and manmade depressions where mosquitoes may breed.

NOTE: Application of **Altosid® SBG II** to sites subject to water flow or exchange will diminish the product's effectiveness.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag. Then offer for recycling if available or dispose of empty bag in trash or in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND CONDITIONS OF SALE

Seller makes no warranty, express or implied, concerning the use and handling of this product other than indicated on the label. To the fullest extent permitted by law, buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

Always read the label before using this product.

For information, call 1-800-248-7763 or visit our web site: **www.altosid.com**

Manufactured for
Central Garden & Pet Company
1501 E. Woodfield Road, 200W
Schaumburg, Illinois 60173, U.S.A.

EPA Reg. No. 75318-8-89459
EPA Est. No. 2724-TX-1

U.S. Patent Pending

ALTOSID and ZOËCON with design are registered trademarks of Wellmark International. Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company. ©2006, 2014 Wellmark International
300516478