



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

8910.91

ADAMS™
Flea & Tick Collar for Dogs



READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID	
If swallowed	Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 800-234-2269 for emergency medical treatment information.

Note to Physicians and Veterinarians:

This product contains tetrachlorvinphos (an organophosphate) which is a cholinesterase inhibitor. Cholinergic symptoms may include salivation, miosis, incoordination, muscle fasciculation and/or weakness, vomiting and diarrhea. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine. Call 800-234-2269 or your local Poison Control Center for further information.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REMOVE COLLAR FROM PACKAGE, UNROLL AND STRETCH TO ACTIVATE INSECTICIDE GENERATOR. As the collar begins to work, a fine white powder will appear on the surface.

Do not use on puppies under 6 weeks of age. Do not unroll collar until ready to use. Place the Adams™ Flea & Tick Collar for Dogs around the dog's neck, adjust for proper fit, and buckle in place. The collar must be worn loosely to allow for growth of the dog and to permit the collar to move about the neck. Generally, a properly fitted collar is one that, when fastened, will snugly slide over the dog's head. Leave 2 or 3 inches on the collar for extra adjustment and cut off and dispose of the extra length.

Consult a veterinarian before using this product on debilitated, aged, or medicated, animals. If used on pregnant or nursing dogs, do not replace collar until the puppies are at least 6 weeks old. Sensitivity may occur after using ANY pesticide product for pets. Some animals may become irritated by any collar if it is applied too tightly. If this occurs, loosen the collar. If irritation continues, remove the collar from the dog. If signs of sensitivity occur, remove the collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Do not use this product on dogs simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea and tick collars may be immediately replaced.

This collar is intended for use only as an insecticide generator.

The collar will begin to kill fleas and ticks immediately. Its full protection against harder to kill ticks will be built up within a few days after being placed on the pet. For continuous flea and tick protection under normal conditions, replace the collar every seven months. The collar kills ticks for 5 months and aids in their control an additional 2 months. Under conditions where dogs are exposed to severe flea and tick infestations, it may be necessary to replace the collar more frequently, however, do not replace the collar more frequently than every 5 months. Wetting will not impair the collar's effectiveness or the pet's protection. If the dog goes swimming or is out in the rain, it is not necessary to remove the collar. The Adams™ collar may be worn with a regular collar. It is recommended that the pet's bedding or sleeping quarters be treated for flea and tick control with products registered for this use such as Adams™ Flea & Tick Home Spray.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. **Pesticide Storage:** Store this product in a cool, dry place that is inaccessible to children. **Pesticide Disposal:** If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured for:

Central Garden and Pet Company
1501 East Woodfield Road 200W
Schaumburg, Illinois 60173

For information or comments, call 800-234-2269
Visit our Web site at adamsfleacontrol.com

Item: 100524188 • 15-10290 • RM: 300519529

EPA REG. No. 2596-84-89459

EPA Est. No. 2724-TX-1

©2004-2014, 2015 Farnam Companies, Inc.

Central Garden & Pet with design is a trademark of Central Garden & Pet Company. All other trademarks are the property of their respective owners.

ADAMS™
Flea & Tick Collar for Dogs

7 MONTH
Protection



Fits Up To
26" Necks

7 MONTH
Protection

ADAMS™

**Flea & Tick
Collar
for Dogs**



**Water
Resistant**

KILLS & REPELS:



fleas



ticks

CAUTION
**DO NOT LET CHILDREN
PLAY WITH THIS COLLAR**

See Back Panel for Additional Precautionary Statements

ACTIVE INGREDIENT:	
Tetrachlorvinphos	14.55%
OTHER INGREDIENTS:	85.45%
TOTAL:	100.00%

Net contents: 1 collar
Net weight: 1.2 OZ (34 g)



0 39079 09318 7

ADAMS™
Flea & Tick
Collar for Dogs

7 MONTH
Protection

7 MONTH
Protection

ADAMS™
Flea & Tick
Collar for Dogs

ADAMS™
Flea & Tick
Collar for Dogs