



DELTA CUP L®

**A Protective and Curative Fungicide
For Use On:**

Citrus, Vegetables, Tree Crops, Small Fruits, Vines and Field Crops

This product is in a liquid form miscible with water and includes 236 g of copper sulfate per liter which corresponds to 60.2 g of metallic copper per liter.



LICENSED

PERIOD **2013-2015** LIC. NO. **8940.1**

Active Ingredient:	
Copper Sulfate Pentahydrate*	19.6%
CAS No. 7758-99-8	
Other Ingredients:	80.4%
Total:	100%

*Copper as metallic: 5%

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear the following:

- Long-sleeved shirt
- Long pants
- Shoes plus socks
- Chemical resistant gloves
- Protective Eyewear

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber ≥14 mils, or nitrile rubber ≥14 mils. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.

Users should wash the outside of gloves before removing.

Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, and then continue rinsing. Call poison control center or doctor for treatment advice.

If on Skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378, 7:30 am to 3:30 pm M-F Pacific Time (NPIC Web site: www.npic.orst.edu.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS:

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirement for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Notify workers of application by warning them orally.

Do not enter or allow worker entry into treated areas during the Restricted-Entry Interval (REI) of 48 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: Coveralls, shoes plus socks, chemical-resistant gloves made of any waterproof material, and protective eyewear.

Not for use in greenhouses.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses that are NOT within the scope of the Worker

Protection Standard for Agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

Spray Drift Management. A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size. Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed. Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are no sensitive areas within 250 feet downwind.

Temperature Inversions. If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements. Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have more stringent regulations, they must be observed.

Equipment. Ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For ground boom application:

- Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Method of application. Product can be applied with any type of application equipment that gives uniform coverage of all foliage, including aerial, ground and low volume sprayers. For every 2 pints of product, apply in 100 gallons of water unless otherwise noted in the "Application Rates" table. Stir or agitate thoroughly before use. Spray carefully for complete plant coverage.



Delta Agrochemicals Co.

15 A, Road No. 6, Maadi, Cairo, Egypt Tel.: (+202) 23787797 - (+202) 23809702

Mob.: (+2) 0100 1605789 Fax: (+202) 23787796

E-mail: info@deltaagro.com web: www.deltaagro.com

Made in Egypt

Lot Number:

Net Contents:

EPA Reg. No. 88633-2

EPA Est. No. 88633-EGP-001

Application Rates

Crop: Pest	Disease	Season	Use Rates in pints per acre	Maximum Application Rate (pints per acre)	Maximum Annual Rate (pints per acre)	Minimum Retreatment Interval	Use Notes
Field Crops							
Alfalfa	Cercospora Leaf Spot, Leptosphaerulina Leaf Spot	Growing Season	1.5 - 2.0	2.0	8.0	30 days	Apply 10 - 14 days before each harvest or earlier if disease threatens. Note. Spay injury may occur with sensitive varieties such as Lohontan.
Corn (Field Corn, Popcorn, Sweet Corn)	Bacterial Stalk Rot	Growing Season	1.5 - 2.0	2.0	8.0	7 days	Begin treatment when disease first appears and every 7 to 10 days or less frequently depending on environmental conditions and disease severity. Use the higher rates and shorter spray intervals when conditions favor disease.
Potato	Early Blight, Late Blight	Growing Season	1.5 - 2.0	2.0	8.0	5 days	Apply 1.5 to 2 pints at 7 to 10 day intervals or less frequently depending on environmental conditions and disease severity starting when plants are 2 to 6 inches high in locations where disease is light. Apply up to 2.0 pints per acre when disease is more severe.
Peanut	Cercospora leaf Spot	Growing Season	1.5 - 2.0	2.0	14.0	7 days	Begin spraying at 35 to 40 days after planting or when disease symptoms first appear and repeat at 10 - 14 day intervals or less frequently depending on environmental conditions and disease severity. Reduce spray to 7 day intervals during humid weather. Use the higher rates when
Sugar Beet	Cercospora Leaf Spot	Growing Season	1.5 - 2.0	2.0	10.0	10 days	Begin application when conditions favor disease development and repeat at 10 - 14 day intervals or less frequently, depending on environmental conditions and disease severity. Use the high rates when conditions favor disease.
Wheat, Barley, Oats, Millet, Sorghum, Rye	Helminthosporium Spot, Blotch Septoria Leaf Blotch	Growing Season	1.5 - 2.0	2.0	4.0	10 days	Make first application at early heading and follow with second spray 10 days later. Use the higher rates when conditions favor disease. For Wheat, Delcup can be applied as a foliar application for early season disease control and again at early heading and followed with another application 10 days later
Small Fruits							
Blackberry (Aurora, Boysen, Cascade, Chehalem, Logan, Marion, Santiam, Thornless Evergreen)	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	Fall, Late Dormant	2.0 - 4.0	4.0	4.0	7 days	Make fall application after harvest. Apply delayed dormant spray after pruning/ training in the spring. If needed, agricultural type spray oil may be added.
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	Growing Season	2.0 - 4.0	4.0	8.0	7 days	Apply when leaf buds begin to open and repeat when flower buds show white. If needed, agricultural type spray oil may be added.
Blueberries	Bacterial Canker	Fall, Late Dormant	2.0 - 4.0	4.0	8.0	30 days	Make first application before fall rains and a second application 4 weeks later. Use the higher rates when conditions favor disease.
	Fruit Rot, Phomopsis Twig Blight	Late Dormant	2.0 - 4.0	4.0	8.0	10 days	Dormant Application: Begin application when bloom buds begin to swell. Make additional applications at 10 - 14 day intervals or less frequently depending on environmental conditions and disease severity before blooms open.
Cranberry	Fruit Rot	Growing Season	4.0	4.0	12.0	10 days	Make first application in the late bloom. Apply one or two additional applications at 10 - 14 day intervals or less frequently depending on environmental conditions and disease severity.
	Rose Bloom	Growing Season	4.0	4.0	12.0	10 days	Apply three sprays on 10 - 14 day schedule or less frequently depending on environmental conditions and disease severity as soon as symptoms are observed.
	Bacterial Stem Canker	Post harvest / dormant	4.0	4.0	12.0	10 days	Apply post harvest and again in spring at bud swell. Apply one or two additional applications at 10 - 14 day intervals or less frequently depending on environmental conditions and disease severity.
	Leaf Blight, Red Leaf Spot, Stem Blight, Tip Blight	Late Dormant	4.0	4.0	12.0	10 days	Apply delayed dormant spray in the spring. Repeat at 10 - 14 day intervals or less frequently depending on environmental conditions and disease severity through pre bloom
Currant Gooseberry	Anthracnose Leaf Spot	Growing Season	4.0	4.0	12.0	10 days	Make initial application after first leaves have expanded. Continue on a 10 - 14 day schedule or less frequently depending on environmental conditions and disease severity during wet conditions in the spring. Make an additional application after harvest
Raspberry	Anthracnose, Cane Spot, Leaf Spot, Pseudomonas Blight, Purple Blotch, Yellow Rust	Fall, Late Dormant	2.0 - 4.0	4.0	8.0	7 days	Make fall application after harvest. Apply delayed dormant spray after training in the spring. If needed, agricultural type spray oil may be added.
	Anthracnose, Cane Spot, Leaf Spot, Purple Blotch, Yellow Rust	Growing Season	2.0 - 4.0	4.0	8.0	7 days	Apply when leaf buds begin to open and repeat when flower buds show white. If needed, agricultural type spray oil may be added.
Strawberry	Angular Leaf Spot (Xanthomonas), Leaf Blight, Leaf Scorch, Leaf Spot	Growing Season	1.5 - 2.0	2.0	14.0	7 days	Begin application when plants are established and continue on a weekly schedule throughout the season. Apply in at least 20 gallons of water. Use the higher rates when conditions favor disease.
Tree Crops							
Almonds only	Bacterial Blast	Growing Season	3.0	3.0	9.0	14 days	Almonds Only: For bacterial blast control in sprinkler irrigated orchards or where disease is severe. Apply 1.50 pts. per acre post bloom at 2 week intervals or less frequently depending on environmental conditions and disease severity or just before sprinkling
Almonds, Apricots, Cherry, Plum, Prune	Bacterial Blast (Pseudomonas), Bacterial Canker, Coryneum Blight (Shot Hole)	Fall, Late Dormant	3.0 - 6.0	6.0	12.0	14 days	Make first application before fall rains and a second at dormant. Use the higher rates when conditions favor disease. If needed, agricultural type spray oil may be added. For Cherries: Where disease is severe, an additional application shortly after harvest may be required.
	Blossom Brown Rot, Coryneum Blight (Shot Hole)	Growing Season (Early Spring)	3.0 - 4.0	4.0	12.0	5 days	Apply during early bloom. Use the higher rates when rainfall is heavy and disease pressure is high.
	Black Knot (Plum)	Growing Season	3.0 - 4.0	4.0	12.0	7 days	Make an application at bud swell up to early bloom for early season disease suppression. Apply before full bloom. Use the higher rates when rainfall is heavy and disease pressure is high.
	Cherry Leaf Spot (Sour Cherries Only)	Growing Season	3.0 - 4.0	4.0	12.0	7 days	Apply at petal fall as well as 1 to 2 times after petal fall. Use the lower rates where disease infection is light and use the higher rates for a dormant application or where disease infection is moderate to heavy. Do not apply to sweet cherry or the English Morello variety as severe injury will result.
Apple	Anthracnose, Blossom Blast, European Canker (Nectria), Shoot Blast (Pseudomonas)	Fall	4.0 - 6.0	6.0	12.0	Only 1 application per season permitted	Apply before fall rains. Use the higher rates when conditions favor disease.
	Apple Scab, Fire Blight	Growing Season	4.0 - 6.0	6.0	6.0	Only 1 application per season permitted	Make application between silver tip and green tip. Apply as a full cover spray for early season disease suppression.
	Apple Scab	Growing Season	3.0 - 4.0	4.0	16.0	5 days	Extend Spray schedule where fruit finish is not a concern. Continued application may be made at 5 to 7 day intervals or less frequently depending on environmental conditions and disease severity between 1/2 inch green tip and first cover spray.
	Fire Blight	Growing Season	3.0 - 4.0	4.0	16.0	5 days	
Collar Rot, Crown Rot	Dormant spring / fall	4.0	4.0	8.0	Only 1 application per season permitted	Mix in 100 gallons of water. Apply 4 gallons of suspension as a drench on the lower trunk area of each tree. Apply in early spring or in fall after harvest for best results. Do not apply to foliage or fruit. Note, Do not use if soil pH is below 5.5 since copper toxicity may result.	
Avocado	Anthracnose, Blotch Scab	Growing Season	3.0 - 4.0	4.0	24.0	14 days	Apply when bloom buds begin to swell and continue application at monthly intervals for five to six applications. Use the higher rates when conditions favor disease.
Banana	Sigatoka (Black and Yellow)	Growing Season	4.0	4.0	40.0	7 days	Apply by air in 3 gallons of water per acre. If needed agricultural type spray oil may be added. Apply on a 14 day schedule or less frequently depending on environmental conditions and disease severity throughout the wet season. Apply at 21 day intervals or less frequently depending on environmental conditions and disease severity during dry periods.

	Black Pitting	Growing Season	4.0	4.0	12.0	7 days	Mix in 100 gallons of water. Apply to the fruit stem and the basal portion of the leaf crown. Apply during the first and second weeks after fruit emergence.
Cacao	Black Pod	Growing Season	3.0 - 4.0	4.0	20.0	14 days	Begin application at the start of the rainy season and continue while infections conditions persist. Apply 2.0 - 3.0 pts. at 14 to 21 day intervals or less frequently depending on environmental conditions and disease severity.
Coffee	Coffee Berry Disease (<i>Colletotrichum coffeanum</i>)	Growing Season	3.0 - 4.0	4.0	20.0	21 days	Apply first spray after flowering and before onset of long rains and then at 21 - 28 day intervals or less frequently depending on environmental conditions and disease severity until picking. Use the higher rates when conditions favor disease.
	Bacterial Blight (<i>Pseudomonas syringae</i>)	Growing Season	3.0 - 4.0	4.0	20.0	14 days	Begin spray program before the onset of long rainy periods and continue through the rainy season at 14 to 21 day intervals or less frequently depending on environmental conditions and disease severity. The critical time for spraying to control this disease is just before, during, and after flowering(s), especially when coinciding with wet weather. Use the higher rates when rainfall is heavy and disease pressure is high.
	Leaf Rust (<i>Hemileia vastatrix</i>)	Growing Season	3.0 - 4.0	4.0	40.0	21 days	Apply before the onset of rain and then at 21 day intervals or less frequently depending on environmental conditions and disease severity while the rains continue. Use the higher rates when rainfall is heavy and pressure is high.
	Iron Spot (<i>Cercospora coffeicola</i>), Pink Disease (<i>Corticium salmonicolor</i>)	Growing Season	3.0 - 4.0	4.0	12.0	30 days	Use concentrate of dilute spray. Begin treatment at the start of wet season and continue at monthly intervals for three applications.
Filberts (Only permitted in Washington and Oregon)	Bacterial Blight	Post harvest	6.0 - 7.5	7.5	15.0	14 days	Apply as a post harvest spray. In season of heavy rainfall, apply a second spray when three fourths of the leaves have dropped. Use the higher rates when rainfall is heavy and disease pressure is high. If needed, agriculture type spray oil may be added.
	Eastern Filbert Blight	Growing Season	6.0 - 7.5	7.5	30.0	14 days	Apply as a dilute spray in adequate water for thorough coverage. Make applications starting at bud swell to bud break and continue at 2 week intervals or less frequently depending on environmental conditions and disease severity until early May. Thorough coverage is essential. Use the higher rates when rainfall is heavy and disease pressure is high. If needed, agricultural type spray oil may be added.
Mango	Anthraxnose	Growing Season	4.0 - 6.0	6.0	18.0	30 days	Apply monthly after fruit set until harvest. Use the higher rate when rainfall is heavy and disease pressure is high.
Olives	Peacock Spot, Olive Knot	Fall, Late Dormant	4.0 - 6.0	6.0	12.0	30 days	Make first application before winter rains begin. A second application in early spring should be made if disease is severe. Apply the higher rates for heavy disease pressure or when conditions favor disease development.
Peach, Nectarine	Bacterial Blast (<i>Pseudomonas</i>), Bacterial Canker, Bacterial Spot (<i>Xanthomonas</i>), Coryneum Blight (Shot Hole), Leaf Curl	Fall, dormant	4.0 - 6.0	6.0	12.0	21 days	Make first application before fall rains and a second application at late dormant. For peach leaf curl, late dormant application must be made before leaf buds swell. Use the higher rates when rainfall is heavy and disease pressure is high. If needed, agricultural type spray oil may be added.
	Blossom Brown Rot, Coryneum Blight (Shot Hole), Leaf Curl	Growing Season	3.0 - 4.0	4.0	12.0	5 days	Full cover spray at pink bud. Use the higher rates when conditions favor disease.
	Bacterial Spot	Growing Season	3.0 - 4.0	4.0	24.0	7 days	Post-bloom application applied at first and second cover sprays. Note: Do not spray 3 weeks prior to harvest
Pear	Fire Blight	Growing Season	3.0	3.0	12.0	5 days	Apply at 5 day intervals or less frequently depending on environmental conditions and disease severity throughout the bloom period.
	Blossom Blast (<i>Pseudomonas</i>)	Fall, Late Dormant	4.0 - 6.0	6.0	12.0	Only one application	Apply before fall rains and again during dormancy, before spring growth starts. Use the higher rates when disease pressure is high or when conditions favor disease development.
Pecan	Kernel Rot, Shuck Rot (<i>Phytophthora caracatorum</i>), Zonate Leaf Spot (<i>Cristulariella pyramidalis</i>)	Growing Season	3.0 - 4.0	4.0	16.0	14 days	For suppression, apply in sufficient water to ensure complete spray coverage at 2 - 4 week intervals or less frequently depending on environmental conditions and disease severity, starting at kernel growth and continue until shucks open. Use the higher rates and shorter spray intervals if frequent rainfall occurs.
	Ball Moss, Spanish Moss	Late Dormant	3.0 - 4.0	4.0	8.0	14 days	Apply in 100 gallons of water in the spring when ball moss is actively growing using 1 1/2 gallons of spray per foot of tree height. Make sure to wet ball moss tufts thoroughly. A second application may be required after 12 months.
Pistachio	Botryosphaeria, Panicle and Shoot Blight, Botrytis Blight, Late Blight (<i>Alternaria alternata</i>), Septoria Leaf Blight	Growing Season	3.0 - 4.0	4.0	12.0	14 days	Make initial application at bud swell and repeat on a 14 to 28 day schedule or less frequently depending on environmental conditions and disease severity. If disease conditions are severe, use the higher rates and shorter spray intervals.
Quince	Fire Blight	Growing Season	1.5	1.5	12.0	5 days	Apply at 5 day intervals or less frequently depending on environmental conditions and disease severity throughout the bloom period. Apply in adequate water for thorough coverage.
Walnuts	Walnut Blight	Early Season	3.0 - 6.0	6.0	30.0	7 days	Apply first spray at early pre-bloom prior to, or when catkins are partially expanded. Make additional applications during bloom and early nutlet stage or less frequently, depending on environmental conditions and disease severity when frequent rainfall or extended periods of moisture occur. Thorough coverage of catkins leaves, and nutlets is essential for effective control. Note. Adequate control may not be obtained when copper tolerant species of <i>Xanthomonas</i> bacteria are present.

Bean (Dry Green)	Brown Spot, Common Blight, Halo Blight	Growing Season	1.5 - 2.0	2.0	10.0	7 days	For protective sprays, make first application when plants are 6 inches high, repeat on a 7 - 14 day schedule or less frequently depending on environmental conditions and disease severity. Use higher rates when conditions favor disease.
Beet (Table Beet, Beet Greens)	Cercospora Leaf Spot	Growing Season	1.5 - 2.0	2.0	8.0	10 days	Begin application when conditions first favor disease development and repeat at 10 - 14 day intervals or less frequently depending on environmental conditions and disease severity. Use the higher rates when conditions favor disease.
Carrot	Alternaria Leaf Spot, Cercospora Leaf Spot	Growing Season	1.5 - 2.0	2.0	8.0	7 days	Begin application when disease first threatens and repeat at 7 to 14 day intervals or less frequently depending on environmental conditions and disease severity.
Celery Celeriac	Bacterial Blight, Cercospora Early Blight, Septoria Late Blight	Growing Season	1.5 - 2.0	2.0	12.0	7 days	Begin application as soon as plants are first established in the field, repeating at 7 to 10 day intervals or less frequently, depending on environmental conditions and disease severity.
Crucifers (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collard Greens, Mustard Greens, Turnip Greens) Chinese Cabbage, Kale, Kohlrabi	Black Leaf Spot (<i>Alternaria</i>), Black Rot (<i>Xanthomonas</i>), Downy Mildew	Growing Season	1.5 - 2.0	2.0	12.0	7 days	Begin application after transplants are set in the field or shortly after emergence of field seeded crops or when conditions favor disease development. Apply at 7 - 10 day intervals or less frequently depending on environmental conditions and disease severity. Use the higher rates when conditions favor disease.
Cucurbits (Cantaloupe, Cucumber, Honeydew, Muskmelon, Pumpkin, Squash, Watermelon) Casaba, Chayote, Citron Melon, Gourd, Waxgourd	Alternaria Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Gummy Stem Blight, Watermelon Bacterial Fruit Blotch (suppression)	Growing Season	1.5 - 2.0	2.0	14.0	5 days	Begin application prior to disease development and continue while conditions are favorable for disease development. Repeat at 5 to 7 day intervals or less frequently depending on environmental conditions and disease severity. Use the higher rates when conditions favor disease.
Eggplant	Alternaria Blight, Anthracnose, Phomopsis	Growing Season	1.5 - 2.0	2.0	12.0	7 days	Begin application prior to development of disease symptoms. Repeat spray at 7 to 10 day intervals or less frequently depending on environmental conditions and disease severity.
Okra	Anthraxnose, Bacterial Leaf Spot, Leaf Spot, Pod Spot	Growing Season	1.5 - 2.0	2.0	12.0	5 days	Begin treatment when disease first threatens and repeat every 5 - 10 days or less frequently depending on environmental conditions and disease severity. Use the higher rates and shorter spray intervals when conditions favor disease.
Onion, Garlic	Bacterial Blight, Downy Mildew, Purple Blotch	Growing Season	1.5 - 2.0	2.0	12.0	7 days	Begin when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals or less frequently depending on environmental conditions and disease severity.
Pepper	Anthraxnose, Bacterial Leaf Spot, Cercospora Leaf Spot	Growing Season	1.5 - 2.0	2.0	20.0	3 days	Begin application when conditions first favor disease development and repeat at 7 - 10 day intervals or less frequently depending on environmental conditions and disease severity. Use the higher rates when conditions favor disease.
Spinach	Anthraxnose, Blue Mold, Cercospora Leaf Spot, White Rust	Growing Season	1.5 - 2.0	2.0	10.0	7 days	Begin application when disease first appears or wet conditions favor disease development. Repeat at 7 to 10 day intervals or less frequently depending on environmental conditions and disease severity. Use the higher rates when conditions favor disease.
Tomato (Fresh Market)	Anthraxnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	Growing Season	1.5 - 2.0	2.0	20.0	3 days	Begin application when disease first threatens and repeat at 5 to 10 day intervals or less frequently depending on environmental conditions and disease severity. Use the higher rates when conditions favor disease.
Watercress	Cercospora Leaf Spot	rowing Seas	1.5 - 2.0	2.0	8.0	7 days	Begin applications when plants are first established in the field, repeating at 7 - 14 day intervals or less frequently, depending on environmental conditions and disease severity.