

AGRI-FOS[®]

400

SYSTEMIC FUNGICIDE FOR THE EFFECTIVE CONTROL OF VARIOUS PLANT DISEASES INCLUDING BLACK SPOT OR SCAB IN APPLE, ROOT ROT IN AVOCADO, BUD ROT AND NUT FALL IN COCONUT, ROOT ROT IN CITRUS AND CUCURBITS, DOWNY MILDEW IN CUCURBITS, GRAPE, LETTUCE, AND ONION, ANTHRACNOSE IN MANGO, ROOT AND HEART ROT IN PINEAPPLE, LATE BLIGHT IN POTATO, ROOT AND COLLAR ROT IN STONEFRUIT, LEATHER ROT AND PHYTOPHTHORA DISEASES IN STRAWBERRY, LATE BLIGHT IN TOMATO, DOWNY MILDEW, PHYTOPHTHORA & PYTHIUM IN ORNAMENTALS, INTERIORSAPES, BEDDING PLANTS, VEGETABLE TRANSPLANTS, AND HERBS AND SPICES, PHYTOPHTHORA AND FUSARIUM IN CONIFERS, PYTHIUM IN TURF, AND PHYTOPHTHORA AND PYTHIUM DISEASES ASSOCIATED WITH STEM AND CANKER BLIGHT (SUDDEN OAK DEATH) AND GENERAL BEECH DECLINE AND VARIOUS TREE STEM CANKERS AND PHLOEM CANKERS.

ACTIVE INGREDIENTS: Mono- and di-potassium salts of Phosphorous Acid*

45.8%

OTHER INGREDIENTS:

54.2%

TOTAL:

100%

*Contains 5.17 lbs/gallon of the active ingredients, mono- and di-potassium salts of Phosphorous Acid.
Equivalent to 3.35 lbs Phosphorous Acid/gallon

KEEP OUT OF REACH OF CHILDREN CAUTION



Department of Agriculture
STATE OF HAWAII

EPA Reg. No. 71962-1

EPA Est. No. 71962-AUS-001

72499-GA-001

71975-WA-001

68253-CO-001

LICENSED

Manufactured for:

LIQUID FERTILISER PTY. LTD. (Trading as Agrichem)

320-222-3026

PERIOD 2015-2017 LIC. NO.

8953.3

NET CONTENTS:

- | | |
|------------------------------------|--------------------------------------|
| <input type="checkbox"/> 1 Pint | <input type="checkbox"/> 25 Gallons |
| <input type="checkbox"/> 2 Quarts | <input type="checkbox"/> 55 Gallons |
| <input type="checkbox"/> 1 Gallon | <input type="checkbox"/> 250 Gallons |
| <input type="checkbox"/> 5 Gallons | |



agrichem[®]

2 Hovey Road

Yatala QLD, 4207

Australia

Phone: +61 7 3451 0000

Fax: +61 7 3451 0093

Web: www.agrichem.com.au

AGRI-FOS[®]

400

Commercial & Agricultural Uses

SYSTEMIC FUNGICIDE FOR THE EFFECTIVE CONTROL OF VARIOUS PLANT DISEASES INCLUDING BLACK SPOT OR SCAB IN APPLE, ROOT ROT IN AVOCADO, BUD ROT AND NUT FALL IN COCONUT, ROOT ROT IN CITRUS AND CUCURBITS, DOWNY MILDEW IN CUCURBITS, GRAPE, LETTUCE, AND ONION, ANTHRACNOSE IN MANGO, ROOT AND HEART ROT IN PINEAPPLE, LATE BLIGHT IN POTATO, ROOT AND COLLAR ROT IN STONEFRUIT, LEATHER ROT AND PHYTOPHTHORA DISEASES IN STRAWBERRY, LATE BLIGHT IN TOMATO, DOWNY MILDEW, PHYTOPHTHORA & PYTHIUM IN HERBS AND SPICES.

ACTIVE INGREDIENTS: Mono- and di-potassium salts of Phosphorous Acid*

45.8%

OTHER INGREDIENTS:

54.2%

TOTAL:

100%

*Contains 5.17 lbs/gallon of the active ingredients, mono- and di-potassium salts of Phosphorous Acid.

Equivalent to 3.35 lbs Phosphorous Acid/gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

Consult page 2 of this booklet for First Aid and Precautionary Statements

EPA Reg. No. 71962-1

EPA Est. No. 71962-AUS-001

72499-GA-001

71975-WA-001

68253-CO-001

Batch No:

Date of

Manufacturer:

Manufactured for:

LIQUID FERTILISER PTY. LTD. (Trading as Agrichem)

320-222-3026

NET CONTENTS:

- | | |
|------------------------------------|--------------------------------------|
| <input type="checkbox"/> 1 Pint | <input type="checkbox"/> 25 Gallons |
| <input type="checkbox"/> 2 Quarts | <input type="checkbox"/> 55 Gallons |
| <input type="checkbox"/> 1 Gallon | <input type="checkbox"/> 250 Gallons |
| <input type="checkbox"/> 5 Gallons | |



agrichem[®]

2 Hovey Road
Yatala QLD, 4207

Australia

Phone: +61 7 3451 0000

Fax: +61 7 3451 0093

Web: www.agrichem.com.au

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice
- Have person sip a glass of water if able to swallow
- Do not induce vomiting unless told to do so by a poison control center or doctor
- Do not give anything to an unconscious person

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye
- Call poison control center or doctor for treatment advice

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing
- Rinse skin immediately with plenty of water for 15-20 minutes
- Call a poison control center or doctor for treatment advice

IF INHALED:

- Move person to fresh air
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible
- Call a poison control center or doctor for further treatment advice

Have the product container or label with you when calling a poison control center or doctor or going for treatment.
Hotline Number: National Poison Control, 1-800-222-1222

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist or vapors. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators, mixers, loaders, and other handlers must wear:

- Protective eyewear
- Long pants and long-sleeved shirt
- Waterproof gloves
- Shoes plus socks.

Follow manufacturer's instructions for maintaining/cleaning personal protective equipment (PPE). If no such instructions for washables, use hot water and detergent. Keep and wash PPE separately from other laundry.

When handlers use closed systems, aircraft or enclosed cabs in a manner that meets the requirements listed in the worker protection standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before drinking, eating, chewing gum, using tobacco or using the toilet.
- Remove PPE clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For Terrestrial Use: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the state or tribal agency responsible for pesticide registration.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, CFR 40 part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours, unless wearing the appropriate PPE.

PPE required for early entry to treated areas that are permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soils or water, is: coveralls worn over short-sleeved shirt and short pants, waterproof gloves, shoes plus socks and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

CHEMIGATION

Use of **AGRI-FOS® 400** through chemigation is not permitted in California.

Apply this product only through center pivot, solid set or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPRINKLER AND DRIP (TRICKLE) IRRIGATION SYSTEMS:

The irrigation system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

(Sprinkler Chemigation Only) Do not apply when wind speed favors drift beyond the area intended for treatment.

Apply **AGRI-FOS® 400** continuously for the duration of the water application. After treatment with **AGRI-FOS® 400** has been completed, avoid further irrigation of the treated area until foliage is dry or for 24 to 48 hours.

GENERAL APPLICATION INSTRUCTIONS

Apply **AGRI-FOS® 400** by various application methods, including foliar spray (aerial and ground), soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply **AGRI-FOS® 400** with sufficient water volumes for adequate coverage of foliage, according to crop and growth stage. To ensure good coverage, spray to wetness, but avoid run-off. This product does not have a pre-harvest interval (PHI). Harvest when dry.

When using **AGRI-FOS® 400** with Supa Stik PN, adhere to all applicable label directions. Only use Supa Stik PN with basal bark applications.

MIXING INSTRUCTIONS

1. Fill the spray tank with $\frac{1}{4}$ - $\frac{1}{2}$ of the volume of water required before adding **AGRI-FOS® 400**
2. Add **AGRI-FOS® 400** slowly to the tank and agitate by mechanical or hydraulic means
3. Continue agitating as tank fills with water to the desired volume
4. Maintain agitation during application

COMPATIBILITY

AGRI-FOS® 400 is compatible with most products used in agriculture. However, individual crop sensitivity to these mixtures may vary. Mixtures of **AGRI-FOS® 400** with some foliar fertilizers and copper products are not always compatible or cause phytotoxicity to some plants. If these combinations or others have not been used previously, do not tank mix without first testing the compatibility of the tank mix. Do not apply tank mixture without first assessing its safety to the crop (phytotoxicity). Tank mix **AGRI-FOS® 400** with fertilizers only if crop safety has been established and the **AGRI-FOS® 400** use rates are followed.

Due to **AGRI-FOS® 400**'s acidic nature, do not use acidifying-type compatibility agents. If spray adjuvants are used test them before use to confirm compatibility with **AGRI-FOS® 400**.

Use a jar test to confirm compatibility with **AGRI-FOS® 400**. In a clean jar using the same water source that is normally used to fill spray tank, add the same proportions of each product and the appropriate quantity of water and mix thoroughly. Let stand for 3 minutes. If mixture remains in solution or is remixed readily the tank mix is compatible.

Spray the solution that results from the above compatibility test onto a few plants and inspect for visual effects of phytotoxicity (leaf burn) 3 days later.

GENERAL HYDROPONIC APPLICATIONS*

For use with plants grown in recirculating hydroponic systems.

Disease	Rate	Application Program
General root rots (Pythium and Phytophthora spp) and root diseases	1 to 2 quarts in 5300 gallons of nutrient solution	Apply every 4 to 6 weeks in the summer and every 8 weeks in the fall. The application time interval may be modified depending on crop load, water quality, and disease pressure.

*Not for use in California

AGRICULTURAL APPLICATIONS

APPLES, CRAB APPLES, LOQUATS, PEARS & QUINCE*

Use AGRI-FOS® 400 for effective control of black spot, root and collar rot and fire blight in apples, crab apples, loquats, pears, and quinces.

Disease	Application Method	Rate	Application Program
Apple black spot and scab (Venturia inaequalis)	Foliar spray Ground	Apply 3 to 4 pints in 100 gallons water per acre.	Apply in combination with a mancozeb-containing product at 3 lbs/acre. Apply at ¼ - ½ inch green tip through first cover at 7 to 10 day intervals or according to forecasted infection events. Continue with AGRI-FOS® 400 and Captan in the remaining applications.
Apple black spot and scab (Venturia inaequalis)	Basal bark spray at bud swell or silver tip stage of growth in early spring	62.4 fl. oz. + 62.4 fl. oz. of water + 1.5 to 3.0 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS® 400 and Supa Stik PN on the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present (treatment generally lasts 8-12 weeks depending on pathogen levels. Higher disease pressure will shorten the length of control).
Root and collar rot (Phytophthora cactorum) Fire Blight (Erwinia amylovora)	Foliar spray Ground	2½ to 5 pints in 100 gallons of water per acre	Thorough spray coverage of plant is required. Start applications when conditions favor disease development. Apply at one to two month intervals. Use the low rate on the shorter interval and the high rate on the longer interval.
	Basal bark spray. Treat in spring and fall for best results.	62.4 fl. oz. + 62.4 fl. oz. of water + 1.5 to 3.0 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS® 400 and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. (Treatment generally lasts 8-12 weeks depending on pathogen levels. Higher disease pressure will shorten the length of control).

*Not for use in California except basal bark spray on apples and crab apples.

ASPARAGUS*Use **AGRI-FOS® 400** for effective control of crown rot & asparagus spear slime disease in asparagus.

Disease	Application Method	Rate	Application Program
Crown rot & Asparagus spear slime (Phytophthora spp)	Foliar spray Ground	1¼ quarts per acre in 25 gallons water to 2½ quarts per acre in 250 gallons water	Apply to ferns that have 2 to 3 inches of new growth. Do not apply to ferns that are starting to die down (senesce). Established plantings, start applications when conditions are favorable to disease (cool wet conditions). Ensure thorough coverage.

*Not for use in California

AVOCADOSUse **AGRI-FOS® 400** for effective control of root rot, trunk cankers and downy mildew disease in avocado.

Disease	Application Method	Rate	Application Program
Root rot (Phytophthora cinnamomi)	Tree injection	Skeletal trees 1st year: ¼ fl. oz. undiluted product per yard of canopy diameter. Other situations: ¾ teaspoon diluted with ½ fl. oz. of water per yard of canopy diameter.	Inject trees at spring flush maturity. Repeat post vegetative at summer flush maturity. Drill holes 3/16 inch (5 mm) in diameter and 1 inch (25 mm) to 2 inches (50 mm) deep with slight downward angle in trunk. Place syringes in the main trunk of the tree and space evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun, Smart Shot injector, or hydraulic tree injection. Do not prune back trees before injection process as burning of new growth may occur. Do not inject trees in winter months. Do not cut back the canopy of injected trees. Do not add any material, other than water, to AGRI-FOS® 400 by trunk injection. Do not inject more liquid in a lesser number of syringes than directed.
	Foliar spray Ground	2 quarts in 100 gallons of water per acre	Spray to run off at 2 – 2½ gallons of spray solution per adult tree. Start applications in spring, up to 4 applications a year at two-month intervals. Ensure thorough coverage.
Canker (Phytophthora citricola)	Trunk spray	1¼ to 2½ quarts mixed with 5 gallons of water with 3.0 fl. oz. of Supa Stik PN Bark Penetrating Surfactant	Apply to trunk lesions using sufficient spray volume to completely wet the trunk and lesions. If lesions absent, apply to trunk from soil level to two feet up trunk. If lesions present, use higher rate.
Downy mildew	Foliar spray Ground	3¼ pints in 100 to 500 gallons water	Spray to run-off, as required for disease control.

BERRIES*

Use **AGRI-FOS® 400** for effective control of root rot, general leaf and berry diseases such as those caused by *Septoria* spp., and downy mildew, and for suppression of *Anthraconose* spp. in bush and cane berries such as, but not limited to, bingleberries, blackberries, black satin berries, blueberries, boysenberries, Cherokee blackberries, Chesterberries, Cheyenne blackberries, coryberries, cranberries, darrowberries, dewberries, Dirksen thornless berries, elderberries, Himalayaberries, huckleberries, hullberries, Lavacaberries, loganberries, lowberries, lucretiaberries, mammoth blackberries, marionberries, mulberries, nectarberries, olallieberries, Oregon evergreen berries, raspberries (red, black, hybrids/cultivars), and youngberries.

Disease	Application Method	Rate	Application Program
Root rot (<i>Phytophthora</i> spp.)	Foliar spray Ground	1-3 quarts per 100 gallons of water per acre. Ensure foliage is completely wet.	New plantings: start application when new growth is 2 to 3 inches long. Established plantings, start applications when cool wet conditions occur which favor disease. West of Rocky Mountains: Autumn applications, apply when conditions favor disease, repeat in 4 weeks. Spring applications, first application after bud break and repeat in 4 weeks. East of Rocky Mountains: First application, spring post bud break (2 to 3 inches new growth) and repeat at 50 to 60 day intervals. Do not exceed 4 applications per season. For blueberries - First application in spring at pink bud and then on a regular schedule of application at two to three intervals.
General leaf and berry diseases such as those caused by <i>Septoria</i> spp and <i>Anthraconose</i> spp	Root dip	Mix a 2.5% v/v solution (1 quart per 10 gallons of water)	Apply as pre-plant dip to the roots for 2-3 minutes. Plant within 48 hours after dipping. Mix a fresh solution daily.
	Chemigation Overhead	1 to 2 quarts in 1000 gallons of water per acre	Apply with normal irrigation schedule. Do not apply more than 4 times per crop cycle.
	Low volume	2 to 3 quarts in 100 gallons of water per acre	
Downy mildew	Foliar spray Aerial	1½ to 2 quarts in 20 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 3 week intervals. Do not apply more than 6 times per crop cycle.
	Ground	1½ to 2 quarts in up to 100 gallons of water per acre	High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle. AGRI-FOS® 400 is best when used in combination with conventional registered fungicides to increase the performance of the disease control program.

*Not for use in California

BRASSICAS*

Use **AGRI-FOS® 400** for effective control of damping-off, root rot, bacterial, and fungal diseases, and downy mildew in brassicas such as, but not limited to, broccoli, Brussels sprouts, cabbage, cauliflower, cavalo broccolo, collards, Chinese cabbage, Chinese mustard cabbage, kale, kohlrabi, mizuna, mustard greens, mustard spinach and rape greens.

Disease	Application Method	Rate	Application Program
Root rot and damping off (Phytophthora spp and Pythium spp)	Foliar spray Aerial Ground Pre-plant seedling nursery treatment Transplant and furrow application	1 to 1½ quarts in 10 gallons of water per acre 1 to 2 quarts in 30 gallons of water per acre 1 quart in 100 gallons of water to nursery plants in seedling trays 1 to 7 days prior to out planting 3 pints at planting or to newly planted seedlings by side dressing or shank application.	Begin application after plants are established and conditions favor disease development. Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Bacterial diseases	Chemigation Overhead: Low volume:	1 to 2 quarts in 1000 gallons of water per acre 2 to 3 quarts in 100 gallons of water per acre	Apply with normal irrigation schedule. Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Downy mildew (Peronospora parasitica)	Foliar spray Ground	1¼ quarts per acre in 25 gallons of water per acre to 2½ quarts per acre in 250 gallons of water per acre California: 1¼ quarts per acre in 100 gallons of water to 2½ quarts per acre in 185 gallons of water	1 to 3 week intervals between applications when conditions favor disease development (cool, moist weather). Use higher rates and shorter intervals when disease pressure increases.
Diseases caused by Septoria, Collectotrichum, Alternaria, and Powdery mildew	Foliar spray Ground	1 to 2 quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

*Use in California limited to control of downy mildew

CEREAL GRAINS*

Apply AGRI-FOS® 400 to crops such as, but not limited to, field corn, ornamental corn, sweet corn, Indian corn, wheat, rye, barley, oats, triticale, and sorghum (milo). Use AGRI-FOS® 400 for effective control of damping-off and root rot diseases.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray Ground	1½ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water	Corn: Apply at 14-day intervals from 4-leaf stage, as needed. Assure good coverage. Other grains: Apply at 14-21 day intervals, as needed. Assure good coverage.

*Not for use in California

CITRUS - Mature trees

Use AGRI-FOS® 400 for effective control of root rot and collar rot diseases in citrus.

Disease	Application Method	Rate	Application Program
Brown rot and root rot Phytophthora spp	Foliar spray Ground	2½ quarts per acre in a maximum of 250 gallons of water	When conditions favor disease. Spray trees to runoff ensure even coverage. Do not apply at high temperatures (above 95°F) particularly if humidity is low or to moisture-stressed trees.
Root rot and collar rot (Phytophthora spp Phytophthora nicotianae and Phytophthora citrophthora)	Trunk spray	Mix 1¼ to 2½ quarts in a minimum of 5 gallons of water with 1 to 3 fl. oz. of Supa Stik PN Bark Penetrating Surfactant	Spray trunk lesions with enough spray volume to ensure lesions are completely wet. When disease levels are high, use higher rate.
* Pre-harvest blue and green mold and brown rot (Phytophthora citricola)	Foliar spray Ground	2 quarts in up to 100 gallons of water per acre	Apply 2 to 4 weeks prior to harvest. Ensure fruit is thoroughly covered by the spray application.

* Not for use in California.

COCONUTS

Use AGRI-FOS® 400 for effective control of bud rot and nut fall in coconuts.

Disease	Application Method	Rate	Application Program
Bud rot – Nut fall (Phytophthora palmivora)	Injection	2 tsp. to 1 fl. oz. per tree	Dilute AGRI-FOS® 400 with water to give final injection volume of 1 fl. oz. to 2 fl. oz. of water and AGRI-FOS® 400. Inject into the trunk or root system.

COTTON*

Use AGRI-FOS® 400 for effective control of damping-off and root rot diseases in cotton.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray Ground	1 to 2 quarts in 30 to 50 gallons of water per acre	Apply at crop emergence every 21 days or during wet conditions that favor pathogen development.

*Not for use in California

CUCURBITS

Use AGRI-FOS® 400 for effective control of sudden wilt, gummy stem blight, downy mildew, powdery mildew, Alternaria leaf blight, and bottom soft rot complex, and for suppression of Anthracnose in cucurbits grown in field situations such as, but not limited to, cucumber, Chinese waxgourd, citron melon, gherkin rock melon, honeydew melon, pumpkin, zucchini, watermelon and squash (summer and winter), momordica spp balsam apple, balsam pear, bitter melon, and Chinese cucumber.

Disease	Application Method	Rate	Application Program
Sudden wilt – Root and fruit rot (Phytophthora spp) Phytophthora blight (Phytophthora capici)	Foliar spray Ground	1-3 quarts per 100 gallons.	Entire spray coverage of plant is required. Do not exceed a total of 6 applications per season.
Gummy stem blight (Mycosphaerella melonis)	Foliar spray Ground	Apply 1¼ quarts per acre in 40 gallons of water. (CA only)	Apply when disease is evident. Continue applications at 21 day intervals until cure is apparent. Do not exceed a total of 6 applications per season.
Downy mildew (Pseudoperonospora cubensis)	Foliar spray Ground		Apply within 7 to 10 days of infection. Repeat as necessary. Do not exceed a total of 6 applications per season.
* Powdery mildew and other leaf diseases such as Anthracnose and Alternaria leaf blight	Foliar spray Ground	2 to 2½ quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
* Bottom soft rot complex	Foliar spray Ground	2 to 2½ quarts in up to 100 gallons of water per acre	Apply after fruit set and during bulking 3 times during the growing season.

*Not for use in California

CUCURBITS, TANK MIXTURES*

For the effective control of downy mildew diseases, tank-mix AGRI-FOS® 400 with a mancozeb-containing fungicide and apply to cucurbits.

Product	Disease	Rate Per Acre	Application Program
AGRI-FOS® 400 + mancozeb-containing fungicide	Downy mildew	Light to medium foliage cover: Apply 1½-2 quarts of product per acre. Heavy foliage cover: Apply 3 quarts of product per acre + Label rate of mancozeb	Apply as a ground foliar spray the in- dicated quantity and dilution rates with water of both products. For best results apply AGRI-FOS® 400 as a tank mix with protectant fungi- cides such as mancozeb, copper oxy- chloride, etc., to ensure both pre- and post-infection activity.

*Not for use in California

FRUITING VEGETABLES*

Use AGRI-FOS® 400 for effective control of damping-off, root rot, gummy stem blight diseases, bacterial diseases, downy mildew, powdery mildew, and Alternaria leaf blight, and for suppression of Anthracnose in eggplant, tomatoes, tomatillos, and peppers such as, but not limited to, bell, chili, cooking, pimento, and sweet.

Disease	Application Method	Rate	Application Program
Eggplant: Pythium and Phytophthora spp, and Gummy stem blight (Mycosphaerella melonis)	Foliar spray Ground	Apply 1¼ quarts per acre in 40 gallons of water.	Entire spray coverage of plant is re- quired. Do not exceed a total of 6 applications per season. Apply when disease is evident. Contin- ue applications at 21-day intervals until cure is apparent. Do not exceed a total of 6 applications per season.
Peppers: Late blight and root rot (Phytophthora infestans and Phytophthora spp)	Foliar spray Ground	1¼ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water.	First application at transplant or when direct seeded crops are at 2-4 true leaf, then at one to two week intervals as required to control disease. In high disease situations, use higher rates and shorter spray intervals.
Tomatoes/Tomatillos: Late blight and root rot (Phytophthora infestans and Phytophthora spp)	Foliar spray Ground	1¼ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water California: 1¼ quarts per acre in 90 gallons of water	First application at transplant or when direct seeded crops are at 2-4 true leaf, then at one to two week intervals as required to control disease. In high disease situations use, higher rates and shorter spray intervals.

FRUITING VEGETABLES* Continued

Disease	Application Method	Rate	Application Program
Root rot and damping off (Phytophthora spp and Pythium spp) Bacterial diseases	Foliar spray Aerial	1 to 1½ quarts in 10 gallons of water per acre	Begin application after plants are established and conditions favor disease development.
	Ground	1 to 2 quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.
	Pre-plant seedling nursery treatment	1 quart in 100 gallons of water to nursery plants in seedling trays 1 to 7 days prior to out planting	High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle. For control of bacterial leaf spot of tomatoes, apply the high rate of AGRI-FOS® 400 with registered bactericides.
	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	Apply with normal irrigation schedule. Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Downy mildew	Foliar spray Aerial	1 to 1½ quarts in 10 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.
	Ground	1 to 2 quarts in 30 gallons of water per acre	High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Powdery mildew and other leaf diseases such as Anthracnose and Alternaria leaf blight	Foliar spray Aerial	1 to 1½ quarts in 10 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle.
	Ground	1 to 2 quarts in 30 gallons of water per acre	High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

* Use in California limited to control of late blight and root rot in tomatoes and tomatillos.

GRAPES*

Use AGRI-FOS® 400 for effective control of root rot, downy mildew, and powdery mildew diseases in grapes.

Disease	Application Method	Rate	Application Program
Downy mildew (Plasmopara viticola)	Foliar spray Ground	Early season small canopy, 1¼ quarts per acre in 100 gallons water Late season/large canopy, 2 quarts per acre in 150 gallons water to 2½ quarts per acre in 200 gallons water	It is essential that the rate of AGRI-FOS® 400 be adjusted to the vine-row volume, i.e., the volume of vine foliage per acre. Spray timing is critical. Apply AGRI-FOS® 400 at times of high disease risk, especially between the time that conditions are conducive to downy mildew infection and the appearance of oil spots. Ensure spray coverage is adequate and that the appropriate rate of AGRI-FOS® 400 is applied to match vine growth, particularly from mid-season onwards, and especially where grapes are grown on root stock
Phytophthora and Pythium root rots	Foliar spray Ground	1½ to 2 quarts in 50 gallons of water per acre	Apply to vines that have stressed root system that can lead to root rots. Mitigating factors such as nematode pressure, water logging, and compaction contribute to vine declines. Table Grapes: Begin application in the Spring at the 4 to 6 inch shoot stage. Continue applications at 1 to 2 week intervals until flowering. Resume applications in the Fall after harvest. Do not apply more than 4 times per crop cycle. Wine and Raisin Grapes: Begin applications in the Spring at the 4 to 6 inch shoot stage. Continue applications at 1 to 2 week intervals through flowering. Do not apply more than 4 times per crop cycle.
Downy mildew (Plasmopara viticola)	Foliar spray Ground	1½ to 2 quarts in 50 gallons of water per acre AGRI-FOS® 400 is most effectively applied to control downy mildew when tank mixed with other registered fungicides.	Apply a bud break with additional applications at 7 to 10 day intervals in a rotational program with other registered fungicides. Use higher rate and volume based on disease severity and density of canopy. Do not apply more than 6 times per crop cycle.
Powdery mildew (Uncinular necator)	Foliar spray Ground	2 to 2½ quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

* Use in California limited to control of downy mildew.

GRAPES, TANK MIXTURES*

For the effective control of downy mildew diseases, tank-mix AGRI-FOS® 400 with a mancozeb-containing fungicide and apply to grapes.

Product	Disease	Rate Per Acre	Application Program
AGRI-FOS® 400 + mancozeb-containing fungicide	Downy mildew	Early season small canopy, 1½ quarts per acre in 100 gallons water Late season/large canopy, 2 quarts per acre in 150 gallons water to 2½ quarts per acre in 200 gallons water + Label rate of mancozeb	Apply as a foliar spray the indicated quantity and dilution rates with water of both products. For best results, apply AGRI-FOS® 400 as a tank mix with protectant fungicides such as mancozeb, copper oxychloride, etc., to ensure both pre- and post-infection activity.

*Not for use in California.

HERBS AND SPICES*

Use AGRI-FOS® 400 for effective control of Downy mildew, Phytophthora spp and Pythium spp diseases of Herbs and Spices grown in fields, nurseries and greenhouses. Apply AGRI-FOS® 400 to plants such as, but not limited to, Anise, Balm, Basil, Chamomile, Caraway, Catnip, Celery, Chives, Coriander, Cumin, Curry Leaf, Dill, Fennel, Marjoram, Mint, Nasturtium, Rosemary, Sage, Savory, Sweet Bay, Tarragon, Thyme, and Wintergreen.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to plants that are heat or moisture stressed.

Disease	Application Method	Rate	Application Program
Downy Mildew	Foliar spray Ground	1¼ to 2½ quarts per 100 gallons of water Or ½ to 1 1/8 fl. oz. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required.
Phytophthora spp and Pythium spp	Foliar spray Ground	1 to 2 quarts per 100 gallons of water Or 2 to 4 tsp. per gallon of water	Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days. Repeat as required. Note: do not apply more than 500 gallons of spray solution per acre.
	Soil drench	6¼ to 12¼ fl. oz. per 100 gallons of water	Apply each 25 gallons of solution to an area of 100 square feet. Follow application with irrigation. Repeat as required. Limit of one application per month.

*Not for use in California.

HOPS*

Use AGRI-FOS® 400 for effective control of downy mildew in hops.

Disease	Application Method	Rate	Application Program
Downy mildew	Foliar spray by ground equipment only.	1-3 quarts per 100 gallons of water per acre	When conditions favor disease, apply when: A. Shoots are ½ to 1 foot long. B. Post-training when vines are 6 feet high. C. 21 days post-application (B). D. During bloom.

*Not for use in California.

LEAFY VEGETABLES*

Use AGRI-FOS® 400 for effective control of root rot, damping-off, downy mildew, powdery mildew, and other fungal leaf diseases, and for suppression of Anthracnose in leafy vegetables such as, but not limited to, amaranth, arugula, cardoon, celery, chervil, chrysanthemum, corn salad, cress, dandelion, dock, endive, fennel, lettuce, orach, parsley, purslane, radicchio, rhubarb, spinach, and Swiss chard. Excludes Brassica vegetables.

Disease	Application Method	Rate	Application Program
Downy mildew (Bremia lactucae)	Foliar spray Ground	½ gallon per acre in 40 gallons water	Ensure spray coverage is adequate to wet the whole plant. During warm, wet conditions, repeat application at 7 to 10 day intervals, if needed.
Root rots and damping off (Phytophthora and Pythium spp)	Foliar spray Ground	Apply 1 to 2 quarts in 40 gallons of water per acre	Begin application after plants are established and conditions favor disease development.
	Pre-plant nursery application	1 quart in 100 gallons of water to nursery plants in seedling trays 1 to 7 days prior to out planting	Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
	Chemigation Overhead	Apply 1 to 2 quarts per acre in a minimum of 1000 gallons per acre of water	Apply with normal irrigation schedule.
	Low Volume	Apply 1 to 2 quarts per acre in a minimum of 100 gallons per acre of water	Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	

LEAFY VEGETABLES* Continued

Disease	Application Method	Rate	Application Program
Powdery mildew and leaf diseases such as Anthracnose (<i>Collectotrichum</i> spp), leaf blights (<i>Septoria</i> and <i>Cercospora</i> spp), and bacterial rots (<i>Erwinia</i> spp)	Foliar spray Ground	1½ to 2 quarts in a 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

*Use in California limited to control of downy mildew in lettuce.

LEGUMES*

Use AGRI-FOS® 400 for effective control of damping-off ,root rot, Fusarium, Rhizoctonia, downy mildew, powdery mildew, and other fungal leaf diseases, and for suppression of Anthracnose in legumes (succulent and dried) such as, but not limited to, green beans, soybeans, wax beans, field beans, navy beans, lima beans, fava beans, kidney beans, pinto bean, mung beans, broad beans, lentils, chickpeas, English peas, snow peas, sugar snap peas, black-eyed peas, cow peas, and pigeon peas.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray Ground	1½ to 2½ quarts in 30 gallons of water per acre	Begin application after plants are established and conditions favor disease development.
Fusarium and Rhizoctonia	Pre-plant nursery application Transplant and furrow application	1 quart in 100 gallons of water to nursery plants in seedling trays 1 to 7 days prior to out planting 3 pints at planting or to newly planted seedlings by side dressing or shank application	Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Downy mildew	Foliar spray Ground	1 to 3 quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1-2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7-10 day intervals. Do not apply more than 6 times per crop cycle.

LEGUMES* Continued

Disease	Application Method	Rate	Application Program
Powdery mildew and leaf diseases such as Anthracnose (<i>Collectotrichum</i> spp), leaf blights (<i>Septoria</i> and <i>Cercospora</i> spp), and bacterial rots (<i>Erwinia</i> spp)	Foliar spray Ground	2 to 2½ quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

*Not for use in California.

MANGOS

Use AGRI-FOS® 400 for suppression of anthracnose in mangos.

Disease	Application Method	Rate	Application Program
Anthracnose (<i>Colletotrichum gloeosporioides</i>)	Foliar spray Ground	2 tsp. per gallon of water	Spray tree every 14 days during blossom period, then monthly until harvest. Spray to the point of run-off.

MISCELLANEOUS CROPS*

Use AGRI-FOS® 400 for effective control of damping-off, root rot, downy mildew, powdery mildew, and other bacterial and leaf diseases, and for suppression of Anthracnose in coffee, okra, papaya, and persimmon.

Disease	Application Method	Rate	Application Program
Bacterial and leaf diseases such as coffee berry disease (<i>Collectotrichum</i> spp), Anthracnose (<i>Collectotrichum</i> spp), and various leaf spots (<i>Septoria</i> and <i>Cercospora</i> spp)	Foliar spray Aerial	1 to 2 quarts in 10 gallons of water per acre	Begin application after plants are established and conditions favor disease development.
	Foliar spray Ground	1 to 2 quarts in 100 gallons of water per acre	Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
	Root dip	Mix a 0.25% v/v solution (1/3 fl. oz. with 1 gallon of water)	Apply as pre-plant dip to transplants immediately prior to planting. Dip plants momentarily and plant within 48. Mix a fresh solution daily.

MISCELLANEOUS CROPS* Continued

Disease	Application Method	Rate	Application
	Chemigation Overhead Low volume Transplant and furrow application	2 to 3 quarts in 1000 gallons of water per acre 2 to 4 quarts in 100 gallons of water per acre 3 pints at planting or to newly planted seedlings by side dressing or shank application	Apply with normal irrigation schedule. Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Downy mildew	Foliar spray Aerial Foliar spray Ground	1 to 3 quarts in 10 gallons of water per acre 1 to 3 quarts in 100 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week in- tervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Powdery mildew	Foliar spray Aerial Foliar spray Ground	1 to 3 quarts in 10 gallons of water per acre 1 to 3 quarts in 100 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.
Pseudomonas garcae	Foliar spray Ground	Mix a 0.50% v/v solution (2 quarts per 100 gallons of water)	Apply prior to onset of disease. Apply spray to saturation or runoff.

*Not for use in California.

NONGRASS ANIMAL FEED*

Use AGRI-FOS® 400 for effective control of damping-off and root rot diseases in forage crops such as, but not limited to, alfalfa, clover, and vetch.

Disease	Application Method	Rate	Application Prgoram
Phytophthora and Pythium spp	Foliar spray	1 1/2 quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water.	Apply at 14-day intervals after plant emergence, as needed. Assure good coverage.

*Not for use in California.

ONIONS

Use AGRI-FOS® 400 for effective control of downy mildew disease in onions, garlic, shallots, and leeks. Use as a preventative control program for best results.

Disease	Application Method	Rate	Application Program
Downy mildew (Peronospora destructor)	Foliar spray Ground	2 quarts per acre in 100 gallons water per acre	As a regular preventative control program or when disease first appears.

ONIONS, TANK MIXTURES*

For the effective control of downy mildew diseases, tank-mix AGRI-FOS® 400 with a mancozeb-containing fungicide and apply to onions, garlic and shallots.

Product	Disease	Rate Per Acre	Application Program
AGRI-FOS® 400 + mancozeb-containing fungicide	Downy mildew	2 quarts per acre in 100 gallons water per acre + Label rate of mancozeb	Apply as a foliar spray the indicated quantity and dilution rates with water of both products. For best results, apply AGRI-FOS® 400 as a tank mix with protectant fungicides such as mancozeb to ensure both pre- and post-infection activity.

*Not for use in California.

PEANUTS*

Use AGRI-FOS® 400 for effective control of damping-off and root rot disease and for suppression of Anthracnose in peanuts.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp	Foliar spray	1½ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water	Apply at 14-day intervals, as necessary. Ensure thorough coverage.
Leaf and crown diseases, Anthracnose (Collectothricum)	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application	Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

*Not for use in California.

PINEAPPLES

Use AGRI-FOS® 400 for effective control of Phytophthora root and heart rot diseases in pineapples.

Disease	Application Method	Rate	Application Program
Phytophthora root and heart rot (Phytophthora cinnamomi and parasitica spp)	Foliar spray	2½ quarts in 25-50 gallons water per acre.	Apply to tops, 14 days prior to harvest of planting material. Will treat enough slips to plant one acre. Established plantings when conditions favor disease. Apply at 90 day intervals. Ensure thorough coverage of plants.
	Pre-plant dip	1¼ quarts in 100 gallons water	
	Foliar spray	2 quarts in 100 gallons water	

POTATOES, POST-HARVEST*

Use AGRI-FOS® 400 only on russet-skinned potatoes intended for processing for suppression of late blight (Phytophthora infestans), pink rot (Phytophthora erythrosepica), and powdery mildew. Apply 16.5 fl. oz. of AGRI-FOS® 400 per ton of tubers in one half gallon of water as a mist spray. For the best control, be sure the tubers are completely and evenly covered.

Disease	Application Method	Rate	Application Program
Potatoes, Post-harvest: Late blight (Phytophthora infestans) and pink rot (Phytophthora erythrosepica)	Spray on tubers	16.5 fl. oz. in ½ gallon of water/ ton of tubers	Apply only to russet-skinned potatoes intended for processing. For best results, be sure tubers are thoroughly and evenly covered.
Late blight (Phytophthora infestans) and pink rot (Phytophthora erythrosepica)	Foliar spray Ground	2 to 2½ quarts in 50 gallons of water per acre	Begin application after plants are established and conditions favor disease development. Low Disease Pressure: Apply lower rate at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle. Late blight management using AGRI-FOS® 400 requires the high application rate and is most effectively applied when tank mixed with other registered fungicides.
	Seed piece spray Tuber spray		Mix a 15% v/v solution (0.3 quart with 2 quarts of water). Treat 2 tons of tubers per 1 gallon of solution.

POTATOES, POST-HARVEST* Continued

Disease	Application Method	Rate	Application Program
Powdery mildew	Foliar spray Ground	1 to 3 quarts in 50 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 4 times per crop cycle. High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 4 times per crop cycle.

*Not for use in California

ROOT AND TUBER VEGETABLES

Use AGRI-FOS® 400 for effective control of foliar and root rot in ginseng, damping-off and root rot diseases in carrots, late blight disease and storage diseases such as pink rot and pythium leak, and downy mildew in potatoes, sweet potatoes, and yams.

Disease	Application Method	Rate	Application Program
* Ginseng: Foliar and root rot (Phytophthora cactorum)	Foliar spray Ground	2½ quarts in 100 gallons of water	In cool wet conditions that favor Phytophthora. Apply at 7 day intervals. Do not exceed a total of 8 applications per season.
* Carrots: Phytophthora and Pythium spp	Foliar spray Ground	1½ quarts per acre in 25 gallons of water to 2 quarts per acre in 250 gallons of water	Apply at 14-day intervals after plant emergence, as needed. Assure good coverage.
Potatoes, Sweet Potatoes, Yams: Pink rot and Pythium leak (Phytophthora erythroseptica and Pythium spp)	In-furrow spray	2-5 quarts per acre in 10 gallons water	Apply in a band spray directly over top of potato seed just before row is closed.
Potatoes, Sweet Potatoes, Yams: Late blight, Pink rot and Pythium leak (Phytophthora infestans, Phytophthora erythroseptica and Pythium spp)	Foliar spray Ground	1¼ quarts per acre in 90-375 gallons water per acre	Apply at 5 to 14 day intervals subject to disease incidence.
* Downy mildew	Foliar spray Aerial Foliar spray Ground	1 to 3 quarts in 10 gallons of water per acre 1 to 3 quarts in 30 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

*Not for use in California

STONE FRUIT*

Use AGRI-FOS® 400 for effective control of root, collar rot, almond pruning wound, and various canker disease in stone fruit such as, but not limited to, apricots, sweet and tart cherries, nectarines, peaches, plums, and fresh prunes.

Disease	Application Method	Rate	Application Program
Root and collar rot (Phytophthora spp)	Foliar spray Basal bark spray, apply in spring and fall	2½ quarts per acre in 250 gallons water 62.4 fl. oz. + 62.4 fl. oz. of water + 3.2 fl. oz. Supa Stik PN Bark Penetrating Surfactant	Three treatments are required 1. Spring 2. Mid summer 3. Fall, post harvest Spray a combination of AGRI-FOS® 400 and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. Treat in spring and fall for best results.
Almond pruning – wound cankers (Phytophthora syringae)	Paint or spray	1¼-2½ quarts in 100 gallons water	Apply to pruning wound and surrounding area, ensure area is thoroughly wet. In high disease situations use higher rate.
Suppression of armillaria root rot (Armillaria luteobubulina)	Basal bark spray	1½ to 2 quarts + 2 quarts water + 1% Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS® 400 and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 18 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Armillaria root rot, make a fall application prior to leaf senescence and a spring application for best results. For trees less than 18 inches DBH, make an early spring application.

*Not for use in California

STRAWBERRIES*

Use AGRI-FOS® 400 for effective control of leather rot, red stele, Phytophthora, foliar and bacterial disease in strawberries and suppression of powdery mildew.

Disease	Application Method	Rate	Application Program
Phytophthora and Pythium spp (red stele, leather rot, and root rot)	Pre-planting dip	1¼ quarts in 100 gallons water	Dip planting material in this solution for 30 minutes, then plant within one day. Use program for annual and perennial varieties.

STRAWBERRIES* Continued

Disease	Application Method	Rate	Application Program
Foliar fungal and bacterial diseases (Rhizopus and Xanthomonas)	Foliar spray Ground	1 to 3 quarts in 50 to 100 gallons of water per acre	Annual crops, first treatment 14 to 21 days post planting, repeat at 1-2 month intervals when disease is evident.
	Transplant and furrow application	3 pints at planting or to newly planted seedlings by side dressing or shank application.	Perennial crops, first treatment during spring growth flush, repeat at 1-2 month intervals when disease is evident. For susceptible varieties use higher rates and shorter spray intervals. For leather rot, apply at 10% bloom and early fruit set, then 1 to 2 week intervals as required for disease control. In high disease situations, use higher rate and shorter spray intervals. Grey mold and Anthracnose suppression using AGRI-FOS® 400 requires use of high application rates and is most effective when tank mixed with other registered fungicides.
Phytophthora spp	Dip	Mix a 0.25% v/v solution (1 quart in 100 gallons of water)	Dip runners in the solution for 1 to 2 minutes. Plant within 48 hours. Mix a fresh solution daily.
Suppression of powdery mildew	Foliar spray Ground	2 to 2½ quarts in 50 gallons of water per acre	Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

* Use in California limited to control of red stele, leather rot, and root rot.

TREE NUTS*

Use AGRI-FOS® 400 for effective control of root and collar rot, and almond pruning-wound canker disease in tree nuts such as, but not limited to, almonds, beech nuts, Brazil nuts, butternuts, cashews, chestnuts, chinquapin, hazelnuts, hickory nuts, macadamia nuts, pecans, pistachios, and walnuts.

Disease	Application Method	Rate	Application Program
Other than macadamia nuts: Root and collar rot (Phytophthora spp)	Foliar spray Ground	1½ quarts per acre in 125 gallons water	Three treatments are required 1. Spring 2. Mid summer 3. Fall, post harvest

TREE NUTS* Continued

Disease	Application Method	Rate	Application Program
Other than macadamia nuts: Almond pruning – wound canker (Phytophthora syringae)	Paint or spray	2½ quarts in 100 gallons water	Apply to pruning wound and surrounding area, ensure area is thoroughly wet.
Macadamia nuts: Raceme blight (Phytophthora spp)	Foliar spray Ground	3¼ quarts per acre in 250 gallons of water	Apply when disease is first seen and reapply at 3 week intervals. Spray to the point of run-off.
Root rot, crown rot, trunk cankers, foliar blights (Phytophthora and Pythium spp)	Foliar spray Ground	2 to 2½ quarts in 100 gallons of water per acre	Begin application after plants are established and conditions favor disease development. Low Disease Pressure: Apply lower rate at 3 month intervals. Do not apply more than 4 times per crop cycle. High Disease Pressure: Apply higher rate at monthly intervals. Do not apply more than 4 times per crop cycle.
	Root dip	Mix a 0.5% v/v solution (2 quarts in 100 gallons of water)	Dip roots in the solution for 30 seconds. Plant within 48 hours. Mix a fresh solution daily.
Pecan scab	Foliar spray Ground	2 quarts in 50 to 100 gallons of water per acre	Apply preventatively with other products shown to be effective against pecan scab.
Pruning wound, crown and trunk cankers (Phytophthora spp)	Trunk spray	2 quarts + 2 quarts of water + 1% Supa Stik PN Bark Penetrating Surfactant	Apply higher rate when lesions are present. Clean wound sites and apply on and around the lesions using enough spray volume to thoroughly wet the lesions. Apply to the trunk from the soil line to 5 feet up the trunk. Apply one time in the Spring, Summer, and Fall.
Downy mildew (Phytophthora and Peronospora spp) Powdery mildew	Foliar spray Aerial Ground	1 to 2 quarts in 20 gallons of water per acre 1 to 2 quarts in 100 gallons of water per acre	Low Disease Pressure: Apply lower rate at first onset of disease. Repeat applications at 1 to 2 week intervals. Do not apply more than 6 times per crop cycle. High Disease Pressure: Apply higher rate at first onset of disease. Repeat applications at 7 to 10 day intervals. Do not apply more than 6 times per crop cycle.

TREE NUTS* Continued

Disease	Application Method	Rate	Application Program
Tree cankers	Basal bark spray	1½ to 2 quarts + 2 quarts water + 1% Supa Stik PN Bark Penetrating Surfactant	Spray a combination of AGRI-FOS® 400 and Supa Stik PN around the complete circumference of the tree trunk until saturation/runoff. Spray from ground level up to 5 feet above the soil line, including the base of the first scaffolding limbs, if present. For large trees larger than 18 inches DBH (Diameter at Breast Height, 4.5 feet above ground) that have been previously attacked by Armillaria root rot, make a fall application prior to leaf senescence and a spring application for best results. For trees less than 18 inches DBH, make an early spring application.

*Not for use in California

SEED TREATMENT APPLICATIONS*

Use AGRI-FOS® 400 for effective control of Phytophthora and Pythium diseases on agricultural crop seeds from crops listed elsewhere on this label. Do not use treated seed for food, feed, or oil. Dye used to color treated seed must be an EPA approved dye (refer to 40 CFR 153-155 (c)). Seed treatment on agricultural establishment in hopper-box, planter box, or other seed treatment application at or immediately before planting is within the scope of WPS, while commercial treatment of seeds is not within the scope of WPS.

Disease	Application Method	Rate
Phytophthora, Pythium and Fusarium spp	Can be applied at-planting or in commercial seed treatment operations.	8-24 fl. oz. AGRI-FOS® 400 per 100 lbs of seed or 4-10 quarts AGRI-FOS® 400 per ton of seed, depending on the size of the seeds to be treated.

*Not for use in California.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE:

Keep this product in containers stored upright and secured with the original closure. Do not store this product near any heat source. Do not store near strong oxidants. If transfer to another container becomes necessary, ensure that the container is clearly labeled, the container is a type suitable for the product, and is clean and free of other materials.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container.

Containers with a capacity of equal to or less than 5 gallons: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Containers with a capacity greater than 5 gallons: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND DISCLAIMER

To the extent consistent with applicable law, all conditions and warranties and statutory or other rights of action which buyer or any other user may have against Liquid Fertiliser Pty. Ltd. are hereby excluded. To the extent consistent with applicable law, Liquid Fertiliser Pty. Ltd. hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by Liquid Fertiliser Pty. Ltd. or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. To the extent consistent with applicable law, Liquid Fertiliser Pty. Ltd.'s liability shall in all circumstances be limited to replacement of product or a refund of the purchase price thereof.