

## FRAGILE. HANDLE WITH CARE.

### IMPORTANT

- The softgel capsule coating is sensitive and can be damaged by certain ambient conditions and rough handling.
- Do not expose capsules to relative humidity above 50% or temperatures above 70F for extended period over 1 hour, or this may affect level of capsule fragmentation - restoring may be possible in some cases, refer 'Capsule Restoration' section of label.
- Do not refrigerate or expose to freezing conditions, this may permanently damage capsules.
- Remove the capsules from container only when ready to apply onto the animals and keep containers sealed at all other times.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.

- In the field do not remove the capsules from container until ready to apply.
- For best results transport the capsules in original sealed packaging until loading into Application Device.
- Apply only with a **SmartVet™ Approved VetCap Application Device** (sold separately) and in accordance with its use directions and treatment method recommendations.
- Never leave the capsules in any Application Device - return all unused capsules to original container and store as directed.
- Only apply when animal is between 15 feet and 30 feet distance from the user.
- Do not use on wet cattle or when rain is expected to wet cattle within twenty-four hours after treatment.
- Do not apply treatment when bystanders are within 30 feet of the animal.
- Do not apply to any animals in or within 10 feet of bodies of water.
- Do not leave intact capsules in the field.
- Do not allow farm dogs or other farm animals to come into contact with the pesticide or residues.

AIM-L is designed for use only with a compatible applicator device that carries the **'SmartVet™ Approved VetCap Application Device'** label. Using any other application device or method could be unsafe, ineffective and/or cause harm or injury.

**How to Apply:** Apply the capsules to the middle of the side of the animal's body anywhere between the front shoulders and 6 inches ahead of the base of the tail.

**Amount to Use:** Apply AIM-L at the dosages shown on the following table. Care must be taken to apply the correct dosage.

Horn Flies, Biting Lice and Sucking Lice	
Animal Weight	Dose Size per Animal
Greater than 600 lbs	1 Capsule (10 ml)

AIM-L can be used on beef cattle of all ages provided they are greater than 600 lbs. Treatment for flies or lice can be repeated as needed but not more often than once every 2 weeks and do not apply more often than four times within any 6-month period. For optimal lice control, an initial application followed by a second treatment 2 weeks later is recommended.

AIM-L does not control cattle grubs.



An Innovative Insecticide Delivery System



Distributed by: Agri Laboratories Ltd, St. Joseph, Missouri, USA, (800) 542 8916  
EPA Reg. No. 88050-2 EPA Est. No. 88050-AR-001

The delivery system is covered by Patents US 8425932B2; US 2013/0216483A1 and United States Patent Application No 14/453930 (amongst others).



## 30 Capsules

Designed Exclusively For Use With



Lambda Cyhalothrin with Piperonyl Butoxide Insecticide Capsules for Control of Horn Flies, Biting Lice and Sucking Lice on Beef Cattle

Keep out of reach of children

### CAUTION

See Back & Side Panels for First Aid and Other Precautionary Statements


ACTIVE INGREDIENTS	% BY WEIGHT
Lambda cyhalothrin	1.50%
Piperonyl Butoxide*	7.50%
<b>OTHER INGREDIENTS</b>	<b>91.00%</b>
Total	100.00%

\*(butyl carbityl) (6-propylpiperonyl) ether and related compounds

### NET CONTENTS:

Thirty 0.338 fl.oz. (10ml) capsules.

BATCH:



## FRAGILE. HANDLE WITH CARE.

To open, cut or apply pressure with thumb along here to break perforation.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION** Harmful if swallowed. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco, using the toilet.

#### FIRST AID

If Swallowed: • Call a poison control center or doctor immediately for treatment advice.  
• Have a person sip a glass of water if able to swallow.  
• Do not induce vomiting unless told to do so by a poison control center or doctor.  
• Do not give anything to an unconscious person.

If in Eyes: • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present after the first 5 minutes, and then continue rinsing eye.  
• Call a poison control center or doctor for treatment advice.

If on Skin or Clothing: • Take off contaminated clothing.  
• Rinse skin immediately with plenty of water for 15-20 minutes.  
• Call a poison control center or doctor for treatment advice.

If Inhaled: • Move person to fresh air.  
• If person is not breathing call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible.  
• Call a poison control center or doctor for treatment advice.

**HOTLINE NUMBERS:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment call 1-800-255-3924. For customer service or to obtain product information, including the SDS, call 1-800 542 8916 or 1-816 233 9533.

**ENVIRONMENTAL HAZARDS:** This pesticide is extremely toxic to aquatic organisms, including fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

**PHYSICAL OR CHEMICAL HAZARDS:** The capsules are sensitive to heat and humidity. Do not expose to relative humidity above 50% or temperatures above 70F for extended period over 1 hour, as this may affect the level of capsule fragmentation.



30 Capsules  
0.338fl.oz. (10ml) each



Designed Exclusively For Use With



An Innovative Insecticide Delivery System

### STORAGE AND DISPOSAL

Not for Use or Storage In or Around the Home.

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Disposal:** Pesticide wastes are hazardous. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** Non-refillable container. Do not reuse or refill any parts of this container. Offer for recycling, if available. Puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Pesticide Storage:** Store in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of reach of children, preferably in a locked storage area. Store under conditions where relative humidity is kept between 30% and 50% and temperatures are maintained between 50F and 70F. Ensure the capsules are never exposed to freezing temperatures during storage, transport or use -this may permanently damage capsules.

**Capsule Restoration:** If inadvertently exposed to high heat and humidity, the capsules can normally be restored to ideal conditions by placing them for 24 to 72 hours in cool & dry conditions where relative humidity is between 30% and 50% and temperatures do not exceed 70F.

**NOTE:** The capsule coating is softgel and sensitive to ambient humidity. Do not expose to relative humidity above 50% or temperatures above 70F for extended period over 1 hour, or this may affect the level of capsule fragmentation. Remove the capsules from container only when ready to apply onto the animals and keep containers sealed at all other times. Handle the container and capsules in a manner as to prevent damage to the capsules. If the capsules or container are leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose as directed for pesticides above.

**Conditions of Sale:** The directions on this label are believed to be adequate and must be followed carefully. This product has been tested under different environmental conditions similar to those that are ordinary and customary for use of the product. However, it is impossible to eliminate all risks associated with the use of this product. Failure to follow label directions may cause injury to animals, humans and damage to the environment. Insufficient control of pests, injury or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of SmartVet USA, its agents, affiliates or assignees. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**Limited Warranty and Limitation of Damages:** SmartVet USA warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SmartVet USA MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL, and no agent of SmartVet USA is authorized to make or modify any warranties beyond those contained herein. Any damages arising from breach of this warranty shall be limited to direct damages and shall not include special, incidental or consequential damages resulting from the use or handling of this product.



SmartVet USA Inc.  
International Animal Health & Food Safety Institute,  
22201 W. Innovation Drive, Olathe, Kansas 66061-1304

SMA 0056-14-C (July2015)Shrinkwrap