

IND/COM



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

9026.41



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

FLYING INSECT
KILLER

Contains the botanical insecticide, Pyrethrum
Kills and Repels Stable Flies, Horse Flies, Face Flies,
Deer Flies, House Flies and Gnats
For use in Animal Quarters, Dairies, Milk Rooms,
Milking Parlors, Calving Areas, Dairy and
Hog Barns as well as outdoor areas
For use on Beef Cattle, Dairy Cattle, Horses,
Hogs, Chickens, and other livestock
Quick-killing pyrethrin controls
flying insects in the dairies, barns, storage sites
and other listed areas

ACTIVE INGREDIENTS:

Pyrethrins	0.25%
*Piperonyl butoxide	2.50%
OTHER INGREDIENTS	97.25%
	100.00%

* (butylcarbityl) (6-propylpiperonyl) ether and 1.00% related compounds.

EPA Reg. No. 1021-1732-7546

EPA Est. 33595-MO-2

EPA Est. 33595-MO-4

EPA Est. No. 706-IL-1

EPA Est. 33595-MO-3

The Est. No. appears after the can code.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side panel for additional precautionary statements.

NET WT. 15 OZ. (425 GRAMS)

FIRST AID
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticides incidents, call 1-888-740-8712.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear the following: Long sleeve shirt, Long pants, Shoes and socks. Chemical-resistant gloves made of any waterproof material.

USER SAFETY RECOMMENDATIONS:

User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

User should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow air tables are more prone to produce runoff that contains this product. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting. Do not use this product in or on electrical equipment due to possibility of shock hazard.

This product contains the quick-killing botanical insecticide, Pyrethrum, which provides effective control of insects upon direct contact, with an added benefit of flushing these insects to aid in more complete kill. This product can be used in listed animal quarters. This product provides fast, effective kill and repellency of the listed pests. This product can also be used on listed livestock and horses for quick kill of pests on these animals. This product can be used in listed pet quarters. This product provides fast, effective kill and repellency of the listed pests. This product can be used in restaurants and other listed areas to kill, on contact, listed pests and is a fast, effective product for fly/ing insect control. Effectively controls exposed stages of stored product pests without persistent residue problems, because it breaks down quickly on treated surfaces exposed to light.

For use in: Animal Holding Areas, Animal Quarters, Attics, Barns, Beef Barns, Bottling Plants, Buildings Housing Animals or Poultry, Calving Barns, Canneries, Commercial Buildings, Crematoriums, Dairy Barns, Dairies, Dried Fruit Storage Areas, Dumpsters, Factories, Farm Structures, Food Processing Plants, Food Storage Areas, Funeral Parlors, Garbage Cans, Goat Holding Areas, Grain Elevators, Grain Storage, Hog Barns, Horse Stables, Hotels, Industrial Plants, Institutions, Kitchens, Milk Sheds, Livestock Holding Areas, Loading Sheds, Museums, Milk Houses, Milk Rooms, Milk Sheds, Milling Rooms, Mortuaries, Pet Sleeping Areas, Poultry Houses, Rail and Cars, Restaurants, Rice Mills, Seed Storage, Sheep Holding Areas, Sheds, Stables, Storage Areas, Stored Products, Supermarkets, Swine Houses, Theaters, Tobacco Warehouses, Trailers, Tree Nut Storage Areas, Trucks, Utility Rooms, Veterinary Clinics, Warehouses, Washrooms, and Wheat Mills.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE

General Directions and Restrictions:

Remove or cover exposed food and drinking water before application. Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces, or wash them before use. Do not use in giraffet cabins.

When used in dairy barns or facilities: Close milk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking. Except when applying to livestock, do not enter or allow others to enter treated area until sprays have dried.

Except when applying to livestock, do not apply this product in a way that will contact workers or other persons, either directly or through drift. Except when applying to livestock, only protected handlers may be in the area during application. Do not apply directly into sewers or drains, or to any area like a gutter where

drainage to storm sewers, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application. For most effective results, follow directions for specific use areas. **SHAKE BEFORE USE AND AT INTERVALS DURING PROLONGED USE.** Hold container upright with nozzle pointing away from face. Press valve down and spray as directed.

FOR USE IN ANIMAL QUARTERS, Barns, Dairies, Milking Parlors, Milking Rooms, Poultry Houses, and other listed animal areas: For rapid kill of Flies, Gnats, Stable Flies, House Flies, Horse Flies, Lesser House Flies, Deer Flies, Horn Flies, and Black Flies: Spray at the rate of 5-10 seconds per 1,000 cubic feet of space, directing the spray toward the ceiling and upper corners of the room and not directly towards the animals or birds. For best results, close doors and windows if possible, before spraying and keep room closed for 10-15 minutes. Applicator should vacate the treated area and ventilate it prior to returning.

FOR USE ON LIVESTOCK:

To protect Beef Cattle, Dairy Cattle, Horses, and other livestock from attacks and to kill and repel Stable Flies, Horse Flies, Deer Flies, House Flies, Horn Flies, Lesser House Flies, and Gnats: Spray about 3 seconds on each side of animal, being careful to spray back, withers, and forelegs thoroughly. To protect from Face Flies, spray the face and head or spray on a cloth and rub on the face, being certain to avoid contact with the eyes. Repeat treatment as necessary when flies are troublesome.

To control Sheep "Tick" or Ked, thoroughly wet all portions of the body by spraying directly into the wool. Treat at a rate to sufficiently wet the animal. To kill Fleas and head ticks on animals and to obtain protection against reinfestation, spray to sufficiently wet animal.

SPACE TREATMENT INDOORS: FOR USE IN FOOD AND NON-FOOD AREAS OF FOOD PROCESSING PLANTS, Industrial Installations, Bakeries, Rice and Wheat Mills, Restaurants, Tobacco Warehouses, Grain Elevators, and Warehouses: Spray as directed to control listed flying insects. Do not make space spray applications when facility is in operation. Prior to space spray applications, cover or remove food. Prior to space spray applications, cover food processing surfaces or clean after treatment and before use. For maximum efficacy, close all doors and windows, turn off fans and air conditioner, and spray up into center of room with slow, sweeping motion. Spray 5-10 seconds for average size room 10 ft. x 12 ft. x 8 ft. Keep room closed for 15 minutes after spraying. Ventilate room for 5 minutes before entry. **FLIES, GNATS, FLYING MOTHS, exposed stages of Indian Meal Moths, Angoumois Grain Moths, Tobacco Moths, Clothes Moths, Chocolate Moths, Almond Moths, Miller Moths, and other listed flying insects:** Direct spray into all parts of the room, paying special attention to hiding places. Sweep up and discard fallen insects.

SPACE TREATMENT OUTDOORS: Do not make applications during rain. Do not wet plants to the point of runoff or drip. Do not water the treated area to the point of run-off. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: (1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf, and other vegetation; (3) Applications to building foundations, up to to a maximum height of 3 feet. Other applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

GNATS, MIDGES, FLIES, and FLYING INSECT CONTROL: Use prior to entry outdoors to reduce annoyance from these and other flying pests. Spray around patios, porches, gazebos, shrubbery, vegetation, and any other listed areas where these insects may rest. For best results, treat when wind velocity does not exceed 5 miles per hour.

EXPOSED STAGES OF Indian Meal Moths, Angoumois Grain Moths, Tobacco Moths, Chocolate Moths, Almond Moths, Millers, and other flying moths: Direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Infested stored products should be fumigated or treated with EPA approved products.

FOR USE IN CANNERIES: Before bringing in produce, thoroughly clean floors, conveying equipment, and walk, paying special attention to cracks and crevices. Spray entire space inside the cannery, including walls, ceilings, and floors, paying special attention to forcing the spray into all cracks and crevices for the control of Fruit Flies, Moths, House Flies, and other listed insects.

FOR USE IN STORAGE SITES: This product can be used in warehouse bins and trucks, cargo ships, mills, bin hoppers, elevators, and conveying equipment as a clean up prior to using them for storage. In mills and elevators, all infested accumulations of grain should be removed from the bin hoppers. All storage and conveying equipment should be thoroughly cleaned by sweeping out waste grain, cobwebs, and other debris from the walls, rafters, floors, and door frames, paying special attention to material lodged in cracks and crevices. All debris should be removed and burned to kill eggs and insects that might be present. **FOR FARMS,** particular attention should be given to cleaning up grain residues from wagons, harvesting equipment and feed troughs, and around used feed and grain bags. Newly harvested grain should not be placed in the same bin with carry-over grain, and all carry-over grain stocks that are not treated with grain protectant should be fumigated. These cleaning operations should be done 2-3 weeks before harvest. After above sanitation measures have been employed, spray all areas prior to use for storage as directed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in a cool dry area. Always store pesticides in the original container. Store away from food and pet food.

Pesticide Disposal and Container Handling

Do Not Puncture or Incinerate! Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions.

Sold By **US Chemical Corporation**
316 Hart Street Watertown, WI 53094-6631