



LICENSED

PERIOD 2011-2013 LIC. NO.



LICENSED

PERIOD 2014-2016 LIC. NO.

9026.47

This product is designed specifically as a ready-to-use cleaner and disinfectant for use in hospitals, nursing homes, shower rooms, locker rooms, public restrooms, restaurants, office buildings, retail businesses, veterinary clinics and pet shops where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, glazed porcelain, glazed ceramic tile, plastic surfaces, bathrooms, shower stalls and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances and stovetops. A rinse with potable water is required for surfaces in direct contact with food.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces.

DISINFECTION, DEODORIZING AND CLEANING: Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 3 minutes. (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

ONE-STEP CLEANING AND DISINFECTION OF TOILET BOWLS AND URINALS: Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of this product evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes) and flush.

TO DISINFECT FOOD SERVICE ESTABLISHMENT FOOD CONTACT SURFACES: Before using this product food products and packaging materials must be removed or carefully protected. For countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow product to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

TUBERCULOCIDAL ACTIVITY - This product exhibits disinfectant efficacy against Mycobacterium tuberculosis H37Rv at 20 degrees Centigrade when the treated surface is allowed to remain wet for 5 minutes.

*VIRUCIDAL ACTIVITY - This product kills on hard, non-porous inanimate surfaces when allowed to remain wet for a 10 minute contact time against: Hepatitis A Virus (HAV), Canine Parvovirus and Poliovirus Type 1.

This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: Hepatitis B Virus (HBV), Hepatitis C Virus (HCV) and Bovine Viral Diarrhea Virus (BVDV) when the treated surface is allowed to remain wet for 5 minutes; Human Coronavirus, SARS associated Coronavirus and Avian Influenza A (Avian Influenza Strains H5N2 and H9N2) for 2 minutes; Rhinovirus type 39 (ATCC VR-340) and Rotavirus for 3 minutes; HIV-1 (associated with AIDS) for 1 minute; and Norovirus (Norwalk virus), Feline Calicivirus and Rabies Virus for 30 seconds.

KILLS HIV-1, HBV, AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

READY-TO-USE DISINFECTANT CLEANER

Cleaner - Deodorizer - Disinfectant
Mildewstat (on hard inanimate surfaces)
*Virucide - Tuberculocidal - Fungicide (against pathogenic fungi)



EPA Reg. No. 1839-83-7546

EPA Est. No. 7546-WI-1

ACTIVE INGREDIENTS:

Table with 2 columns: Ingredient name and percentage. Includes n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18), dimethyl benzyl ammonium chlorides (0.105%), n-Alkyl (68% C12, 32% C14), dimethyl ethylbenzyl ammonium chlorides (0.105%), and TOTAL (99.790% and 100.000%).

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

NET CONTENTS:

1 qt. (946 mL)

www.uschemical.com

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings. CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. CONTACT TIME: Allow surface to remain wet for HBV and HCV 5 minutes, for HIV-1 1 minute. These contact times will not control other common types of viruses and bacteria. DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

FUNGICIDAL ACTIVITY - This product is fungicidal against the pathogenic fungus, Trichophyton mentagrophytes (Athlete's Foot Fungus), when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

Disinfection of animal quarters and kennels: Before using this product, remove all animals and feed from premises. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Apply this product for disinfection of pre-cleaned animal quarters and kennels. Saturate the surfaces with the product for a period of 3 minutes (for Bovine Viral Diarrhea Virus (BVDV) 5 minutes)(for Feline Calicivirus and Rabies virus 30 seconds)(for Canine Parvovirus 10 minutes). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse. Using approved AOAC test methods under Good Laboratory Practices (GLPs), in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, this product kills the following organisms on hard non-porous inanimate surfaces: BACTERICIDAL ACTIVITY (BACTERIA): Pseudomonas aeruginosa, Listeria monocytogenes, Staphylococcus aureus, Yersinia enterocolitica, Salmonella (choleraesuis) enterica, Enterococcus faecium, Escherichia coli, Corynebacterium ammoniagenes, Escherichia coli O157:H7, Salmonella (typhi) enterica, Streptococcus pyogenes (Necrotizing Fasciitis - Group A) (V.A. Medical Center Isolate 04001) ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecalis (VRE), Vancomycin intermediate resistant Staphylococcus aureus (VISA), (CDC Isolate 99287), Methicillin resistant Staphylococcus epidermidis (MRSE).

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - Nonrefillable container. Do not reuse or refill container. Clean container promptly after emptying. Triple rinse as follows: Fill container 1/4 full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.

For Medical Emergency ONLY call: 24 Hour Emergency Response Number 1-866-923-4913 For Product Information, Contact Your Local Distributor



Distributed by: U S Chemical Corporation 316 Hart Street Watertown, WI 53094



0 97146 02183 9

5576076 1A 11/13/7