

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds. Dogs, cats, and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

**Important: Do not expose children, pets, or nontarget animals to rodenticides. To reduce exposure:**

- Store product not in use in a location out of reach of children and pets.
- Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. All above ground, outdoors applications must be in tamper resistant bait stations. These stations must be resistant to destruction by dogs and by children under 6 years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
- Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor above ground use. Tamper resistant bait stations must be used if children, pets, non-target mammals or birds may access the bait.

#### USE RESTRICTIONS

This product may only be used to control Norway rats, roof rats and house mice *in and within* 100 feet of agricultural buildings and man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or harboring or attracting rodent infestations. Fence and perimeter baiting beyond 100 feet from and agricultural structure as defined above is prohibited. This product must not be applied directly to food or feed crops. Do not broadcast bait.

Do not use in homes or other human residences. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs food or feed handling equipment, milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities this product must be applied in tamper resistant bait stations. Do not use in sewers.

This product may also be applied to active rodent burrows to control Norway rats and roof rats *within* 100 feet of buildings and man-made structures provided that infestations of these rodents have been confirmed.

Efforts should be made to remove food trash, garbage, clutter, and debris.

Bait must be placed not less than 6 inches into active Norway/roof rat burrows.

Do not broadcast bait.

Do not sell this product in individual containers holding less than 8 pounds of bait.

#### SELECTION OF TREATMENT AREAS

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much alternative food as possible.

#### APPLICATION DIRECTIONS

##### Norway and Roof Rats

Apply 4 to 16 oz. of bait (usually at intervals of 15-30 ft.) per placement. Bait must be placed not less than 6 inches into active Norway/roof rat burrows. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

##### House Mice

Apply ¼ - ½ oz of bait per placement. Space placements at intervals of 8-12 ft. Larger placements (up to 2 oz) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

##### Follow-up

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water and harborage as much as possible. Where a continuous source of infestations is present maintain permanent bait stations and replenish with census or toxic bait as needed.

# Havoc® Rodenticide Pellets Kills Rats and Mice

**For Control of Commensal Rats and House Mice  
Second Generation Anticoagulant**

Kills Warfarin-Resistant Norway Rats and House Mice

Norway rats and house mice may consume a lethal dose in one night's feeding, with first dead rodents appearing four or five days after feeding begins.

**For Use In and Around Agricultural Buildings Only**

**Distribution to and sales in “consumer” stores including grocery stores, drug stores, hardware stores, club stores is prohibited.**

#### ACTIVE INGREDIENT:

Brodifacoum -  
 3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-tetrahydro-1-naphthalenyl]-4-hydroxy-2H-1-benzopyran-2-one.....0.005%  
**OTHER INGREDIENTS:**..... 99.995%  
**TOTAL** ..... 100.000%

EPA Reg. No. 100-1052-61282

EPA Est. No. 61282-WI-01

## KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements and directions for use on label.

#### PRECAUTIONARY STATEMENTS

##### Hazard to humans and domestic animals

##### CAUTION:

Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin, or on clothing.

All handlers (including applicators) must wear long sleeve shirt long pants, shoes plus socks and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

##### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying the bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

<b>FIRST AID</b> <b>HAVE THE LABEL WHEN OBTAINING TREATMENT ADVICE</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center, doctor, or <b>1-800-498-5743</b> immediately for treatment advice.</li> <li>• Have a person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 – 20 minutes.</li> <li>• Call a poison control center, doctor, or <b>1-800-498-5743</b> immediately for treatment advice</li> </ul>
<b>HOT LINE NUMBER</b> For 24-Hour Medical Emergency Assistance (Human or Animal) Call <b>1-800-498-5743</b> For Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call <b>1-800-424-9300</b>	
<b>NOTE TO PHYSICIAN</b> If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K <sub>1</sub> intramuscularly or orally. Vitamin K <sub>1</sub> is antidotal at doses of 10-20 mg ( <b>not</b> mg/kg). Repeat as necessary based on monitoring of prothrombin times.	
<b>TREATMENT FOR PET POISONING</b> If animal eats bait, call veterinarian or <b>1-800-498-5743</b> at once. <b>NOTE TO VETERINARIAN</b> <b>Anticoagulant Brodifacoum:</b> For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K <sub>1</sub> at doses of 2-5 mg/kg. If needed, check prothrombin times every 3 day until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.	

Manufactured for:  
HACCO, Inc.  
110 Hopkins Drive  
Randolph, WI 53956

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Dispose of wastes resulting from the use of this product at an approved waste disposal facility or call your local solid waste agency for alternative disposal instructions. Never dispose of unused product down any indoor or outdoor drain.

**Container Handling:** Non refillable container. Do not reuse or refill this container. When completely empty, offer for recycling is available, or dispose of empty container in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of HACCO, Inc. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold HACCO, Inc. and Seller harmless for any claims relating to such factors. HACCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or HACCO, Inc., and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, HACCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall HACCO, INC. be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF HACCO, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF HACCO, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

HACCO, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitations of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of HACCO, Inc.

For non-emergency (e.g., current product information)  
call: **1-800-621-8829**

Code (NT.RE.) NL.RG.

Havoc® is a registered trademark of Syngenta Group Company.  
Bitrex® is a registered trademark of Macfarlan Smith Ltd., Edinburgh, Scotland.



**LICENSED**

**PERIOD 2014-2016 LIC. NO. 9084.17**  
**Net Contents: 50 pounds**

**Net Weight: 50 pounds**