




LICENSED

PERIOD 2014-2016 LIC. NO.

9113.114
Disc.



2002-2004



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2008-2010 LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READY TO USE -- No dilution necessary, see table below.

Animal Treatments

Lactating and Non-Lactating Dairy Cattle, Beef Cattle, and Calves

Target Insects: Controls Horn Flies, Face Flies, Lice, and Aids in the Control of Horse flies, Stable Flies, and House Flies.

Application: *Apply along back and down face. Use 1/2 fl oz (15 mL) per 100 lbs. body wt. of animal, up to 5 fl oz for any one animal.

Sheep

Target Insects: Sheep Keds, Lice

Application: * Apply along back. Use 1/4 fl oz (7.5 mL) per 50 lbs. body wt. of animal, up to a maximum of 3 fl oz for any one animal.

Goats

Target Insect: Lice

Application: * Apply along back. Use 1/4 fl oz (7.5 mL) per 50 lbs. body wt. of animal, up to a maximum of 3 fl oz for any one animal.

Repeat treatment as needed, but not more than once every 2 weeks. For optimum lice control a double treatment at 14 day intervals is recommended.

**Duration of activity for horn fly control may vary due to pyrethroid resistance, environmental conditions and management factors (i.e.; sanitation, animal husbandry, facilities, etc.)

Special Note: DURASECT Long-Acting Livestock Pour-On is not effective in controlling cattle grubs.

*DURASECT Long-Acting Livestock Pour-On is to be used in conjunction with the DURASECT application gun or dipper. Do not use this product as a spray.



Durasect

Long-Acting Livestock Pour-On

- Pour-on for beef cattle, lactating and non-lactating dairy cattle, sheep and goats
- Patent-pending ready-for-use water repellent formulation for use on livestock
- No withdrawal time
- Does not leave an oily residue
- 4 to 11 weeks control of horn flies**
- Controls face flies for at least 1 month
- Controls lice

Active Ingredients:


*Permethrin (3-phenoxyphenyl) methyl ± cis, trans-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate 1.0%
 INERT INGREDIENTS 99.0%
 *cis/trans ratio: Min 35% (±)cis and max 65% (±)trans

Keep Out of Reach of Children

CAUTION

See Side Panel for Precautionary Statements

Net Contents: 1 gallon



Regis.

1005-5002

411-811P 1005-5001

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

Statement of Practical Treatment

If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If in Eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If Swallowed: Call a physician immediately.

Environmental Hazards

This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not add directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product as specified on the label.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Keep container sealed when not in use. Do not store near food or feed.

Disposal: Triple rinse (or equivalent). Then offer for recycling, or puncture and discard in trash. The container is made of High Density Polyethylene (HDPE) #2 and can be offered for recycling.

Distributed by:



Animal Health

Kenilworth, PA 19031, USA
Div. of Pfizer Inc., NY, NY 10017

EPA Est. No. 54487-GA-1

EPA Reg. No. 1007-90

03-0649-00-0

Product