

Detail[®]

Group 14 Herbicide



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

9131.159

Powered by Kixor[®] Herbicide

For postemergence and residual weed control in noncropland areas

Active Ingredient*:

saflufenacil: N'-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-(trifluoromethyl)-3,6-dihydro-1(2H)-pyrimidinyl)benzoyl]-N-isopropyl-N-methylsulfamide..... 29.74%

Other Ingredients: 70.26%

Total: 100.00%

* Contains 2.85 pounds active ingredient saflufenacil per gallon formulated as a water-based suspension concentrate

EPA Reg. No. 7969-297

EPA Est. No. 241-PR-002

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents: 1 Gallon

81066058 NVA 2014-05-324-0591
Product of U.S.A.

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

 **BASF**
We create chemistry



FIRST AID

If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give any liquid to the person.• DO NOT give anything to an unconscious person.
If in eyes	<ul style="list-style-type: none">• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eyes.• Call a poison control center for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.• Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).	

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (such as natural rubber)
- Protective eyewear such as face shield, goggles, or safety glasses

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for **applicators and other handlers** and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses, **DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

Groundwater Advisory. Saflufenacil has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory. Saflufenacil may impact surface water due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several weeks after application. A level, well-maintained buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this chemical from runoff water and sediment. Runoff of this product will be reduced by avoiding application when rainfall is forecast to occur within 48 hours.

Endangered Species Protection Requirements

This product may have effects on federally listed threatened or endangered plant species or their critical habitat. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-800-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will generally be available from the above sources 6 months prior to their effective dates.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at time of herbicide application.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Observe all precautions and limitations in this label and the labels of products used in combination with **Detail® herbicide**. The use of **Detail** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions, and **Conditions of Sale and Warranty** are to be followed.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

EXCEPTION: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as natural rubber \geq 14 mils
- Shoes plus socks
- Protective eyewear

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter treated areas without protective clothing until sprays have dried.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

DO NOT use or store near heat or open flame. Store in original container in a well-ventilated area separately from fertilizer, feed, or foodstuffs. Avoid cross-contamination with other pesticides.

Pesticide Disposal

Wastes resulting from this product may be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity \leq 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Steps to take if material is released or spilled:

- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

Product Information

Detail® herbicide provides both contact burndown (postemergence) and rate-dependent residual preemergence weed control (refer to **Table 1** and **Table 2** for list of weeds controlled). **Detail** does not control grass weeds and must be tank mixed with a grass herbicide for a complete weed control program.

Good coverage of weed foliage is critical for optimum postemergence control. Higher spray volumes should be used when targeting larger weeds or higher weed densities.

Postemergence applications of **Detail** should be made when weeds are small (less than 6 inches) and actively growing. An adjuvant is required with **Detail** for optimum burndown activity (refer to **Additives** section for details). Burndown activity may be slowed or reduced under cloudy and/or foggy or cooler weather conditions, or when weeds are growing under drought or other stress conditions. When targeting dense weed populations and/or larger weeds, use higher spray volumes and/or a higher application rate within an application rate range. Angling nozzles forward (to 45 degrees) may improve penetration of denser weed canopies.

Length of weed control from residual preemergence applications will be affected by use rate, soil characteristics (texture, organic matter, cation exchange capacity), as well as the amount of rainfall after application.

Table 1. Weeds Controlled with a Postemergence Application of Detail® herbicide¹

Common Name	Scientific Name
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Bedstraw, catchweed	<i>Galium aparine</i>
Beggarticks, hairy	<i>Bidens pilosa</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Bindweed, field ³	<i>Convolvulus arvensis</i>
Buckwheat, wild	<i>Polygonum convolvulus</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed, common	<i>Stellaria media</i>
Cocklebur, common	<i>Xanthium strumarium</i>
Cowcockle	<i>Vaccaria pyramidata</i>
Dandelion ³	<i>Taraxacum officinale</i>
Eveningprimrose, cutleaf	<i>Oenothera laciniata</i>
Falseflax, smallseed	<i>Camelina microcarpa</i>
Filaree, redstem	<i>Erodium cicutarium</i>
Fleabane, hairy	<i>Conyza bonariensis</i>
Flixweed	<i>Descurainia sophia</i>
Groundcherry, cutleaf	<i>Physalis angulata</i>
Groundsel, common	<i>Senecio vulgaris</i>
Henbit ²	<i>Lamium amplexicaule</i>
Horseweed (marestail)	<i>Conyza canadensis</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lambsquarters, narrowleaf	<i>Chenopodium pratericola</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Mallow, common	<i>Malva neglecta</i>
Mallow, little (cheeseweed)	<i>Malva parviflora</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Marestail (horseweed)	<i>Conyza canadensis</i>
Morningglory, entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>
Morningglory, palmleaf	<i>Ipomoea wrightii</i>

(continued)

Table 1. Weeds Controlled with a Postemergence Application of Detail® herbicide¹
(continued)

Common Name	Scientific Name
Morningglory, pitted	<i>Ipomoea lacunosa</i>
Morningglory, tall	<i>Ipomoea purpurea</i>
Mustard, black	<i>Brassica nigra</i>
Mustard, tumble	<i>Sisymbrium altissimum</i>
Mustard, wild	<i>Sinapis arvensis</i>
Nettle, burning	<i>Urtica urens</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, cutleaf	<i>Solanum triflorum</i>
Nightshade, Eastern black	<i>Solanum ptycanthum</i>
Nightshade, hairy	<i>Solanum saccharoides</i>
Pennycress, field	<i>Thlaspi arvense</i>
Pigweed, prostrate	<i>Amaranthus blitoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pine ⁴	<i>Pinus</i> spp.
Puncturevine	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida ²	<i>Richardia scabra</i>
Ragweed, common ⁵	<i>Ambrosia artemisiifolia</i>
Ragweed, giant ⁵	<i>Ambrosia trifida</i>
Rapeseed (canola), volunteer	<i>Brassica</i> spp.
Sesbania, hemp	<i>Sesbania exaltata</i>
Shepherd's-purse	<i>Capsella bursa-pastoris</i>
Sida, prickly	<i>Sida spinosa</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Sowthistle, annual	<i>Sonchus oleraceus</i>
Sowthistle, spiny	<i>Sonchus asper</i>
Spurge, leafy ⁶	<i>Euphorbia esula</i>
Sunflower, common	<i>Helianthus annuus</i>
Tansymustard, pinnate	<i>Descurainia pinnata</i>
Thistle, Canada ³	<i>Cirsium arvense</i>

(continued)

Table 1. Weeds Controlled with a Postemergence Application of Detail® herbicide¹
(continued)

Common Name	Scientific Name
Thistle, Russian	<i>Salsola kali</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp ⁵	<i>Amaranthus tuberculatus</i>
Willowweed	<i>Epilobium adenocaulon</i>

¹ For best control, target applications when weeds are less than 6 inches. Larger weeds or heavy infestations will require higher use rates (see **Table 3**) or tank mixes.

² Suppression only

³ Control of seedling stage and suppression of perennial growth stage

⁴ See **Right of Way** specific use pattern directions for additional information. Tank mix partners, such as glyphosate, are required.

⁵ Populations of noted weeds exist that are known to be resistant to burndown applications of **Group 14/Group E** herbicides and will not be controlled by herbicides like **Detail**. See the **Resistance Management** section for practices to manage and minimize the impact of resistant weeds (e.g. tank mixes or alternation with other herbicide modes of action and mechanical control).

⁶ Control of leafy spurge requires tank mix with **Plateau® herbicide**.

Table 2. Preemergence Weed Control with Detail® herbicide¹

Common Name	Scientific Name
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Powell	<i>Amaranthus powellii</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Buckwheat, wild	<i>Polygonum convolvulus</i>
Burcucumber ²	<i>Sicyos angulatos</i>
Chickweed, common	<i>Stellaria media</i>
Cocklebur, common	<i>Xanthium strumarium</i>
Copperleaf, Virginia	<i>Acalypha virginica</i>
Galinsoga, smallflower	<i>Galinsoga parviflora</i>
Groundcherry, cutleaf	<i>Physalis angulata</i>

(continued)

Table 2. Preemergence Weed Control with Detail® herbicide¹ (continued)

Common Name	Scientific Name
Horseweed (marestail)	<i>Coryza canadensis</i>
Jimsonweed	<i>Datura stramonium</i>
Kochia	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Morningglory, entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>
Morningglory, pitted	<i>Ipomoea lacunosa</i>
Morningglory, tall	<i>Ipomoea purpurea</i>
Mustard, wild	<i>Sinapis arvensis</i>
Nightshade, black	<i>Solanum nigrum</i>
Pennycress, field	<i>Thlaspi arvense</i>
Pigweed, prostrate	<i>Amaranthus blitoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Puncturevine ²	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Pusley, Florida	<i>Richardia scabra</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Sida, prickly	<i>Sida spinosa</i>
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Starbur, bristly	<i>Acanthospermum hispidum</i>
Sunflower, common	<i>Helianthus annuus</i>
Thistle, Russian	<i>Salsola kali</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp	<i>Amaranthus tuberculatus</i>

¹ For effective **residual** preemergence weed control from postemergence applications, **Detail** must be used at the maximum use rate of 6 fl ozs/A (see **Table 3**) and be activated by a minimum of 1/2 inch of rainfall before weed seedling emergence. When **Detail** is not activated, a labeled postemergence herbicide may be required to improve weed control.

² Suppression only

Mode of Action

Detail is a potent inhibitor of protoporphyrinogen-oxidase belonging to herbicide mode of action **Group 14** (WSSA)/**Group E** (HRAC). **Detail** is rapidly absorbed by roots and foliage. Following inhibition of protoporphyrinogen-oxidase, plant death is the result of membrane damage. Under active growing conditions, susceptible emerged weeds usually develop chlorotic and necrotic injury symptoms within hours and die within a few days. Susceptible emerging weed seedlings will usually die as they reach the soil surface or shortly after emergence.

Resistance Management

While weed resistance to protoporphyrinogen-oxidase inhibiting herbicides is relatively infrequent, populations of resistant biotypes are known to exist. Resistance management practices include:

1. Following labeled application rate and weed growth stage recommendations
2. Avoiding repeated applications of herbicides with the same mode of action
3. Using tank mixes and sequential applications with other effective herbicides possessing different modes of action

Application Instructions

Detail may be applied in a single application or sequentially with an interval of 14 days or more.

Application Rates

Application rates for **Detail** when applied alone, in tank mix, or sequentially are given in **Table 3**. **DO NOT** apply more than a maximum cumulative amount of 6 fl ozs/A of **Detail** per year.

Table 3. Application Rates for Detail® herbicide

Application	Application Target	Application Rate (fl ozs/A)
Postemergence	Weed size < 6 inches	2 to 4
	Weed size ≥ 6 inches and/or heavier weed infestations	4 to 6 ^a
Postemergence + Residual	Burndown + Residual pre-emergence weed control	6 ^b
Tank Mixes with Glyphosate		
Accelerated Burndown	Accelerated burndown of weeds and/or control of glyphosate-tolerant species [such as horseweed (marestalk)]	1 to 5
Accelerated Burndown + Residual	Accelerated burndown of weeds plus control of glyphosate-tolerant species with residual pre-emergence weed control	6 ^b

^a Partial control or suppression may result with application to weeds greater than 6 inches.

^b For effective residual control of labeled weed species, **Detail** must be used at the maximum use rate of 6 fl ozs/A.

Application Methods and Equipment

Detail may be applied by ground or air. Good spray coverage is important for optimum weed control and can be improved with proper adjuvant, nozzle, and spray volume selection.

Use and configure application equipment to provide an adequate spray volume, an accurate and uniform distribution of spray droplets over the treated area, and to avoid spray drift to nontarget areas. Equipment should be adjusted to maintain continuous agitation during spraying with good mechanical or bypass agitation. Avoid overlaps that will increase rates above the use rates specified in this label.

Aerial Application Requirements

Water Volume. Use 10 or more gallons of water per acre.

The following measures must be followed to reduce the potential of spray drift to nontarget areas from aerial application:

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the fixed wingspan or 90% of rotor blade diameter.
2. Use low-drift nozzles such as straight-stream nozzles (D-8 or larger). **DO NOT** use nozzles producing a mist droplet spray.
3. Nozzles must always point backward parallel with the airstream and never be pointed downward more than 45 degrees.
4. Without compromising aircraft safety, applications should be made at a height of 10 feet or less above the crop canopy or tallest plants.
5. **DO NOT** apply during periods of temperature inversions or stable atmospheric conditions.
6. Avoid potential adverse effects to nontarget areas by maintaining a (xx)^a foot buffer between the point of direct application and the **closest downwind edge** of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, and shrub lands).

^a The buffer zone size is determined by use rate. Refer to the table below for the minimum buffer zone distance required for the intended use rate. Utilize the appropriate buffer zone distance from the table below in the buffer zone statement above.

Detail® herbicide Use Rate (fl ozs/A)	Buffer Zone Distance (feet)
1	26
2	66
4	100
6	150

Ground Application (broadcast) Requirements

Water Volume. Use 20 or more gallons of water per acre.

The following measures must be followed to reduce the potential of spray drift to nontarget areas from ground application:

1. Apply this product using nozzles which deliver **medium-to-coarse spray droplets** as defined by ASAE standard S-572 and as shown in nozzle manufacturer's catalogs. Flat-fan nozzles are recommended for burndown applications while flood-jet type nozzles are recommended for residual soil surface applications. Nozzles that deliver coarse spray droplets may be used to reduce spray drift provided spray volume per acre (GPA) is increased to maintain coverage of target (i.e. weeds or soil surface). **DO NOT** use nozzles that produce fine (e.g. cone) spray droplets.
2. Apply this product only when the potential for drift to adjacent nontarget areas is minimal (e.g. when the wind is **10 MPH or less and is blowing away** from sensitive areas). **DO NOT** apply during periods of temperature inversions or stable atmospheric conditions.
3. Avoid potential adverse effects to nontarget areas by maintaining a (xx)^a foot buffer between the application area and the **closest downwind edge** of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, and shrub lands).

^a The buffer zone size is determined by use rate. Refer to the table below for the minimum buffer zone distance required for the intended use rate. Utilize the appropriate buffer zone distance from the table below in the buffer zone statement above.

Detail Use Rate (fl ozs/A)	Buffer Zone Distance (feet)
1	13
2	33
4	50
6	75

Ground Application (spot)

Postemergence spot applications may be made with **Detail**. Spray volumes should be sufficient to thoroughly wet targeted foliage but not to the point of runoff, i.e. a spray-to-wet basis. Use a 0.25% to 0.50% volume/volume (v/v) spray solution for control of weeds less than 6 inches. For larger weeds or under heavy weed infestations, increase the spray solution to 0.50% to 1.00% v/v. Spot applications will also require the use of an adjuvant; add methylated seed oil (MSO) at the rate of 1% v/v. The following chart provides the amount(s) of **Detail** to use to prepare spray solutions for spot applications.

Detail Required (fl ozs) for Spot Application Treatments				
Spray Solution to Prepare (gals)	Desired Concentration (v/v)			
	0.25%	0.50%	0.75%	1.0%
1	0.3	0.6	1.0	1.3
3	1.0	1.9	2.9	3.8
4	1.3	2.6	3.8	5.1
5	1.6	3.2	4.8	6.4
50	16.0	32.0	48.0	64.0
100	32.0	64.0	96.0	128.0

2 tablespoons = 1 fluid ounce, 1 pint = 16 fluid ounces, 1 quart = 32 fluid ounces, 1 gallon = 128 fluid ounces

Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions, followed by triple rinsing the equipment before and after applying this product.

Spray Drift Management

It is the responsibility of the applicator to avoid spray drift at the application site, especially onto nontarget areas. The interaction of many equipment-related and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The applicator should be familiar with and take into account the information covered in the following spray drift reduction advisory information.

Controlling Droplet Size. The most effective way to reduce drift potential is to apply the largest droplets that provide sufficient coverage and control.

Volume. Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure. DO NOT exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles. Use the minimum number of nozzles that provide uniform coverage.

Nozzle Type. Use a nozzle type designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets.

Swath Adjustment. When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the application equipment (e.g. aircraft, ground) upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

Wind. Drift potential is lowest between wind speeds of 3 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. If applying at wind speeds less than 3 mph, the applicator must determine if:

1. Conditions of temperature inversion exist, or
2. Stable atmospheric conditions exist at or below nozzle height.

DO NOT make applications into areas of temperature inversions or stable atmospheric conditions.

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Wind Erosion. Avoid treating powdery, dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

Additives

For optimum burndown activity with **Detail® herbicide** and to achieve consistent weed control in post-emergence use patterns, an adjuvant system must be used that includes the following:

Adjuvant ¹	Rate
Methylated seed oil (MSO) ²	1 gal/100 gals (1% v/v) ³

¹ Use of ammonium sulfate (AMS) fertilizer at 8.5 to 17 lbs/100 gallons (1% to 2% weight/volume) or urea ammonium nitrate (UAN) at 1.25 to 2.5 gals/100 gals (1.25% to 2.5% v/v) is highly recommended when mixing **Detail** with glyphosate-based herbicides.

² MSO-based adjuvant **MUST** contain at least 60% methylated seed oil. Poor performance may occur with adjuvants containing less than 60% methylated seed oil.

³ **DO NOT** use less than 1 pint/A of MSO with low-volume (< 12.5 gallons per acre) aerial or ground applications.

DO NOT use a nonionic surfactant (NIS) or a crop oil concentrate (COC) as a substitute for MSO, or poor performance on weeds will occur.

Tank Mixing Information

Detail may be tank mixed with one or more registered herbicide products according to the specific tank mixing instructions in this label and respective product labels. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Always follow the most restrictive label use directions. Refer to **Specific Use Information** section for details.

Tank mixtures with contact herbicides (e.g. carfentrazone, paraquat) may reduce the burndown activity of **Detail**.

Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

1. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.
2. Add components in the sequence indicated in the **Mixing Order** section using 2 teaspoons for each pound or 1 teaspoon for each pint of label use rate per acre.
3. Always cap the jar and invert 10 cycles between component additions.
4. When the components have all been added to the jar, let the solution stand for 15 minutes.
5. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, or fine particles that precipitate to the bottom, or thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, **DO NOT** mix the ingredients in the same tank.

Mixing Order

1. **Water** - Fill tank 1/2 to 3/4 full with clean water and start agitation.
2. **Agitation** - Maintain continuous and constant agitation throughout mixing.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-soluble additives** (including dry and liquid fertilizers such as ammonium sulfate or urea ammonium nitrate)
6. **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
7. **Water-soluble products**
8. **Emulsifiable concentrates** (including methylated seed oil adjuvants)
9. **Remaining quantity of water**

Maintain agitation throughout application until spraying is completed. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

Use Precautions

- **Maximum annual use rate** - **DO NOT** apply more than a maximum cumulative amount of 6 fl ozs/A of **Detail® herbicide** (0.134 pound active ingredient saflufenacil per acre) per year from broadcast or banded applications.
- **Rainfastness** - **Detail** is rainfast 1 hour after application. Burndown activity may be reduced if rain occurs within 1 hour of application.
- **DO NOT** contaminate irrigation ditches or water used for domestic purposes.
- **DO NOT** apply through any type of irrigation system (e.g. chemigation).
- **Detail is not for sale, distribution, or use in Nassau and Suffolk counties in New York State.**

Specific Use Information

Detail may be used for selective or nonselective weed control in conifer and hardwood plantations and various noncropland areas. This section provides use directions for **Detail** in various noncrop situations. Be sure to read product information, mixing, application, weeds controlled and adjuvant instructions in preceding sections of the label. Read and follow tank mix product labels for restrictions, precautions, instructions, and rotational crop restrictions.

Conifer and Hardwood Plantations

Application Method, Rate, and Timing

Apply **Detail** for the control of undesirable plants during site preparation operations conducted before the planting and establishment of conifer and hardwood plantations. Refer to **Table 1**, **Table 2**, and **Table 3** for lists of weeds controlled and application rates.

DO NOT apply **Detail** as an over-the-top spray on desirable conifer or hardwood plantings or severe injury will occur.

DO NOT plant tree seedlings within 2 months after **Detail**® herbicide application.

Site Preparation Application

Apply **Detail** plus the recommended adjuvant (refer to **Additives** section for details) as a uniform broadcast application during preplant site preparation for the control of undesirable weed species in plantations and for enhanced brownout with other site preparation tank mixes.

Wildling Pine Control

For best control of wildling pine, tank mix **Detail** with glyphosate (refer to specific glyphosate label for appropriate use rates) in addition to other tank mix herbicides listed following, and make foliar applications in the spring to summer when wildling pine seedlings are actively growing. Fall applications may not provide consistent control. Proper spray coverage is essential for best control. Use a minimum spray volume of 15 gallons of water per acre for aerial applications. For ground applications, use a minimum spray volume of 30 gallons of water per acre for broadcast foliar applications to provide maximum spray coverage.

Tank Mixtures

Broad-spectrum burndown and/or residual control of grasses, pine, or additional broadleaf species will require a tank mix with another herbicide. **Detail** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Arsenal**® herbicide Applicators Concentrate
- **Chopper**® Gen2™ herbicide
- glyphosate (e.g. **Accord**® XRT herbicide)

Noncropland Areas

Detail may be used in noncropland areas including fence rows, nonirrigation ditchbanks, petroleum tank farms, pumping installations, railroad, rights-of-way (utility, pipeline, highway), storage areas, and utility plant sites.

Application Method, Rate, and

Timing

Selective Weeding

Apply **Detail** as a postemergence spray plus the recommended adjuvant (refer to **Additives** section for details) as a uniform broadcast application for selective

weed control. Refer to **Table 1**, **Table 2**, and **Table 3** for lists of weeds controlled and application rates.

Tank Mixtures. Broad-spectrum postemergence and/or residual control of grasses or additional broadleaf species will require a tank mix with another herbicide. **Detail** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Overdrive**® herbicide
- **Plateau**® herbicide
- glyphosate

Bareground

Detail will provide contact burndown plus residual preemergence control of annual weeds. Apply **Detail** plus the recommended adjuvant (refer to **Additives** section for details) as a uniform broadcast application. To provide effective residual broadleaf weed control, **Detail** must be applied at the maximum use rate of 6 fl ozs/A. The actual length of residual control depends on factors such as application rate, soil type, organic matter, weed pressure, and rainfall amounts after application. Adequate precipitation is necessary to activate **Detail**. Dry weather following application may reduce effectiveness. Refer to **Table 1**, **Table 2**, and **Table 3** for lists of weeds controlled and application rates.

Tank Mixtures. Broad-spectrum postemergence and/or residual control of grasses or additional broadleaf species will require a tank mix with another herbicide. **Detail** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Arsenal**® herbicide
- **Arsenal**® PowerLine™ herbicide
- **Frequency**® herbicide
- **Pendulum**® AquaCap™ herbicide
- **Plateau**® herbicide
- **Sahara**® herbicide
- diuron
- glyphosate

Right of Way

Broadcast Application. Apply **Detail** to provide rapid brownout of volunteer pine, including loblolly pine (*Pinus taeda*) and Virginia pine (*P. virginiana*). For best results, apply **Detail** plus the recommended adjuvant

(refer to **Additives** section for details) as a uniform broadcast application at 3 to 6 fl ozs/A in a tank mix with glyphosate (refer to glyphosate label for specific use rates). Make foliar applications in the spring to summer when volunteer pine are actively growing. Fall applications may not provide consistent control. Use a minimum spray volume of 50 gallons water per acre for broadcast foliar applications to provide the maximum spray coverage.

Selective Stem Application. Apply **Detail**[®] herbicide in a tank mix with glyphosate or other labeled products, plus the recommended adjuvant(s), to provide rapid brownout of woody species using a directed-foliar individual plant treatment. For enhanced brownout of pine species (including loblolly pine [*Pinus taeda*] and Virginia pine [*P. virginiana*]), tank mix with glyphosate or other pine control products. Make selective stem applications of **Detail** using backpack or hydraulic handgun equipment. For best results, apply **Detail** at a rate range of 0.125% to 1.0% v/v with a tank mix partner (refer to tank mix partner label for the recommended use rate). The proper spray pattern for selective stem applications is to uniformly wet all the foliage on the target plant, but **DO NOT** drench the target vegetation causing spray solution to run off. Excessive wetting of foliage to runoff is not recommended. For best results, make selective stem applications with methylated seed oil at 1% v/v as the adjuvant. Apply **Detail** up to but **DO NOT** exceed 16 fl ozs/A (0.356 pound active ingredient saflufenacil per acre) with selective stem applications.

Tank Mixtures. **Detail** may be tank mixed or applied sequentially with one or more of, but not limited to, the following herbicide products:

- **Arsenal**
- **Arsenal PowerLine**
- **Milestone**[®] herbicide
- glyphosate

Leafy Spurge Control

Detail applied in tank mixture with **Plateau**[®] herbicide will control leafy spurge when applied late spring/early summer in noncropland areas as described above. This tank mixture will also control additional weeds listed on the respective **Detail** and **Plateau** labels. **Detail** plus **Plateau** tank mix may be applied by ground or air.

Apply **Detail** at 1.0 to 2.0 fl ozs/A plus **Plateau** at 4.0 to 6.0 fl ozs/A to leafy spurge when it reaches the yellow bract (pre-bloom) stage in late spring/early summer. **DO NOT** apply this tank mix as a fall application because resultant control may not be satisfactory.

DO NOT feed or allow animals to graze areas of grass treated with **Detail** within 365 days of treatment.

Spray Additives for Leafy Spurge Control. **Detail** plus **Plateau** tank mix requires the use of an effective adjuvant system. For best results, use a non-ionic surfactant at 0.25% v/v plus ammonium sulfate at 8.5 to 17.0 lbs/100 gals [1% to 2% weight/volume (w/v)]. Crop oil concentrate or methylated seed oil may also be used with this tank mixture when injury (stunting, necrosis) to grasses is acceptable.

Water Volume for Leafy Spurge Control. For ground applications, use 10 or more gallons of water per acre. Thorough coverage of weeds is essential and higher spray volumes may be necessary for better performance on a heavy population of leafy spurge. For aerial applications, use 5 or more gallons of water per acre.

Native Grass Areas

Detail may be used for the establishment and maintenance of native grass and natural areas (such as wildlife management areas, wildlife openings, wildlife food plots, and wildlife habitats).

Application Method, Rate, and Timing

Apply **Detail** as a postemergence spray plus the recommended adjuvant (refer to **Additives** section for details) as a uniform broadcast application for selective broadleaf weed control in native grass areas and unimproved turf. Refer to **Table 1**, **Table 2**, and **Table 3** for lists of weeds controlled and application rates. Transitory injury may be observed on certain grass species such as Bermudagrass and Bahiagrass at higher use rates.

DO NOT feed or allow domestic animals to graze areas of grass treated with **Detail** within 365 days of treatment.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

***Arsenal, Chopper, Detail, Frequency, Kixor, Overdrive, Pendulum, Plateau, and Sahara** are registered trademarks of BASF.*

***AquaCap, Gen2, and PowerLine** are trademarks of BASF.*

***Accord and Milestone** are registered trademarks of Dow AgroSciences LLC.*

© 2014 BASF Corporation
All rights reserved.

007969-00297.20141104.**NVA 2014-04-324-0182**

Based on: NVA 2014-04-324-0067

Supersedes: NVA 2012-04-324-0032

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry

NOTES

Detail[®]

Group 14 Herbicide

Powered by **Kixor[®]** Herbicide

For postemergence and residual weed control in noncropland areas

Active Ingredient*:

saflufenacil: N'-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-(trifluoromethyl)-3,6-dihydro-1(2H)-pyrimidinyl)benzoyl]-N-isopropyl-N-methylsulfamide..... 29.74%

Other Ingredients:..... 70.26%

Total:..... 100.00%

* Contains 2.85 pounds active ingredient saflufenacil per gallon formulated as a water-based suspension concentrate
EPA Reg. No. 7969-297 EPA Est. No. 241-PR-002

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID: If swallowed: Call a poison control center or doctor immediately for treatment advice. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. **DO NOT** give any liquid to the person. **DO NOT** give anything to an unconscious person. **If in eyes:** Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eyes. Call a poison control center for treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth, if possible. Call a poison control center or doctor for further treatment advice. **HOTLINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357). **Precautionary Statements: Hazards to Humans and Domestic Animals. CAUTION.** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Remove and wash contaminated clothing before reuse. **Environmental Hazards:** For terrestrial uses, **DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. **Groundwater Advisory.** Saflufenacil has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. **Surface Water Advisory.** Saflufenacil may impact surface water due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several weeks after application. A level, well-maintained buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this chemical from runoff water and sediment. Runoff of this product will be reduced by avoiding application when rainfall is forecast to occur within 48 hours.

STORAGE AND DISPOSAL: do NOT contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. **Pesticide Storage:** **DO NOT** use or store near heat or open flame. Store in original container in a well-ventilated area separately from fertilizer, feed, or foodstuffs. **Avoid cross-contamination with other pesticides. Pesticide Disposal:** Wastes resulting from this product may be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Handling: Nonrefillable Container. DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities. See attached booklet for complete container handling directions including triple rinsing and pressure rinsing instructions.

See attached booklet for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific use site restrictions. **In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).**

Net Contents: 1 Gallon
Product of U.S.A.

81066058 NVA 2014-05-324-0591
Product of U.S.A.

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

 **BASF**
We create chemistry

Supplemental Label

Detail®

Powered by **Kixor®** Herbicide

For aerial application by helicopter

This supplemental label expires December 31, 2018, and must not be used or distributed after this date.

Active Ingredient:
saflufenacil: N'-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-(trifluoromethyl)-3,6-dihydro-1(2H)-pyrimidinyl)benzoyl]-N-isopropyl-N-methylsulfamide..... 29.74%

Other Ingredients:..... 70.26%

Total: 100.00%

Contains 2.85 pounds active ingredient saflufenacil per gallon formulated as a water-based suspension concentrate

EPA Reg. No. 7969-297

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Detail® Powered by KIXOR® herbicide** (henceforth in this label referred to as **Detail herbicide**) container label, EPA Reg. No. 7969-297, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Detail herbicide** before applying.
- Use of **Detail herbicide** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Detail herbicide**.

Product Information

Detail herbicide provides wildling pine control and both contact burndown (postemergence) and rate-dependent residual preemergence weed control (refer to **Table 1** and **Table 2** on product container label for list of weeds controlled) in conifer and hardwood plantations, pasture and rangeland, and in vegetation control and management sites in noncropland areas. **Detail herbicide** does not control grass weeds and must be tank mixed with a grass herbicide for a complete weed control program.

Refer to the product container label and follow all application instructions, application rates, application sites, and spray additives.

Aerial Application by Helicopter

Detail herbicide may be applied by air with a helicopter. Thorough spray coverage is important for optimum weed control and can be improved with proper adjuvant, nozzle, and spray volume selection.

Use and configure application equipment to provide an adequate spray volume, an accurate and uniform distribution of spray droplets over the treated area, and to avoid spray drift to nontarget areas. Adjust equipment to maintain continuous agitation during spraying with good mechanical or bypass agitation. Avoid overlaps that will increase rates above use rates specified in the product container label.

Aerial Application Requirements with Helicopter

Water Volume. Use 15 or more gallons of water per acre.

DO NOT apply aerially in California.

The following measures must be followed to reduce the potential of spray drift to nontarget areas from aerial application with helicopter:

1. The distance of the outermost nozzles on the boom must not exceed 75 to 80% of rotor blade diameter.
2. Use Accu-Flo™ .028 nozzles or larger. **DO NOT** use nozzles producing a smaller droplet size than Accu-Flo .028.
3. Orient nozzles so spray is released parallel to the airstream.
4. Without compromising aircraft safety, applications should be made at a height of 10 feet or less above the target vegetative canopy.
5. **DO NOT** apply when wind speed is greater than 10 miles per hour, during periods of temperature inversions or stable atmospheric conditions.
6. Avoid potential adverse effects to nontarget areas by maintaining a (XX, see table below)^a foot buffer between the point of direct application and the **closest downwind edge** of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, and shrub lands).

^a Buffer zone size is determined by use rate. Refer to table below for minimum buffer zone distance required for the intended use rate. Use the appropriate buffer zone distance from the table below in the buffer zone statement above.

Buffer Zone Distances for Helicopter Applications

Detail® herbicide Use Rate (fl ozs/A)	Buffer Zone Distance (feet)
1	5
2	5
4	15
6	30

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Detail and Kixor are registered trademarks of BASF.

Accu-Flo is a trademark of Bishop Equipment Mfg., Inc.

© 2016 BASF Corporation
All rights reserved.

007969-00297.20160106.NVA 2016-04-324-0002
Based on: NVA 2015-04-324-0046

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry

Supplemental Label



For broadleaf weed control in perennial forage grasses grown in rangeland

This supplemental label expires August 31, 2017, and must not be used or distributed after this date.

Active Ingredient:

saflufenacil: N'-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-(trifluoromethyl)-3,6-dihydro-1(2H)-pyrimidinyl)benzoyl]-N-isopropyl-N-methylsulfamide..... 29.74%

Other Ingredients:..... 70.26%

Total: 100.00%

Contains 2.85 pounds active ingredient saflufenacil per gallon formulated as a water-based suspension concentrate

EPA Reg. No. 7969-297

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Detail**™ herbicide container label, EPA Reg. No. 7969-297, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Detail** before applying.
- Use of **Detail** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Detail**.

Use Information

Detail may be applied for broadleaf weed control (refer to **Table 1** and **Table 2** on the product container label for list of weeds controlled) in perennial cool-season and warm-season forage grasses grown in rangeland for livestock grazing.

Before applying **Detail** to forage grasses, verify the selectivity of **Detail** on your variety with your local seed company (supplier) to help avoid potential injury to sensitive varieties.

Application Timings and Rates

Apply **Detail** only to established (defined as planted in fall or spring which has gone through a first cutting/mowing) stands of perennial cool-season and warm-season forage grasses.

Detail may cause transitory injury to forage grasses (leaf necrosis) under certain conditions, but new growth is normal and vigor is not reduced.

Disease, extremely cold weather, drought, extensive frost heaving, low or high pH, salinity, and other environmental pressures may weaken grass stands and make the crop more susceptible to herbicidal injury.

Dormant-season Applications for Burndown and Residual Weed Control in Warm-season and Cool-season Grasses

Apply **Detail** at 1.0 to 2.0 fl ozs/A as a broadcast burndown spray to emerged broadleaf weeds in the dormant season [i.e. when grasses are not actively growing in the fall (postharvest), during the winter dormancy period, or in the early spring before greenup]. An adjuvant system is required for optimum broadleaf burndown activity.

For additional residual broadleaf weed control, **Detail™ herbicide** can be applied anytime in the dormant season (as described above) at rates of 3.0 to 4.0 fl ozs/A.

Sequential applications of **Detail** may be made within the dormant season as long as the maximum cumulative amount does not exceed 4.0 fl ozs/A of **Detail**. Apply dormant-season burndown applications sequentially where the first burndown application is made fall (postharvest) or during the winter dormancy period, and the second application is made in the early spring before greenup. Separate sequential dormant-season burndown applications by at least 14 days.

Specific Adjuvant Requirements for Dormant-season Applications in Warm-season and Cool-season Grasses. For optimum postemergence control of emerged broadleaf weeds, the following adjuvants are required for use with **Detail**:

- Methylated seed oil (MSO) at 1% volume/volume (v/v) plus ammonium sulfate (AMS) at 8.5 to 17.0 lbs/100 gallons

In-season Postemergence Applications for Weed Control in Cool-season Grasses

Apply **Detail** at 1.0 to 2.0 fl ozs/A as a broadcast postemergence spray to control emerged broadleaf weeds in season (i.e. actively growing cool-season forage grasses). Make in-season applications before weeds reach the maximum size listed in **Table 1** on the product container label. Postemergence applications require the addition of an adjuvant system.

Specific Adjuvant Requirements for In-season Postemergence Applications in Cool-season Grasses. For optimum postemergence control of emerged broadleaf weeds, the following adjuvant is required for use with **Detail**:

- MSO at 1% v/v

In-season Postemergence Applications for Weed Control in Warm-season Grasses

Apply **Detail** at 1.0 to 2.0 fl ozs/A as a broadcast postemergence spray to control emerged broadleaf weeds in season (i.e. actively growing warm-season forage grasses). In-season postemergence applications can be made in the spring after greenup. Make in-season applications before weeds reach the maximum size listed in **Table 1** on the product container label. Postemergence applications require the addition of an adjuvant system.

DO NOT apply **Detail** in-season postemergence on Bahiagrass or buffalograss.

DO NOT exceed 1.0 fl oz/A of **Detail** in forage Bermudagrass and switchgrass applied in-season postemergence (i.e. after greenup) because higher rates may cause unacceptable grass injury.

Specific Adjuvant Requirements for Postemergence Applications in Warm-season Grasses. For optimum postemergence control of emerged broadleaf weeds, the following adjuvant is required for use with **Detail**:

- MSO at 1% v/v
- **DO NOT** add nitrogen-containing fertilizers when applying **Detail** to warm-season grasses.

Sequential Applications in Warm-season and Cool-season Grasses

Detail may be applied as a sequential or split program where application(s) is made in the dormant season and subsequent application(s) is made postemergence in-season after greenup. **DO NOT** apply more than a maximum cumulative amount of 6.0 fl ozs/A of **Detail** per cropping season.

In-season postemergence applications of **Detail** may also be applied sequentially; separate sequential applications by at least 14 days. The maximum cumulative amount for in-season postemergence applications must not exceed 2.0 fl ozs/A of **Detail**.

Tank Mixtures

Broad-spectrum control of grasses and/or additional broadleaf weeds will require a tank mix with another herbicide. Read and follow the applicable restrictions and limitations and directions for use on the other product label. The most restrictive labeling applies to tank mixes. **Detail** may be tank mixed or applied sequentially with other herbicide products.

Leafy Spurge Control

Detail applied in tank mixture with **Plateau®** herbicide will control leafy spurge when applied late spring/early summer. This tank mixture will also control additional weeds listed on the respective **Detail** and **Plateau** labels.

Apply **Detail** at 1.0 to 2.0 fl ozs/A plus **Plateau** at 4.0 to 6.0 fl ozs/A to leafy spurge when it reaches the yellow bract (pre-bloom) stage in late spring/early summer. **DO NOT** apply this tank mix as a fall application because resultant control may not be satisfactory.

Spray Additives. **Detail™** herbicide plus **Plateau®** herbicide tank mix requires the use of an effective adjuvant system. For best results, use nonionic surfactant (NIS) at 0.25% v/v plus AMS at 8.5 to 17.0 lbs/100 gallons [1% to 2% weight/volume (w/v)]. Crop oil concentrate (COC) or MSO may also be used with this tank mixture when injury (stunting, necrosis) to grasses is acceptable.

Detail plus **Plateau** tank mix may be applied by ground or air.

Water Volume. For ground application, use 10 or more gallons of water per acre. Thorough coverage of weeds is essential, and higher spray volumes may be necessary for better performance on a heavy population of leafy spurge. For aerial application, use 5 or more gallons of water per acre.

Crop-specific Restrictions and Limitations

- **DO NOT** apply more than a maximum cumulative amount of 6.0 fl ozs/A of **Detail** per cropping season.
- For a mixed stand of cool-season and warm-season grasses, follow use directions for warm-season grasses when applying **Detail** in-season postemergence.
- **DO NOT** apply **Detail** to mixed stands of grass and forage legumes or to grass stands containing other desirable broadleaf species. **Detail** application will kill or cause severe injury to alfalfa, clover, other legumes, and most broadleaf species.
- There is no preharvest or pre-grazing interval for **Detail**-treated grass forage or hay.
- **DO NOT** apply **Detail** to stands of annual forage (e.g. forage sorghum, Sudangrass).
- Refer to **Table 4** on the product container label for crop rotation intervals.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Kixor and Plateau are registered trademarks of BASF.

*Detail and the **X** symbol are trademarks of BASF.*

© 2014 BASF Corporation
All rights reserved.

007969-00297.20140929.**NVA 2014-04-324-0242**
Based on: NVA 2013-04-324-0092

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

 **BASF**

We create chemistry