



LICENSED

PERIOD 2014-2016 LIC. NO.

9131.176

# Sultan™

miticide

Group 25 Acaricide

For use in mite control on ornamentals grown in outdoor nurseries, retail nurseries, residential and commercial landscapes, interiorscapes, greenhouses, lathouses and shadehouses, containers, and on forest and conifer nurseries and plantations

### Active Ingredient:

cyflumetofen: (2-methoxyethyl(*RS*)-2-(4-*tert*-butylphenyl)-2-cyano-3-oxo-3-( $\alpha,\alpha,\alpha$ -trifluoro-*o*-tolyl)propionate) ..... 18.7%

Other Ingredients: ..... 81.3%

Total: ..... 100.0%

Sultan™ miticide contains 1.67 pounds of cyflumetofen per gallon.

EPA Reg. No. 7969-337 EPA Est. No. 5905-GA-001

Product of Japan and China

**KEEP OUT OF REACH OF CHILDREN  
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

**Net Contents: 16 fl. oz.**

81085120 NVA 2014-05-391-0168

**In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).**

Produced for: BASF Corporation  
26 Davis Drive, Research Triangle Park, NC 27709



## FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information at 1-800-832-HELP (4357).

---

## Precautionary Statements

---

### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist.

### Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to **Category A** on an EPA chemical-resistance category selection chart.

#### **Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material (such as nitrile, butyl, neoprene, and/or barrier laminate)
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## Environmental Hazards

**DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

---

## Directions For Use

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, made of any waterproof material (such as nitrile, butyl, neoprene, and/or barrier laminate)
- Shoes plus socks

## NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard of agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses.

**DO NOT enter or allow others to enter treated areas until sprays have dried.**

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

### Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed.

### Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

*(continued)*

## STORAGE AND DISPOSAL *(continued)*

### Container Handling

**Nonrefillable Container. DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Triple rinse containers small enough to shake (capacity  $\leq$  5 gallons) as follows:**

Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## **In Case of Emergency**

In case of large-scale spill of this product, call:

- CHEMTREC            1-800-424-9300
- BASF Corporation    1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation    1-800-832-HELP (4357)

### **Steps to take if this material is released or spilled:**

- In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

---

## Product Information

---

**Sultan™ miticide**, a suspension concentrate (SC), is a selective contact miticide for control of a variety of mite pests on ornamental plants. When used as directed, **Sultan** is effective, providing knockdown and residual control of tetranychid mites on ornamental plants listed on this label. **Sultan** is not effective against broad mite, bulb mite, cyclamen mite, flat mite, or rust mite. **Sultan** is active on egg and motile stages. Common mite species controlled by **Sultan** include:

### Pests Controlled

Common Name	Scientific Name
Banks grass mite	<i>Oligonychus pratensis</i>
Brown almond mite	<i>Bryobia rubrioculus</i>
Brown wheat mite	<i>Petrobia latens</i>
Carmine spider mite	<i>Tetranychus cinnabarinus</i>
Citrus red mite	<i>Panonychus citri</i>
European red mite	<i>Panonychus ulmi</i>

(continued)

## Pests Controlled *(continued)*

Common Name	Scientific Name
McDaniel spider mite	<i>Tetranychus mcdanieli</i>
Pacific spider mite	<i>Tetranychus pacificus</i>
Spruce spider mite	<i>Oligonychus ununguis</i>
Strawberry spider mite	<i>Tetranychus turkestanii</i>
Texas citrus mite	<i>Eutetranychus banksi</i>
Two-spotted spider mite	<i>Tetranychus urticae</i>
Willamette spider mite	<i>Eotetranychus willamettei</i>
Yuma spider mite	<i>Eotetranychus yumensis</i>

**Sultan™ miticide** can be used on ornamental plants, including annual and perennial herbaceous plants such as foliage plants and woody trees and shrubs at the following use sites:

- Commercial and retail nurseries, field or container
- Commercial, public, and recreational gardens
- Forest tree and conifer nurseries or plantations
- Greenhouses, shadehouses, and lathhouses

- Interiorscapes
- Ornamentals in landscape
- Other commercial and residential landscape maintenance areas

**Sultan™ miticide** is compatible with most biological control organisms used for mite control.

**NOTE:** Because of its selective activity on the *Tetranychid* family of mites, **Sultan** will not have activity on other species such as broad mite, bulb mite, cyclamen mite, flat mite, or rust mite. With selective activity against tetranychid mites, **Sultan** is also compatible with the use of beneficial/predacious insects or mites (including those listed following) within integrated pest management (IPM) programs.

### **Beneficial Insects**

<b>Common Name</b>	<b>Scientific Name</b>
Common lacewing	<i>Chrysopa carnea</i>
Insidious flower bug	<i>Orius insidiosus</i>
Predatory mite	<i>Amblyseius fallacies</i>
	<i>Phytoseiulus persimilis</i>
	<i>Typhlodromus pyri</i>
	<i>Zezellia mali</i>

(continued)

## Beneficial Insects *(continued)*

Common Name	Scientific Name
Seven-spotted lady beetle	<i>Coccinella septempunctata</i>
Six-spotted thrips	<i>Scolothrips sexmaculatus</i>
Spider mite destroyer	<i>Stethorus punctum</i>
Western predatory mite	<i>Galendromus occidentalis</i>

## Resistance Management

The active ingredient in **Sultan™ miticide** is cyflumetofen, a member of the beta-ketonitrile class of chemistry. **Sultan's** mode of action is respiration inhibition classified by the Insecticide Resistance Action Committee (IRAC) as **Mitochondria complex II electron transport inhibitor** under **Group 25**. Unwarranted multiple use of a miticide with a given mode of action can result in the development of resistance.

**DO NOT** exceed the maximum seasonal use rate or the total number of applications of **Sultan** per season. To reduce the potential for developing mite resistance, rotate to a miticide with a mode of action outside IRAC **Group 25**.

Monitor treated mite populations to determine the need for reapplication, product rotation, or additional management practices. Read and follow product label before applying any miticide. Contact your local extension specialist, certified crop advisor, and/or BASF representative for additional IPM recommendations and specific steps for avoiding resistance. More information on managing mite resistance can be found on the IRAC website at <http://IRAC-online.org>.

---

## Use and Application Instructions

---

For **Sultan™ miticide** to be most effective, apply it at the first sign of mites and before the build-up of heavy pressure. Application must be timed to coincide with locally recommended treatment thresholds in developing mite populations.

**Sultan** is not systemic or translaminar. Thorough and uniform coverage of the leaf surface and/or direct contact of the spray mixture with the target pest is required for satisfactory control. Foliar application may be made using properly calibrated ground sprayers. Apply **Sultan** in a minimum of 100 gallons/acre (or 100 gallons spray solution applied per acre). Because **Sultan** is **NOT** a systemic miticide, complete coverage is necessary for effective control. Applying **Sultan** at spray volumes lower than directed can make it harder to obtain thorough coverage and may reduce performance. Consult with your local or state extension personnel for advice on miticide use and selection.

**DO NOT** apply where thorough coverage of plants is not possible. Application made with less than thorough coverage may result in slower activity and/or inadequate control.

To provide optimum coverage of ornamentals with hard-to-wet foliage, an adjuvant can be used in the **Sultan™ miticide** spray solution. Test the safety and compatibility of all adjuvants before use. Always read and follow the specific adjuvant label using the proper concentration of adjuvant to avoid plant injury.

**DO NOT** apply **Sultan** if rain or irrigation event is expected within 1 hour following application. The product can be expected to be rainfast 1 hour after application.

## Sultan™ miticide Use-specific Application Instructions

Use Sites	Application Rate per 100 gallons	Application Method
<b>Outdoor Uses</b>		
<p><b>Nurseries</b> (container, bench, flat, plug, bed-grown, field-grown, shadehouses, screenhouses, lathouses)</p> <p><b>Forest and conifer plantations</b></p> <p><b>Residential and commercial landscape</b></p> <p><b>Recreational areas and gardens with ornamental landscapes</b></p>	<p>13.7 fl ozs (0.18 pound active ingredient)</p>	<ul style="list-style-type: none"> <li>• When mites appear, apply in sufficient volume of water to ensure thorough coverage of foliage. Apply in a minimum of 100 gallons of finished spray solution per acre by ground equipment including tractor ground boom, backpack, or hand wand. Applying <b>Sultan</b> at spray volumes lower than directed can make it harder to obtain thorough coverage and may reduce performance.</li> <li>• <b>DO NOT</b> exceed the maximum seasonal use rate of 27.4 fluid ounces of <b>Sultan</b> per crop growing cycle.</li> <li>• <b>DO NOT</b> make more than two (2) applications of <b>Sultan</b> per crop per year.</li> </ul>

## Sultan™ miticide Use-specific Application Instructions *(continued)*

Use Sites	Application Rate per 100 gallons	Application Method
<b>Indoor Uses</b>		
<b>Greenhouses</b> (container, bench, flat, plug)  <b>Interiorscapes</b>	13.7 fl ozs (0.18 pound active ingredient)	<ul style="list-style-type: none"><li>• When mites appear, apply in sufficient volume of water to ensure thorough coverage of foliage. Apply in a minimum of 100 gallons of finished spray solution per acre by ground equipment including backpack or hand wand. Applying <b>Sultan</b> at spray volumes lower than directed can make it harder to obtain thorough coverage and may reduce performance.</li><li>• Use higher specified dosage for increased mite pressure or increased residual control. Lower rates can be used where mite infestations are light.</li></ul>

*(continued)*

## Sultan™ miticide Use-specific Application Instructions *(continued)*

Use Sites	Application Rate per 100 gallons	Application Method
<b>Indoor Uses</b>		
		<ul style="list-style-type: none"><li>• <b>DO NOT</b> exceed the maximum seasonal use rate of 27.4 fl ozs of <b>Sultan</b> per crop growing cycle.</li><li>• <b>DO NOT</b> make more than two (2) applications of <b>Sultan</b> per crop growing cycle.</li></ul>

---

## Restrictions and Limitations

---

- **DO NOT** exceed the maximum seasonal use rate of 27.4 fluid ounces (0.36 lb ai/A) of **Sultan™ miticide** per crop per year.
- **DO NOT MAKE MORE THAN 1 APPLICATION BEFORE ROTATING TO A PRODUCT OF A DIFFERENT MODE-OF-ACTION GROUP.**
- **DO NOT** make more than two (2) applications of **Sultan** per crop per year.
- **DO NOT** make Sultan applications at intervals shorter than 14 days.
- **DO NOT** tank mix **Sultan** with:
  - Insect growth regulators
  - Plant growth regulators
  - Carbamate insecticides
  - Organophosphate insecticides
  - Pyrethroid insecticides
- For use in ground application equipment only.
- **DO NOT** apply by chemigation or aerial application.
- Not for use on plants intended for feed or forage.
- **DO NOT** use on vegetable transplants.
- **DO NOT** apply to plants exhibiting symptoms of stress or injury, such as stunting, wilting, leaf burn or drop, or abnormal growth.

---

## Use Precautions

---

**Sultan™ miticide** has been applied to a wide variety of common ornamental plants without observed plant injury. Not all species, varieties, and cultivars have been tested for tolerance to **Sultan**, possible tank mix combinations with **Sultan**, pesticide treatments preceding or following those with **Sultan**, and combinations of **Sultan** with surfactants or adjuvants. Local conditions can also influence plant tolerance and may not match those under which BASF has conducted testing. Because many cultivars within a plant species vary in tolerance to chemical applications and growing conditions, the grower must recognize these differences and test the product accordingly. At a minimum, always test a small group of representative plants for tolerance to **Sultan** under local growing conditions and before large-scale use.

Grower assumes responsibility for testing species suitability under local growing conditions by treating a small number of plants at the specified rate. At a minimum, this should include evaluating treated plants for several weeks following treatment for possible injury or other effects. To the extent consistent with applicable law, by applying **Sultan**, the user assumes responsibility for any crop damage or other liability associated with factors beyond the manufacturer's control, such as weather, presence of other materials, and manner or use of application.

---

## Spray Drift Management

---

**Water Volume** - Apply in at least 100 gallons/A spray solution for optimum performance. Because **Sultan™ miticide** is not systemic or translaminar, thorough coverage of plant surfaces is necessary for effective control. Applying **Sultan** at spray volumes lower than directed can make it harder to obtain thorough crop coverage and may reduce performance.

**Fixed Boom Nozzle Height** - Maximum nozzle height (release height) should be no greater than 2 feet above the plant canopy.

**Variable Winds** - **DO NOT** apply in variable wind conditions.

**Wind Direction** - Base wind direction (e.g. away from nontarget areas) on the average direction (not instantaneous).

**Wind Speed** - Maximum sustained wind must not exceed 15 mph measured at nozzle height (release height).

**Inversion Restriction** - **DO NOT** apply if conditions will capture and potentially transport spray droplets off target (consult Pasquill stability categories).

**Terrain Precaution** - Local terrain can influence wind patterns. The applicator must be familiar with these patterns and understand their impact on drift.

**Sensitive Areas** - To ensure the protection of threatened or endangered species, it is important to maintain spray drift loadings below levels of concern for any area adjacent to the application site that is not excluded as possible habitat for these organisms. The following restrictions apply:

- **Buffer distance** - 15 feet
- **Release height** - 20 inches
- **Droplet size** - A combination of spray nozzles and appropriate pressure must be selected to provide ASABE standard S571.1 droplet size category of fine ( $DV_{0.5}$  of  $\geq 180$  microns) or coarser than fine.
- **Maximum wind speed** - 15 mph

---

## Additives and Tank Mixing Information

---

Shake container well before use.

**Sultan™ miticide** can be tank mixed with most recommended fungicides, insecticides, liquid fertilizers, biological control products, adjuvants, and additives **except as noted in the Restrictions and Limitations section of this label.**

Under some conditions, the use of additives or adjuvants may improve the performance of **Sultan™ miticide**. Local conditions can influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced mite control, or crop injury may result from mixing **Sultan** with other products.

Before using any tank mix (fungicides, insecticides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Treat a small group of plants including all cultivars, and evaluate for 3 to 7 days before applying to the entire crop.

When an adjuvant is to be used with this product, BASF recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

---

## Tank Mixing Order

---

1. **Water** - Begin by agitating a thoroughly clean sprayer tank 3/4 full of clean water.
2. **Agitation** - Maintain constant agitation throughout mixing and application.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-dispersible products** (such as **Sultan™ miticide**, dry flowables, wettable powders, other suspension concentrates, or suspo-emulsions)
6. **Water-soluble products**
7. **Emulsifiable concentrates** (such as oil concentrates when applicable)
8. **Water-soluble additives**
9. **Remaining quantity of water**

Thoroughly mix and suspend each component before adding tank mix partners.

Maintain constant agitation during application.

---

## Cleaning Application Equipment

---

Before application, start with clean, well-maintained application equipment. Following spray application, thoroughly clean all application equipment. Drain application equipment of any excess product. Thoroughly rinse application equipment and flush hoses, boom, and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean equipment near wells, water sources, or desirable vegetation. Dispose of waste rinsewater in accordance with local regulations.

Spraying equipment must be cleaned thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used before **Sultan™ miticide**.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

*(continued)*

## **Conditions of Sale and Warranty** *(continued)*

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF. 1108

**Sultan** is a trademark of BASF.

© 2014 BASF Corporation  
All rights reserved.

007969-00337.20111205g.**NVA 2011-04-391-0232**

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



The Chemical Company

## NOTES

## NOTES

## NOTES

## NOTES

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that proper record-keeping is essential for transparency and accountability, particularly in the context of public administration and government operations. This section also highlights the role of technology in streamlining record management processes and reducing the risk of errors or data loss.

2. The second part of the document focuses on the implementation of robust internal controls and risk management frameworks. It outlines the need for regular audits and assessments to identify potential vulnerabilities and ensure that organizational policies are effectively enforced. This section also discusses the importance of employee training and awareness programs to foster a culture of compliance and ethical behavior within the organization.

3. The third part of the document addresses the challenges of data security and privacy protection in the digital age. It highlights the increasing threat of cyberattacks and the need for comprehensive security protocols, including encryption, access controls, and regular security updates. This section also discusses the importance of data backup and recovery strategies to ensure business continuity in the event of a data breach or system outage.

4. The fourth part of the document explores the role of leadership and governance in driving organizational success. It emphasizes the importance of clear communication, strategic vision, and effective decision-making. This section also discusses the need for strong ethical leadership and the establishment of a code of conduct that guides the behavior of all employees and stakeholders.

5. The fifth and final part of the document provides a summary of the key findings and recommendations. It reiterates the importance of a holistic approach to organizational management, one that integrates financial, operational, and human resources. The document concludes by encouraging organizations to embrace change and innovation as a means of staying competitive in a rapidly evolving market.

# Sultan™

Group 25 Acaricide

miticide

For use in mite control on ornamentals grown in outdoor nurseries, retail nurseries, residential and commercial landscapes, interiorscapes, greenhouses, lathhouses and shadehouses, containers, and on forest and conifer nurseries and plantations

**Active Ingredient:**

cyflumetofen: (2-methoxyethyl(*RS*)-2-(4-*tert*-butylphenyl)-2-cyano-3-oxo-3-( $\alpha,\alpha,\alpha$ -trifluoro-*o*-tolyl)propionate) ..... 18.7%

**Other Ingredients:** ..... 81.3%

**Total:** ..... 100.0%

**Sultan™ miticide** contains 1.67 pounds of cyflumetofen per gallon.

EPA Reg. No. 7969-337 EPA Est. No. 5905-GA-001

Product of Japan and China

**KEEP OUT OF REACH OF CHILDREN  
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail.)

See attached booklet for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

**Net Contents: 16 fl. oz.**

81085120 NVA 2014-05-391-0168

**In case of an emergency endangering life or property involving**

**this product, call day or night 1-800-832-HELP (4357).**

Produced for: BASF Corporation  
26 Davis Drive, Research Triangle Park, NC 27709

 **BASF**  
The Chemical Company