

Reg B.



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO.

9138.25

Disc.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** For control of Norway rats, roof rats, and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Wear protective gloves when mixing and applying bait. With detergent and hot water, wash all implements used for mixing and applying bait. Do not use these implements for mixing, holding, or transferring foods or feeds. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not apply bait directly to ground surface or in grass or other ground cover.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

**MIXING DIRECTIONS**

Thoroughly mix one part of concentrate with 19 parts of suitable bait materials (cereal, grain, fish, meat, or vegetables, etc.), including binding agents and sweeteners if needed. For example, mix eleven parts of rolled oat groats, six parts of whole yellow ground cornmeal, one part sugar, and one part corn oil with one part of Diphacinone Concentrate. Blend for 20 minutes in a rotary blender.

# DIPHACINONE CONCENTRATE

## A CONCENTRATE FOR PREPARING TO KILL NORWAY RATS, ROOF RATS, AND HOUSE MICE.

**ACTIVE INGREDIENT:**

Diphacinone	
2-(Diphenylacetyl)-1,3-indandione .....	0.1%
INERT INGREDIENTS .....	99.9%
<b>TOTAL</b>	<b>100.0%</b>

### KEEP OUT OF REACH OF CHILDREN CAUTION

(See right panel for additional  
precautionary statements.)


EPA REG. NO. 12455-9  
EPA EST. NO. 12455-WI-1

**NET CONTENTS: 50 LBS. (22.7 kg)**

Manufactured by:



Bell Laboratories, Inc.  
Madison, WI 53704 U.S.A.



**APPLICATION DIRECTIONS**

**RATS:** Using a measuring scoop or other suitable implement, apply 4-16 oz. of bait per placement. Space placements at intervals of 15 to 30 feet. Maintain an uninterrupted supply of fresh bait for a least 10 days or until signs of rat activity cease.

**MICE:** Apply 1/4 to 1/2 oz. (1 - 2 level tablespoons) of bait at each placement location. Space placements at 8- to 12- foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

**RATS AND MICE:** Replace contaminated or spoiled bait immediately.

If perishable baits (meat, fish, or vegetables) are used, place fresh baits in late afternoon. Replace daily with fresh baits.

Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store only in original container, in a dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag and carton. Then dispose of empty bag and carton in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**PRECAUTIONARY STATEMENTS**

**HAZARD TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

Keep away from humans, domestic animals, and pets. Exposure during pregnancy should be avoided. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. When mixing baits, wear dust respirator and rubber gloves. If on skin, wash with soap and water.

**NOTE TO PHYSICIAN:** If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

**ENVIRONMENTAL HAZARD**

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.