



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

9165.510

DISC.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2005-2007** LIC. NO.

ORTHO[®]

BUGB-GON[®]

MAX[®]

GARDEN INSECT DUST



BEETLE



APHID



CATERPILLAR

**Kills
60+
Insects**

**New Easy
To Use
Applicator!**



**Use on Ornamentals,
Fruits & Vegetables**

**KEEP OUT OF REACH OF CHILDREN
CAUTION** SEE BACK PANEL BOOKLET FOR ADDI-
TIONAL PRECAUTIONARY STATEMENTS

Active Ingredient

Permethrin*.....0.25%
Other Ingredients...99.75%

*cis/trans ratio:

min. 35%(+) cis,
max. 65%(+) trans

NET WT 1 LB/453g



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2008-2010** LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2011-2013** LIC. NO.

Back Cover (Last Page - Pg 6)

2"

Front Cover (Page 1)

2-1/8"

FIRST AID



IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

PHYSICAL AND CHEMICAL HAZARDS: Do not use or store near heat or open flame.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

OPEN

Resealable Label
For Directions &
Precautions

BUGGON®

ORTHO®

MAX

GARDEN INSECT DUST

PRODUCT FACTS

WHERE TO USE	On ornamentals including roses, vegetables, melons, fruit and nut trees.
INSECTS CONTROLLED	Aphids, beetles, caterpillars, cutworms, leafhoppers, and other garden insects.
WHEN TO USE	Use at first signs of insect infestation, preferably before damage from feeding occurs.



Questions, Comments or Medical Information?

Call 1-800-225-2883



www.ortho.com

Manufactured for
The ORTHO Group
P.O. BOX 190
Marysville, OH 43040
Form LBXXXXXXX

EPA Reg. No. 28293-126-239
EPA Est. 58996-MO-1

Made in USA

80% SIZE
12-digit UPC
(non-suppressed)
For Position Only

1005

0 71549 01735 ?



FPO

ITF CODE IS
??????

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE**HOME GARDENS**

For use on home gardens. Not for use on plants being grown for SALE or other COMMERCIAL USE.

HOW TO APPLY

1. Apply when air is calm to avoid drift and contact with eyes and skin.
2. Wearing household latex or rubber gloves, unscrew cap and remove seal from bottle top.
3. Tightly screw cap back on bottle and flip top open.
4. From a distance of 12 inches point bottle towards plant to be treated.
5. Point bottle downward in a 45° angle, and with one hand, apply light constant pressure to the center of the bottle while shaking sideways to produce a light dusting.
6. Dust lightly to cover upper and lower leaf surfaces with a thin, even film of dust. Many insects hide on the underside of the leaf.
7. To continue applying, release pressure from bottle before squeezing again.

WHEN TO APPLY

Use at first signs of insect infestation, preferably before damage from feeding occurs.

Note the pre-harvest (time between use and harvest) interval with each crop listed.

VEGETABLES & MELONS	APPLICATIONS PER SEASON (MAX)	DAYS TO WAIT TO HARVEST
---------------------	-------------------------------	-------------------------

Broccoli, brussels sprouts & cauliflower	8	1
Cabbage & Chinese cabbage (tight-heading varieties only)	5	1
Cucumbers, gourds, melons & squash	8	0
Eggplant	10	3
Peppers (bell)	8	3
Tomatoes	6	0
Amaranth, arugula, celery, lettuce (head & leaf), parsley, radicchio, rhubarb, spinach & Swiss chard	5	1
Onions (dry bulb & garlic)	6	1
Potatoes	8	14
Sweet corn	6 (FL only: 8)	1
Artichokes	5	0
Asparagus	4	1

FRUIT & NUT TREES	APPLICATIONS PER SEASON (MAX)	DAYS TO WAIT TO HARVEST/ COMMENTS
-------------------	-------------------------------	-----------------------------------

Apples		Do not apply after petal fall
Cherries (in the East)	6	No more than 4 applications after petal fall
Cherries (in the West)	4	No more than 3 applications after petal fall
Peaches	5	14
Pears (Dormant)	2	May be applied during the dormant through the delayed dormant growth periods only
Pears (Pre-bloom)	2	May be applied from dormant through bud burst
Walnuts	8	1

INSECTS CONTROLLED

On Fruit & Nut Trees

Apple aphid, caterpillars, codling moth, filbertworm, fruit fly, green fruitworm, lepidopterous larvae, leafhoppers, leafrollers, lesser peachtree borer, mirid bug, navel orangeworm, oriental fruit moth, peach twig borer, pear psylla, plum curculio, redbanded leafroller, rose chafer, spotted tentiform leafminer, tarnished plant bug, thrips, white apple leafhopper and walnut husk fly

On Vegetables & Melons

Aphids, armyworms, artichoke plume moth, asparagus beetle, Aster leafhopper, cabbage aphid, cabbage looper, Colorado potato beetle, corn earworm, corn rootworm (adult), cutworm complex, cucumber beetle (adult), cutworms, diamondback moth, European corn borer, fall armyworm, flea beetles, granulate cutworm, hornworms, imported cabbageworm, leafhoppers, leafminers, melon worm, onion maggot (adult), onion thrips, pepper weevil, pickle worm, plant bugs (including lygus, squash and stink bugs), potato psyllid, potato aphid, potato flea beetle, potato leafhopper, potato tuberworm, rindworm, southern armyworm, squash vine borer, tomato fruitworm and tomato pinworm, vegetable leafminer

On Ornamentals (Including Roses)

Asparagus beetle, beet armyworm, cabbage looper, Japanese beetle, lygus bug and omnivorous leafroller

STORAGE AND DISPOSAL



STORAGE: Store in a cool dry place. Do not contaminate feed or foodstuffs by storage or disposal.

DISPOSAL: IF EMPTY: Do not reuse this container. Place in trash or offer for recycling if available.

IF PARTLY FILLED: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS



HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Humans: Harmful if swallowed, inhaled. Avoid breathing dust. Avoid contact with eyes and skin. Wash thoroughly with soap and water after handling. Avoid contamination of feed and foodstuffs. **Animals:** Harmful if swallowed, inhaled.

Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using **ANY** pesticide product for pets.

If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

THE ORTHO GUARANTEE

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.



Questions, Comments or Medical Information?

Call 1-800-225-2883



www.ortho.com

© 2005, Licensed by The ORTHO Group.
World rights reserved.

Manufactured for **The ORTHO Group**
P.O. Box 190, Marysville, OH 43040

Form LBXXXXXXXX

EPA Reg No 28293-126-239

EPA Est. 58996-MO-1

Made in USA