



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

9200.121

DuPont™ Avaunt®
INSECTICIDE



FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Remove contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing (dust, vapor or spray mist). Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category A on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all ≥14 mils.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS. **IMPORTANT:** when reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicator and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

**Sold by: E.I. du Pont de Nemours and Company,
Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.
Made in U.S.A.**





DuPont™ Avaunt®

insecticide

GROUP	22	INSECTICIDE
-------	----	-------------

Dispersible Granules

Active Ingredient	By Weight
--------------------------	------------------

Indoxacarb

(S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxy-carbonyl)[4(trifluoromethoxy)phenyl]amino]-carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a-(3H)-carboxylate	30%
--	-----

Other Ingredients	70%
--------------------------------	-----

TOTAL	100%
-------	------

EPA Reg. No. 352-597

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

(continued)

FIRST AID (cont'd)

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Remove contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-441-3637** for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing (dust, vapor or spray mist). Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category A on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

Long-sleeved shirt and long pants.

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all ≥ 14 mils.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS. **IMPORTANT:** when reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicator and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

PRECAUTIONARY STATEMENTS (cont'd)

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing and/or PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, birds, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Cover, incorporate, or clean up granules that are spilled.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

DuPont™ AVAUNT® must be used only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and

(continued)

AGRICULTURAL USE REQUIREMENTS (cont'd)

restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all crops.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

Coveralls over long-sleeved shirt and long pants,

Socks plus chemical resistant footwear,

Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all ≥ 14 mils.

DuPont™ AVAUNT® must be used only in accordance with the directions on this label or in separate DuPont supplemental labeling available as a result of new EPA approvals.

AVAUNT® is a water dispersible granule that can be applied as a foliar spray to control many important insects. AVAUNT® is mixed with water for application.

Use only in commercial and farm plantings. Not for use in home plantings.

Do not formulate this product into any other End-use products without written permission of DuPont.

CHEMIGATION: Do not apply this product through any type of irrigation system except for application to cranberries, mint, potatoes and sweet corn and as allowed by Federal Supplemental and Special Local Need (SLN) labeling. (See "Application By Chemigation" section of the label.)

INTEGRATED PEST MANAGEMENT

DuPont supports the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an IPM program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, rotation of insecticides with different modes-of-action, and treating when target pest populations reach locally determined

action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

SCOUTING

Monitor insect populations to determine whether or not there is a need for application of DuPont™ AVAUNT® based on locally determined economic thresholds. More than one treatment of AVAUNT® may be required to control a population of pests.

RESISTANCE MANAGEMENT

For resistance management, AVAUNT® is a group 22 insecticide. Repeated and exclusive use of AVAUNT® or other group 22 insecticides may lead to the buildup of resistant strains of insects in some crops.

Some insects are known to develop resistance to products used repeatedly for control. When this occurs, the recommended dosages fail to suppress the pest population below the economic threshold. Because the development of resistance cannot be predicted, the use of this product should conform to resistant management strategies established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of mode-of-action classes of insecticides on succeeding generations and targeting the most susceptible life stage. Consult your local or state agricultural authorities for details.

Unless directed otherwise in the specific crop/pest sections of this label, the following best practices are recommended to delay the development of insecticides resistance:

- Make no more than 2 successive applications per generation or within a 30 day period to the same insect species on a crop.
- The following application to the target pest(s) must be with an effective product with a different mode of action (for example ASANA® XL or DuPont™ LANNATE® insecticides).

If resistance to AVAUNT® develops in your area, AVAUNT®, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company

representative or agricultural advisor for the best alternate method of control for your area. For additional information on insect resistance monitoring, visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irc-online.org>.

APPLICATION

Apply at the listed rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Follow-up treatments of AVAUNT® should be applied, as needed, to keep pest populations within threshold limits. Apply AVAUNT® on most crops every 3 to 5 days, as specified in the specific crop sections, to maintain control. For bushberry, cranberry, dry bean, pome and stone fruit the minimum interval between treatments is 7 days.

Use sufficient water to obtain thorough, uniform coverage.

Because AVAUNT® is most effective through ingestion of treated plant material, thorough spray coverage is essential for optimum control of targeted pest insects. Using increased water volumes will typically result in better spray coverage, especially under adverse conditions such as dry, hot weather or dense plant foliage. AVAUNT® may be applied by ground, aerial or overhead sprinkler chemigation application equipment. For aerial application use the following directions unless otherwise specified in this label: use a minimum of 5 gallons per acre (gpa) of water, except in tree and vine crops use a minimum of 10 gpa. For ground applications, use the following directions unless otherwise specified in this label: use a minimum of 10 gallons per acre (gpa) of water, except in tree and vine crops use a minimum of 50 gpa and a maximum of 200 gpa of water.

Use of Adjuvants: In some situations where coverage is difficult to achieve such as closed canopy, dense foliage, plants with waxy leaf surfaces, or less than optimum application equipment, an adjuvant may improve performance. Use only adjuvant products that are labeled for agricultural use and follow the directions on the manufacturer's label. For uses in fruit crops, use a proven and recommended adjuvant that does not affect fruit finish.

Do not use adjuvant on bushberries or garden beets.

SPRAY PREPARATION

Spray equipment must be clean and free of previous pesticide deposits before applying AVAUNT®. Fill spray tank 1/4 to 1/2 full of water. Add AVAUNT® directly to spray tank. Mix thoroughly to fully disperse the insecticide; once dispersed continued agitation is required. Use mechanical or hydraulic means; do not use air agitation. Spray mix must not be stored overnight in spray tank.

Compatibility - Since formulations may be changed and new ones introduced, it is recommended that users premix a small quantity of a desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.). Avoid mixtures of several materials and very concentrated spray mixtures.

This product can be mixed with pesticide products labeled for use on crops on this label in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

Tank Mixing Sequence - Add different formulation types in the sequence indicated below. Allow time for complete mixing and dispersion after addition of each product.

1. Water soluble bags.
2. DuPont™ AVAUNT® and other water dispersible granules.
3. Wettable powders.
4. Water based suspension concentrates.
5. Water soluble concentrates.
6. Oil based suspension concentrates.
7. Emulsifiable concentrates.
8. Adjuvants, surfactants, oils.
9. Soluble fertilizers.
10. Drift retardants.

Follow local practice and manufacturer's recommendations.

SPRAY TANK CLEANOUT

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water.

Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

CROP ROTATION

Crops that are on this label and alfalfa, cotton, peanuts and soybeans may be planted immediately following harvest. Do not plant for food or feed any other crops not registered for use with indoxacarb for 30 days after last use.

APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH* AND SWEET CORN

*Use on spinach via overhead sprinkler irrigation is allowed only in the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling.

Instructions for the Use of AVAUNT® in Overhead Sprinkler Chemigation Systems.

Overhead chemigation applications offer the advantage of greater penetration and coverage of the target plant. However, typical chemigation applications are more dilute than ground or aerial applications. For best results, it is recommended to keep the concentration of AVAUNT® as high as possible in the application. Apply AVAUNT® in 0.1 to 0.2 inches of water per acre.

AVAUNT® is most active as an ingestion insecticide, although it does have activity as a direct contact insecticide. For best results, applications of AVAUNT® should ensure thorough coverage of the target plant to maximize the opportunity for target insects to ingest AVAUNT®.

Types of Chemigation Systems:

AVAUNT® may be applied only through overhead sprinkler irrigation systems. Overhead irrigation systems include the following; center pivot, end tow, hand move, lateral move, side roll, solid set and wheel line.

Center pivot and lateral move irrigation systems are preferred. Other overhead sprinkler systems may be used if they provide uniform water distribution. Do not apply DuPont™ AVAUNT® through any other type of irrigation system. Do not use filter screens smaller than 50 mesh throughout the system, due to possible build up of material on 100 mesh or smaller screens.

Directions for Chemigation:

Preparation

Use a pesticide tank for the application of AVAUNT® in chemigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. With the mix tank 1/4 to 1/2 full with water and the agitator running, measure the required amount of AVAUNT® and add it to the tank. Then add additional water to bring your total pesticide mixture up to the desired volume for your application. Note: Always add the AVAUNT® to water, never put AVAUNT® into a dry tank or other mixing equipment without first adding water. See container label for tank mixing sequence. Continue to agitate the mixture throughout the application process. Use mechanical or hydraulic agitation, do not use air agitation. Highly alkaline water must be buffered so that the pH of the spray solution is in the range of neutral to slightly acidic.

Injection Into Chemigation Systems

Inject the proper amount of AVAUNT® into the irrigation water flow using a positive displacement injection pump. Inject the mixture at a point in the main irrigation water flow to ensure thorough mixing with the irrigation water. For continuously moving systems, inject the solution containing AVAUNT® into the irrigation water line continually and uniformly throughout the irrigation cycle. Apply in no more than 0.2 inches of water per acre. For overhead sprinkler systems that are stationary, add the solution containing AVAUNT® to the irrigation water line and apply no more than 0.2 inches of water per acre just before the end of the irrigation cycle.

Uniform Water Distribution

The irrigation system used for application of AVAUNT® must provide for uniform distribution of AVAUNT® treated water. Non-uniform distribution might result in crop injury, lack of effectiveness or illegal pesticide residues in or on the crop being treated. Ensure the irrigation system is calibrated to uniformly distribute the chemigation application to the crop. Contact the equipment manufacturer, the local University Extension agent or other experts if you have questions about achieving uniform distribution of the application.

Equipment Calibration

Calibrate the irrigation system and injector before applying AVAUNT®. Calibrate the injection pump while the system is running using the expected irrigation rate. If you have questions about calibration, you should contact your state extension service specialists, equipment manufacturer or other experts.

Monitoring of Chemigation Applications

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of a responsible person, shall shut the system down and make necessary adjustments should the need arise. Wear the personal protective equipment as defined in the PPE section of the label for applicators and other handlers when making adjustments or repairs on the chemigation system when AVAUNT® is in the irrigation water.

Required System Safety Devices

Do not connect any irrigation system used for pesticide applications to a public water system unless the pesticide label-prescribed safety devices are in place. Public water system means a system for the provision to the public of piped water for human consumption, if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals at least 60 days out of the year.

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Operation

Start the water pump and sprinkler, and let the system achieve the desired pressure and speed before starting the injector. Start the injector and calibrate the injection system according to the directions above. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

- End guns must be turned off during the application, if they irrigate nontarget areas or if they do not provide uniform application and coverage.
- Plug nozzles in the immediate area of control panels, chemical supply tanks and system safety devices to prevent contamination of these areas.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Do not apply when system connections or fittings leak or when nozzles do not provide uniform distribution.
- Do not allow irrigation water to collect or run-off during chemigation.

Cleaning the System

Thoroughly clean the injection system and tank of any fertilizer or chemical residues using a standard clean-out procedure. Dispose of any residues in accordance with State and Federal laws. Consult your owner's manual or your local equipment dealer for cleanout procedures for your injection system.

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Bean, Dried (except soybean) Including: Dried cultivars of bean (Lupinus) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (Phaseolus) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (Vigna) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil	Corn Earworm European Corn Borer	0.065 - 0.11	3.5 - 6.0	7	12 hrs
	Make no more than 4 applications per acre per crop. Do not apply more than 24 oz AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 72 oz of AVAUNT® or 1.32 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 7 days. For ground applications, make a uniform application in approximately 20-100 gallons of water per acre.				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Bean, Succulent (except soybean) Including: Bean (Phaseolus) includes (lima bean, green; broad bean, succulent; runner bean, snap bean, wax bean); bean (Vigna) (includes asparagus bean, blackeyed pea, Chinese longbean, cowpea, moth bean, southern pea, yardlong bean); jackbean; sword bean	Corn Earworm European Corn Borer	0.065 - 0.11	3.5 - 6.0	3	12 hrs
	Make no more than 4 applications per acre per crop. Do not apply more than 14 oz AVAUNT® or 0.26 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 56 oz of AVAUNT® or 1.04 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 7 days. For ground applications, make a uniform application in approximately 20-100 gallons of water per acre.				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Brassica (Cole) Leafy Vegetables	Beet Armyworm	0.065*	3.5*	3	12 hrs
	Diamondback moth				
Including: Broccoli, Chinese broccoli, Broccoli raab, Brussels sprouts, Cabbage, Chinese cabbage (napa and bok choy), Chinese mustard cabbage, Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens and Turnip tops**	Cabbage looper Cabbage webworm (except California) Cross striped cabbageworm (except California) Imported cabbageworm	0.045 - 0.065*	2.5 - 3.5*		
	<p>Make no more than 4 applications per acre per crop. Do not apply more than 14 oz AVAUNT® or 0.26 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 56 oz of AVAUNT® or 1.04 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 3 days. Do not apply to greenhouse or field grown brassica crops grown for transplant. Resistance Management for Diamondback Moth: Do not apply AVAUNT® more than twice to any generation of diamondback moth larvae. After the second application, rotate to another insecticide with a different mode of action (ie. a product with a different IRAC group number). Do not apply less than 3.5 ounces of AVAUNT® per acre. If applications of AVAUNT® do not result in reduction in diamondback moth larvae populations, immediately stop use of AVAUNT® and apply a registered insecticide with a different mode of action. Do not make more than 6 total applications of AVAUNT® per calendar year for control of diamondback moth per farm location. In the State of Georgia: Do not apply more than 4 applications of AVAUNT® per calendar year for the control of diamondback moth per farm location. *Add a wetting agent to improve spray coverage. **For use on turnips grown for tops or greens, not for the production of turnip roots.</p>				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Bushberries Including: Aronia berry, Blueberries (Highbush blueberry, and Lowbush blue- berry), Chilean guava, Currants (Black currant, Buffalo currant, Native currant and Red currant), European barberry, Elderberry, Gooseberry, Highbush cranberry, Honeysuckle, Huckleberry, Jostaberry, Juneberry, Salal, Sea buckthorn; cultivars, varieties and/or hybrids of these.	Cranberry fruitworm, Cherry fruitworm Winter moth	0.065 - 0.11	3.5 - 6.0	7	12 hrs
	Bruce spanworm Cranberry weevil (adult) Plum curculio (adult)	0.11	6.0		
<p>Make no more than 4 applications per season. Do not apply more than 24 oz of AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per year. Do not apply dilute applications of more than 200 gallons of water per acre. Do not use adjuvants. For best results apply in 20 - 50 gallons of water per acre by ground or 10 gallons of water per acre by air. Minimum interval between treatments is 7 days.</p>					

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Corn (sweet) For application through tassel push only.	European corn borer (except California) Fall armyworm Corn earworm	0.045 - 0.065	2.5 - 3.5	3 35 - fodder & stover	12 hrs
	<p>Make no more than 4 applications per acre per crop. Whorl stage through tassel push (prior to silking) application only. Do not apply more than 14 oz AVAUNT® or 0.26 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 42 oz of AVAUNT® or 0.78 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 3 days. Overhead Chemigation - AVAUNT® may be applied to sweet corn by overhead chemigation. For specific guidance see label section titled APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH* AND SWEET CORN. Begin application when sweet corn is in the V1 (1st collar) stage of growth up to tassel push (V15) when damage from larvae populations exceed recommended thresholds. For best results, a slurry of AVAUNT®, vegetable oil and an emulsifier must be kept continuously agitated in the injection tank to keep the mixture in suspension and to ensure application of the proper rate per acre.</p>				14 days for hand harvesting

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Low growing berry subgroup, (except lowbush blueberry and strawberry) Including: Bearberry; bilberry; cloudberry; cranberry; lingonberry; muntries; partridgeberry; cultivars, varieties and/or hybrids of these	Cranberry weevil* Blackheaded fireworm Black vine weevil (adult)** (OR and WA only) Spanworm	0.11	6.0	30	12 hrs
<p>Do not apply to flow through bogs or allow release of irrigation water from bogs for at least 1 day following application. Make no more than 3 applications per acre per season. Do not apply more than 24 oz of AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per season. The minimum interval between sprays is 7 days. *Apply up to two applications to the spring (overwintering) generation of adult cranberry weevil prior to bloom. Do not apply more than 12 oz. AVAUNT® (0.22 lb ai) per acre per season for control of cranberry weevils. ** Black vine weevil adults are nocturnal feeders - it is important to monitor adult emergence by regular sweeping or trapping in the evening hours. Make repeat applications on a 7 to 10 day schedule if monitoring indicates continued adult feeding activity. Broadcast applications may need supplemental spot treatments in localized areas of heavy insect pressure. Allow 5 to 7 days to achieve maximum results, then follow-up with night time monitoring.</p>					

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Cucurbit vegetables Including: Chayote (fruit), Chinese waxgourd (Chinese) preserving melon) Citron melon, Cucumber, Gherkin, Edible gourd (including hyotan, cucuzza, hechima and Chinese okra), Momordica species (including balsam apple, balsam pear, bitter melon and Chinese cucumber), Muskmelon (including true canteloupe, canteloupe, casaba, crenshaw melon, (continued)	Cabbage Looper Melonworm Pickleworm	0.045 - 0.11	2.5 - 6.0	3	12 hrs
	Beet Armyworm	0.065 - 0.11	3.5 - 6.0		
Make no more than 4 applications per acre per crop Do not apply more than 24 oz AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 72 oz of AVAUNT® or 1.32 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 5 days. For ground applications, apply using a minimum of 10 gallons per acre of water.					

(continued)

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Cucurbit vegetables <i>(continued)</i> golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon), Pumpkin, Summer squash (including crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini), Winter squash (including butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) and Watermelon	Cabbage Looper Melonworm Pickleworm	0.045 - 0.11	2.5 - 6.0	3	12 hrs
	Beet Armyworm	0.065 - 0.11	3.5 - 6.0		
<p>Make no more than 4 applications per acre per crop. Do not apply more than 24 oz AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 72 oz of AVAUNT® or 1.32 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 5 days. For ground applications, apply using a minimum of 10 gallons per acre of water.</p>					

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Fruiting Vegetables and Okra Including: eggplant, groundcherry, pepino, peppers (bell, chili, cooking, pimento and sweet), tomatillo and tomato	Beet armyworm European Corn Borer* (except California) - bell pepper only Leafminer (Use on Florida tomatoes only - suppression only)** Southern armyworm Tomato fruitworm Tomato pinworm Western yellowstriped armyworm	0.065	3.5	3	12 hrs
	Hornworms Loopers	0.045 - 0.065	2.5 - 3.5		
<p>Make no more than 4 applications per acre per crop. Do not apply more than 14 oz AVAUNT® or 0.26 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 42 oz of AVAUNT® or 0.78 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 5 days. *- European corn borer applications for use only on bell peppers - For best results, begin applications of AVAUNT® following two applications of an organo-phosphate insecticide labeled for European corn borer control in bell pepper. ** Suppression of leafminer on Florida tomatoes- Use of an adjuvant may improve performance.</p>					

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Garden beet	Beet armyworm	0.065 - 0.11	3.5 - 6.0	7	12 hrs
	<p>Make no more than 4 applications per acre per crop. Do not apply more than 24 oz AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 96 oz of AVAUNT® or 1.76 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 3 days. Do not use adjuvants.</p>				
Grape	Grape leaf folder Japanese beetle (except California) Western grape leaf skeletonizer	0.065 - 0.11	3.5 - 6.0	7	12 hrs
	European grapevine moth Grape berry moth (except California) Leafhoppers (suppression only) Light brown apple moth	0.09 - 0.11	5.0 - 6.0		

(continued)

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Grape (continued)	Katydid (nymphs)* Omnivorous leafroller	0.11	6.0	7	12 hrs
	<p>Make the first application at initiation of egg hatch or at the first signs of infestation for each generation. Use the higher application rate for moderate to heavy insect pressure. Make application before pests reach damaging levels. Monitor fields and make an additional application if populations rebuild to potentially damaging levels. Apply in sufficient water to obtain thorough coverage of foliage. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for these pests.</p> <p>Make no more than 2 applications per season. Do not apply more than 12 oz AVAUNT® or 0.22 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 21 days. For best results, use an adjuvant to help increase coverage, penetration and thus performance. * Forktailed bush katydid (<i>Scudderia furcata</i>) and Angularwinged katydid (<i>Microcentrum retinerve</i>). Correct timing of spray application is to the early nymphal stages; thorough spray coverage is critical to achieve best results. Make repeat applications on a 7 to 10 day schedule if monitoring indicates continued feeding activity.</p>				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Small fruit vine climbing subgroup, (except fuzzy kiwifruit) Including: Amur river grape; gooseberry; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these	Grape leafroller Japanese beetle (except California) Western grapeleaf skeletonizer	0.065 - 0.11	3.5 - 6.0	7	12 hrs
	Grape berry moth (except California) Leafhoppers (suppression only)	0.09 - 0.11	5.0 - 6.0		
	Omnivorous leafroller	0.11	6.0		
<p>Make the first application at initiation of egg hatch or at the first signs of infestation for each generation. Use the higher application rate for moderate to heavy insect pressure. Make application before pests reach damaging levels. Monitor fields and make an additional application if populations rebuild to potentially damaging levels. Apply in sufficient water to obtain thorough coverage of foliage. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for these pests. Make no more than 2 applications per season. Do not apply more than 12 oz AVAUNT® or 0.22 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 21 days. For best results, use an adjuvant to help increase coverage, penetration and thus performance.</p>					

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Leafy Green Vegetables, (except spinach and spinach varieties) Including: Arugula (Roquette), Chervil, Edible-leaved chrysanthemum, Garland chrysanthemum, Corn salad, Garden cress, Upland cress (yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Head and Leaf Lettuce, Orach, Parsley, Garden Purslane, Winter purslane and Radicchio (red chicory)	Beet armyworm	0.065 - 0.11	3.5 - 6.0	3	12 hrs
	Corn earworm				
	Cabbage looper	0.045 - 0.065	2.5 - 3.5		
Make no more than 4 applications per season. Do not apply more than 24 oz AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 96 oz of AVAUNT® or 1.76 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 3 days.					

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Leafy petioles Including: Cardoon, Celery, Chinese celery, Celtuce, Florence fennel (finocchio), Rhubarb and Swiss chard	Beet Armyworm Cabbage looper	0.065	3.5	3	12 hrs
	Make no more than 4 applications per season. Do not apply more than 14 oz AVAUNT® or 0.26 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 56 oz of AVAUNT® or 1.04 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 3 days.				
Mint (Peppermint and Spearmint)	Cabbage looper Spotted cutworm	0.065	3.5	7	12 hrs
	Make no more than 4 applications per acre per crop. Do not apply more than 14 oz AVAUNT® or 0.26 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 3 days. For ground applications, apply using a minimum of 20 gallons per acre of water. AVAUNT® may be applied to mint by overhead chemigation. For specific guidance see label section titled APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH* AND SWEET CORN.				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Pear	Codling moth - East of the Rocky Mountains	0.09 - 0.11	5.0 - 6.0	28	12 hrs
	Codling moth - West of the Rocky Mountains*	0.09 - 0.11	5.0 - 6.0		
	Light brown apple moth Oriental fruit moth Pandemis leafroller (except California) Redbanded leafroller White apple leafhopper (except California)	0.09 - 0.11	5.0 - 6.0		
	<p>Make no more than 3 applications prior to hand-thinning. No hand-thinning after the 4th application. Make no more than 4 applications per season. Do not apply more than 24 oz of AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per year. Do not apply dilute applications of more than 200 gal water per acre. For best results apply 50 - 150 gal water per acre. The minimum interval between treatments is 7 days.</p> <p>*West of the Rockies. For use against low to moderate infestations in conjunction with alternate control measures such as established Mating Disruption blocks.</p>				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Pome Fruit (except pear) Including: Apple, Crabapple, Loquat, Mayhaw, and Quince	Codling moth - East of the Rocky Mountains	0.09 - 0.11	5.0 - 6.0	14	12 hrs
	Codling moth - West of the Rocky Mountains*	0.09 - 0.11	5.0 - 6.0		
	European apple sawfly (except California) Green fruitworm (except California) Lesser appleworm Light brown apple moth Oriental fruit moth Pandemis leafroller Plum curculio Potato leafhopper Redbanded leafroller Spotted tentiform leafminer - suppression only** (except California) Tarnished plant bug Tufted apple bud moth White apple leafhopper***	0.09 - 0.11	5.0 - 6.0		

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Pome Fruit (continued)	Lacanobia fruitworm (except California)	0.056 - 0.11	3.0 - 6.0	14	12 hrs
	Apple maggot****	0.11	6.0		
<p>Make no more than 3 applications prior to hand-thinning. No hand-thinning after the 4th application. Make no more than 4 applications per season. Do not apply more than 24 oz of AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per year. Do not apply dilute applications of more than 200 gal water per acre. For best results apply 50 - 150 gal water per acre. The minimum interval between treatments is 7 days.</p> <p>*West of the Rockies. For use against low to moderate infestations in conjunction with alternate control measures such as established Mating Disruption blocks.</p> <p>**Use of an adjuvant may improve performance. For best results, especially when using the lower use rate, use an adjuvant.</p> <p>***White apple leafhopper (OR and WA only)-application rates of 2.5 - 4.9 ounces per acre (0.045 - 0.089 lbs. A.I./acre) may be used for suppression of light infestations.</p> <p>****Apple maggot - apple maggot entering the orchard from border areas may not be controlled if they do not feed on treated apples prior to oviposition.</p>					

(continued)

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Spinach, New Zealand spinach, Vine spinach and Amaranth (leafy amaranth, Chinese spinach amaranth)	Beet armyworm Cabbage looper	0.065	3.5	3	12 hrs
	<p>Make no more than 4 applications per acre per crop. Do not apply more than 14 oz AVAUNT® or 0.26 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 56 oz of AVAUNT® or 1.04 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 3 days. Make sequential applications at 3 day intervals or until insect populations are brought below threshold. Use on spinach via overhead sprinkler irrigation is allowed only in the states of Arkansas, Georgia, Missouri, North Carolina, New Mexico, Oklahoma, and Texas unless otherwise permitted in supplemental labeling. For specific guidance see label section titled APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH* AND SWEET CORN.</p>				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Stone Fruit Including: Apricot, Sweet cherry, Tart cherry, Nectarine, Peach, Plum, Chicksaw plum, Damson plum, Japanese plum, Plumcot and Prune	Light brown apple moth Plum curculio	0.09 - 0.11	5.0 - 6.0	14	12 hrs
	Katydid (nymphs)* Oriental fruit moth*** Peach twig borer**	0.11	6.0		
<p>Make no more than 3 applications prior to hand-thinning. No hand-thinning after the 4th application. Make no more than 4 applications per season. Do not apply more than 24 oz of AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per year. Do not apply dilute applications of more than 200 gal water per acre. For best results apply 50 - 150 gal water per acre. The minimum interval between treatments is 7 days.</p> <p>* Forktailed bush katydid (<i>Scudderia furcata</i>) and Angularwinged katydid (<i>Microcentrum retinerve</i>) - Correct timing of spray application is to the early nymphal stages; thorough spray coverage is critical to achieve best results. Make repeat applications on a 7 to 10 day schedule if monitoring indicates continued feeding activity.</p> <p>** Oriental fruit moth (OFM) - For applications East of the Rockies: AVAUNT® is effective for control of OFM when used as part of an effective IPM program. Rotate to a product with another mode of action after each AVAUNT® application. West of the Rockies: AVAUNT® provides suppression only of OFM.</p> <p>*** Peach twig borer - AVAUNT® may be used as an early bloom or in-season spray for peach twig borer.</p> <p>AVAUNT® provides control of fruit strikes by peach twig borer and suppression of shoot strikes. Peach twig borer (dormant and delayed dormant, CA only) - Avaunt may be used as a dormant or delayed-dormant spray for the control of first generation peach twig borer. Make application with an EPA registered dormant oil; for specific recommendations on the use of oil consult the manufacturer's label. For best performance, ground application equipment is recommended.</p>					

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Tuberous and Corm Vegetables Including: Arracacha, Arrowroot, Chinese Artichoke, Jerusalem Artichoke, Edible Canna (Queensland arrowroot), Bitter and Sweet Cassava, Chayote (root), Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet Potato, Tanier (cocoyam), Tumeric, Yam Bean (jicama, manioc pea), and True Yam	Cabbage looper	0.045 - 0.11	2.5 - 6.0	7	12 hrs
	Colorado potato beetle* European corn borer (except California)	0.065 - 0.11	3.5 - 6.0		
	Potato tuberworm**	0.056 - 0.11	3.0 - 6.0		
	Make no more than 4 applications per acre per crop. Do not apply more than 24 oz of AVAUNT® or 0.44 lbs a.i. of indoxacarb containing products per acre per crop. Do not apply more than 96 oz of AVAUNT® or 1.76 lbs a.i. of indoxacarb containing products per acre per year. The minimum interval between sprays is 5 days. For POTATO only - AVAUNT® may be applied by overhead chemigation. For specific guidance see label section titled APPLICATION BY CHEMIGATION – CRANBERRY, MINT, POTATOES, SPINACH* AND SWEET CORN. <div style="text-align: right;"><i>(continued)</i></div>				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Tuberous and Corm Vegetables <i>(continued)</i>	<p>*Colorado potato beetle - In situations where Colorado potato beetle populations are known or suspected to be difficult to control with current insect control products, the inclusion of piperonyl butoxide (PBO), a synergist, with AVAUNT® may be necessary to achieve optimum control. In these situations, a combination of AVAUNT® applied at a rate of 3.5 - 6.0 oz.per acre combined with 0.25 lb a.i.per acre of PBO may be necessary to achieve the most effective control of Colorado potato beetle larvae.</p> <p>Apply the low rates on small plants, small insects and light infestations of insects. Use intermediate rates on large insects and heavier infestations of insects. Use the highest specified rate for controlling severe infestations. Apply AVAUNT® insecticide as a thorough coverage spray using properly calibrated air or ground spray equipment. Use sufficient water to obtain thorough and uniform coverage. For aerial application, use a minimum of 5 gallons of water per acre.</p> <p style="text-align: right;"><i>(continued)</i></p>				

Crops	Insects	DuPont™ AVAUNT® Rate Per Acre		Last Application (Days to Harvest)	REI
		Lbs. A.I.	Ounces		
Tuberous and Corm Vegetables (continued)	**Potato tuberworm foliar feeding larvae - AVAUNT® is most effective when applied by ground, air or overhead chemigation to vigorously growing plants through tuber bulking prior to the beginning of crop senescence. For control of potato tuberworm foliar feeding larvae, apply AVAUNT® insecticide when tuberworm larvae and/or moth counts reach locally established treatment threshold populations. AVAUNT® is absorbed into leaf tissue via translaminar movement and is most effective when applied to vigorously growing plants through tuber bulking (Growth Stage IV) prior to the beginning of crop senescence (Growth Stage V). Repeat applications of effective insecticides may be needed to keep tuberworm larvae populations as low as possible prior to harvest in order to reduce the risk of tuber damage. Failure to adequately control tuberworm larvae prior to crop senescence or vinekill increases the risk of tuber damage. To improve control of adults (moths), apply AVAUNT® in a tankmix with a pyrethroid insecticide such as "Asana" XL. Potato tuberworm is a difficult pest to control due to several factors; eggs can be laid deep in the canopy and on the underside of the leaf, and larvae feed inside the leaves prior to moving to the soil to feed on the tubers. An integrated spray approach is essential. Foliar sprays alone (ground or air) may not provide adequate control of larvae in the mid to lower crop canopy. For best results, apply via chemigation or integrate chemigation applications into the foliar spray program. Ensure thorough coverage by using sufficient spray volumes. For ground applications use at least 10 gallons of water per acre. For aerial applications, use at least 5 gallons of water per acre. For best results with foliar sprays, add Methylated Seed Oil (MSO) as a spray adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v). For chemigation applications, apply in 0.1 to 0.2 acre inch of water and add MSO at 12 to 16 fl oz/A. Do not make more than two sequential applications of AVAUNT® for control of potato tuberworm before rotating to another registered insecticide having a different mode of action.				

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity**, and **Surface Temperature Inversions** sections of this label.

Controlling Droplet Size - General Techniques

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size - Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.

- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- Do not apply as a ULV application.

BOOM LENGTH AND HEIGHT

- **Boom Length (aircraft)** - The boom length should not exceed 3/4 of the wing length, using shorter booms decreases drift potential. For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.
- **Boom Height (aircraft)** - Application more than 10 ft above the canopy increases the potential for spray drift.
- **Boom Height (ground)** Setting the boom at the lowest height which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to variable direction and inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they effect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the

movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates a surface inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

AIR ASSISTED (AIR BLAST) TREE AND VINE SPRAYERS

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream.

In addition to the general drift management principles already described, the following specific practices will further reduce the potential for drift.:

- *Adjust deflectors and aiming devices so that spray is only directed into the canopy.*
- *Block off upward pointed nozzles when there is no overhanging canopy.*
- *Use only enough air volume to penetrate the canopy and provide good coverage.*
- *Movement of spray that goes beyond the edge of the cultivated area may be minimized by practices such as spraying the outside row only from outside the planting.*

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not subject to temperatures below 32 degrees F. Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Refer to the Net Contents section of this product's labeling for the applicable "Refillable Container" or "Nonrefillable Container" designation.

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning; if burned, stay out of smoke.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

NOTICE TO BUYER--Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo, DuPont™, AVAUNT® and LANNATE® are trademarks or registered trademarks of E. I. duPont de Nemours and Company.

"Asana" is a registered trademark of Sumitomo Chemical Company, Limited.

"Asana" XL and DuPont™ LANNATE® insecticides are restricted use pesticides.

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT

Internet address: <http://cropprotection.dupont.com/>

© 2000-2015 E.I. du Pont de Nemours and Company,

Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.

All rights reserved.



DuPont™ Avaunt®

INSECTICIDE

GROUP 22 INSECTICIDE

Dispersible Granules

Active Ingredient By Weight

Indoxacarb	
(S)-methyl 7-chloro-2,5-dihydro-2-[[[methoxy-carbonyl]4(trifluoromethoxy)phenyl]amino]-carbonyl]indeno[1,2-e][1,3,4]oxadiazine-4a-(3H)-carboxylate	30%
Other Ingredients	70%
	TOTAL 100%

EPA Reg. No. 352-597

EPA Est. No. 67545-AZ-001

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See back and side panels and attached booklet for First Aid, Precautionary Statements, and Storage and Disposal.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing and/or PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, birds, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Cover, incorporate, or clean up granules that are spilled.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.

Refer to accompanying labeling for additional precautions and complete directions for use.

AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under (Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Notice to Buyer: Purchase of this material does not confer any rights under patents of countries outside of the United States.

A01780022 (SL-1970 062415 09-24-15)

Net: 1 lb., 2 oz.
Nonrefillable Container

DuPont™ Avaunt®
INSECTICIDE



FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Remove contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-441-3637** for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if inhaled. Avoid breathing (dust, vapor or spray mist). Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category A on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical Resistant Gloves Category A (such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber), all ≥14 mils.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS. **IMPORTANT:** when reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicator and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

Sold by: E.I. du Pont de Nemours and Company,
Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.
Made in U.S.A.

OPEN AT ARROW & PRESS TO RESEAL