

930 mm

120 mm

385 mm


100 mm

TRIM
10 10
mm mm

10 10
mm mm

10
mm

810 mm



Bag Description: 6#
Bag Style: Fouch
Special Information:
1x260(cut-off)

AREA: TRIM Dimensions are in millimeters

260 mm

10 mm

10 mm
25 mm

Tear Notch
25 MM From
Open End

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Caution! Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistant category selection chart.

Pilots, flaggers and groundboom applicators must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Mixers, loaders, other applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride
- A NIOSH approved dust/mist filtering respirator with any N, R, P, or HE filter or with approval number prefix TC-21C.
- Chemical resistant apron when mixing, loading, or cleaning equipment or spills.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

See "Engineering Control Statement" for additional requirements.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

Flaggers supporting aerial applications must use an enclosed cab that meets the definition in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)] for dermal protection.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Cover or incorporate spills.

Refer to accompanying labeling for additional precautions and complete directions for use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to product labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.



Apply DFU Folder Here.

Sold by: E. I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, DE 19898 U.S.A.

Made in U.S.A.

A01442987

EPA Est. 11773-IA-001

385 mm

TRIM
10 mm
10 mm
10 mm

810

Fres-go
SYSTEM USA, INC.
3005 State Road
Telford, PA 18969
215-721-4600

Customer: Dupont
Schematic #: 2182Schem41
Description: 385(face)x100(guess)x260(cut-o-
Machine(s))
Date Created: 09/08/2008

NO COPY AREA: FOLD LINES: SAFE COPY AREA:

DIRECTIONS FOR USE
A01462777 (SL - 165EA 04/25/11 04-20-11)

| | |
|--|---------------------|
| Diuron Dry Flowable Herbicide | |
| Active Ingredients | By Weight |
| Diuron: 3- [3,4-dichlorophenyl]-1,1-dimethylurea | 80.0% |
| Other Ingredients | 20.0% |
| EPA Reg. No. 352-692 | TOTAL 100.0% |

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies involving this product, call toll free 1-800-441-3637.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution: Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

- Pilots, flaggers and groundboom applicators must wear:**
- Long-sleeved shirt and long pants
 - Shoes plus socks
- Mixers, loaders, other applicators, and other handlers must wear:**
- Long-sleeved shirt and long pants
 - Shoes plus socks
 - Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinylchloride
 - A NIOSH approved dust/mist filtering respirator with any N, R, P, or HE filter or with approval number prefix TC-21C.
 - Chemical resistant apron when mixing, loading, or cleaning equipment or spills.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

See "Engineering Control Statement" for additional requirements.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

Flaggers supporting aerial applications must use an enclosed cab that meets the definition in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)] for dermal protection.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Cover or incorporate spills.

PRODUCT INFORMATION

DuPont™ KARMEX® XP herbicide must be used only in accordance with instructions on this label, or in separate published instructions.

KARMEX® XP is a dispersible granule to be mixed with water and applied as a spray for selective control of weeds in certain crops and for nonselective weed control on non-cropland areas. It is non-corrosive to equipment, non-flammable and non-volatile.

KARMEX® XP may be applied to soil prior to emergence of weeds to control susceptible weed seedlings for an extended period of time. The degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Soils high in clay or organic matter require higher dosages than soil low in clay or organic matter for equivalent herbicide performance. Moisture is required to activate the herbicide. Best results occur if rainfall (or sprinkler irrigation) occurs within 2 weeks of application.

KARMEX® XP applied before emergence of crop and weeds, is an effective profligate because susceptible weeds are controlled in an early, vulnerable seedling stage before they compete with the crop. With favorable moisture conditions, KARMEX® XP continues to control weeds for some time as the crop becomes better able to compete. Should weed seedlings begin to break through the preemergence treatment in significant numbers, secondary weed control procedures should be implemented; these include cultivation and post-emergence herbicide applications.

KARMEX® XP may also be used to control emerged weeds. Results vary with rate applied and environmental conditions. Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher. Addition of a surfactant to the spray (where recommended) increases contact effects of KARMEX® XP.

KARMEX® XP may be used as a directed postemergence application. Contact of crop foliage and/or fruit with spray or mist must be avoided on the following crops: artichoke, corn (field), cotton, sorghum (grain), sugarcane and established plantings of apples, bananas, plantains, blueberries, caneberrys, gooseberries, citrus, grapes, macadamia nuts, olives, papayas, peaches, pears, pecans, walnuts and certain tree plantings as injury may occur.

KARMEX® XP may be used on alfalfa (see CROP USES). KARMEX® XP without surfactant may be applied over the top of alfalfa established, dormant or semi-dormant, asparagus (established), birdfoot trefoil (established, dormant), grass seed crops (established), oats, red clover (established, dormant), sugarcane, wheat and pineapple.

Weed species vary in susceptibility to KARMEX® XP and they may be more difficult to control when under stress. Combinations of KARMEX® XP with other herbicides (as registered) increase the number of weed species controlled. Consult labels of the companion product for this and other information. Observe all precautions and limitations on labeling of all products used in mixtures.

Since the effect of KARMEX® XP varies with soils, uniformity of application and environmental conditions, it is suggested that growers limit their first use to small areas.

IMPORTANT PRECAUTIONS

Do not use on home plantings of trees, shrubs or herbaceous plants or lawns, walks, driveways, tennis courts or similar areas.

Do not apply this product through any type of irrigation system.

Thoroughly clean all traces of KARMEX® XP from application equipment immediately after use. Flush tank, pumps, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

Draining or flushing equipment on or near desirable trees or other plants or in areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots may injure the plants.

Trees or other desirable plants whose roots extend into a treated crop use area may be injured.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only pro- tected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for Injunctive Relief in Washington Toxics Coalition et al vs EPA, C01-132C (W.D. W.A.). For information, please refer to www.epa.gov/espp/wv/.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material.
- Shoes plus socks

SELECTIVE USE IN CROPS

Preemergence Use (Germinating Weeds): DuPont™ KARMEX® XP, at directed rates, controls annual weeds and grasses such as:

0.75 to 1 Pound/Acre

Barnyardgrass (Watergrass) Lambsquarter Purslane
Crabgrass Pigweed Ragweed

1.5 to 2 Pounds/Acre

Bluegrass, Annual Groundcherry, Annual Tansymustard
Chickweed Knaveel Velvetgrass
Corn Spurry Morningglory, Annual Vernalgrass, Sweet, Annual
Hawksbeard Pennycress Wild Buckwheat
Field Bindweed Rattail Wild Lettuce
Foxtail Red Spangletop Wild Mustard
Gromwell Shepherdspur

2 to 6 Pounds/Acre

Ageratum Marigold Ryegrass, Annual
Corn Speedwell Mexican Clover Sandbur
Dayflower Orchardgrass Smartweed, Annual
Flora's Paintbrush Pigeongrass Velvetleaf
Hawkbear Pineappleweed Spanish Needles
Johnsongrass (Seedling) Peckev Velvetleaf (Buttonweed)
Kyllinger (Kylling) Rabbit Tobacco Wild Radish
Lovegrass, Annual Ricegrass

PARTIAL CONTROL

1 Pound/Acre

Chickweed Morningglory, Sickledod
Cenchrus annual Vetchleaf
Cot. Prickly sida (teaweed) Waterhemp
Cyperus Anisima Waterhemp
Liriodendron Marestail Shepherd's-purse

4 Pounds/Acre

Horsenettle Quackgrass

8 to 10 Pounds/Acre

Guineagrass Maidencane Pangolagrass

APPLICATION INFORMATION

AERIAL APPLICATION: For alfalfa, barley (winter), cotton (preplant or preemergence only), grass seed crops (PIW only), sugarcane and wheat (winter), application may be made by aircraft at 5 to 10 gallons of water per acre. Avoid overlapping of spray swath and avoid application under conditions where excessive drift may occur. Where land is bedded, make application parallel to rows.

GROUND APPLICATION: Use a boom power sprayer properly calibrated to a constant speed and rate of delivery. Openings in screen should be 50 mesh or larger. Continuous agitation in the spray tank is required to keep the material in suspension. Agitate by mechanical or hydraulic means. If by-pass or return line is used, it should terminate at bottom of tank to minimize foaming. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping or injury to crop may result.

PREEMERGENCE: For preemergence application, use sufficient spray volume and pressure to uniformly distribute the spray solution over treated soil. Preemergence weed control will be reduced on high organic matter soils such as peat or muck.

POSTEMERGENCE: For postemergence application, use sufficient spray volume and pressure for thorough coverage of weed foliage. For selective applications and applications near sensitive crops, use low spray pressure to keep spray drift to a minimum. DuPont™ KARMEX® XP at directed rates controls seedling annual weeds such as annual morningglory, barnyardgrass (watergrass), crabgrass, crowfoot, goosegrass, pigweed and purslane. Addition of a surfactant to the spray (where recommended) increases contact effects of KARMEX® XP. Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher.

SPRAY PREPARATION: Mix proper amount of KARMEX® XP into necessary volume of water. Where use of a surfactant is recommended, dilute with ten parts of water and add as last ingredient to nearly full spray tank.

TANK MIXTURES: KARMEX® XP may be tank mixed with other herbicides and/or adjuvants registered for crop or non-crop use in this label. Refer to the label of the tank mixture partner(s) for any additional use instructions or restrictions.

REPLANTING: If initial seeding fails to produce a crop, any crop registered for the rate of KARMEX® XP that was applied may be replanted immediately. Thoroughly rework the soil before replanting. Do not retreat field with a second application as injury to the crop may result.

CROP ROTATION

Unless otherwise directed on a specific crop section of this label, do not rotate treated areas to any crop within 2 years after last application as injury to subsequent crop(s) may result.

For crops grown in the arid west and other arid areas: Reductions in normal irrigation practices for the crop in production or a summer fallow period without supplemental irrigation may require the crop rotation intervals to be extended (see specific Crop Uses sections for crop-specific rotation intervals or refer to the Crop Rotation section above).

When such conditions occur, a field bioassay should be completed prior to planting any desired crop. A successful bioassay means growing to maturity a test strip of the crop(s) intended for production. The test crop(s) strip should cross the entire field including knolls, low areas and areas where any berms were located. The results of this bioassay may require the rotation intervals to be extended.

RATES: All rates of KARMEX® XP are expressed as broadcast rates. Where band applications are specified use proportionately less. For example, use 1/3 of the broadcast rate when treating a 14 inch band where row spacing is 42 inches. When a range of dosages is given, use the lower rate on coarse textured soils low in clay or organic matter and the higher rate on fine textured soils high in clay or organic matter.

For postemergence application, use the lower rate on smaller weeds and the higher rate on larger weeds.

SOIL LIMITATIONS: Crop injury may result from failure to observe the following: Unless otherwise directed, do not use on sand, loamy sand, gravelly soils or exposed subsoils; nor on pecans where organic matter is less than 0.5%; nor on alfalfa, apples, artichoke, barley (winter), citrus, corn, cotton, grapes, oats, olives, papayas, peaches, pears, sorgho, sugarcane, walnuts and winter wheat where organic matter is less than 1%; nor on blueberries, birdfoot trefoil, caneberrys, gooseberries, macadamia nuts and peppermint where organic matter is less than 2%.

FIELD CROPS (SEE SOIL LIMITATIONS): A good seedbed must be prepared before preemergence use of KARMEX® XP, as crop injury may result if application is made to ground which is cloddy or compacted resulting in improperly planted seed. Plant seed to depth specified. Unless otherwise directed, the surface of the soil should not be cultivated or disturbed after application of KARMEX® XP and before emergence of the crop as weed control may be reduced and crop injury may result. However, if moisture is insufficient to activate the herbicide, a shallow cultivation (rotary hoe preferred) should be made after emergence of crops while weeds are small enough to be controlled by mechanical means.

FRUIT AND NUT CROPS (SEE SOIL LIMITATIONS): Unless otherwise directed, make a single application per year as a directed spray, avoiding contact of foliage and fruit with spray or drift. Do not graze livestock in treated orchards or groves.

CROP USES

ALFALFA

Treat only vigorous, healthy stands of alfalfa that have been established for at least one full growing season. Do not apply to seedling alfalfa nor to alfalfa/grass mixtures. Do not apply to alfalfa under stress from disease, insect damage, shallow root penetration (such as on shallow hard pans), alkali spots, nor to flooded fields as crop injury may result. Do not spray on snow-covered or frozen ground. In alfalfa, KARMEX® XP may only be applied once per year and not exceed 3 pounds per acre per year. Avoid overlapping of spray swaths and shut off spray booms while starting, turning, slowing or stopping or crop injury may result.

ARIZONA, CALIFORNIA, IDAHO, NEVADA, OREGON, WASHINGTON

APPLICATION INFORMATION: Severe alfalfa injury may result following application after cutting if either the regrowth is more than 2 inches high, significant stubble is left after alfalfa cutting or grazing, or the air temperature is above 90 degrees F. For best results, apply before weeds have emerged or become established (2 inches in height or diameter). Control of established weeds is improved by applying DuPont™ KARMEX® XP with a suitable contact herbicide (except Non-Dormant varieties). Sufficient rainfall or irrigation is needed for soil activation of KARMEX® XP. Treated areas may be replanted to any crop after 1 year from last application if the rate does not exceed 2 pounds per acre.

NON-DORMANT AND SEMI-DORMANT VARIETIES: Make a single application of 1.5 to 3 pounds per acre during the winter months when alfalfa plants are in the least active stage of growth.

DORMANT VARIETIES: Make a single application of 1.5 to 3 pounds per acre after alfalfa becomes dormant and before new growth exceeds 2 inches in height in the spring. Where weeds have emerged, include a surfactant or use a tank mixture with the recommended rate of paraquat.

EASTERN COLORADO, KANSAS: For control of tansymustard, apply 1 pound per acre shortly after emergence of mustard in the fall or winter. Use 2 pounds per acre if weeds are 12 to 4 inches in height. Alternatively, if other annual weeds are present, use 3 pounds per acre in February or March.

OTHER AREAS WHERE ALFALFA BECOMES WINTER DORMANT: Use 1.5 to 3 pounds per acre (1.5 to 2 pounds per acre East of Appalachian Mountains). Apply in March or early April, but before spring growth begins.

APPLE

Use KARMEX® XP only under trees established in the orchard for at least 1 year. Do not treat young trees grafted on full-dwarf root stocks. Apply 4 pounds per acre in the spring from March through May. In the Far West, apply 4 pounds per acre to small weeds less than 2 inches in height or diameter under dormant trees. Alternatively, treatments to small weeds may be applied at 2 pounds per acre post-harvest followed by 2 pounds per acre prior to bud break.

Do not apply more than 90 days per acre per year. When using KARMEX® XP in a sequential treatment program, allow a minimum of 90 days between applications. Do not make more than two applications of KARMEX® XP per year.

GEORGIA: Apply 2 to 3 pounds per acre in the spring. Repeat application in the fall but do not use more than 4 pounds per acre per year. Add a surfactant to improve control of small, emerged weeds.

KANSAS, TX PLUS TERBACIL: Apply only under trees established in the orchard for at least 2 years. Apply either in the spring or after harvest in the fall before weeds emerge or during early seedling stage of weed growth.

| | | | | | |
|-----------------------|----------------------------------|-----------------|-----------------------------|-----------------|-----|
| | RATES POUNDS PER ACRE | | | | |
| | 1 to 2% Organic Matter | | More than 2% Organic Matter | | |
| Soil Texture | Karmex® XP | Terbacil | Karmex® XP | Terbacil | |
| Sandy loam | 1.0 | + | 1.0 | + | 1.5 |
| Loam, Silt loam, Silt | 1.5 | + | 1.5 | + | 2.0 |
| Clay loam, Clay | 2.0 | + | 2.0 | + | 2.0 |

Where crop is grown under furrow irrigation or under raised-berm flood irrigation (trees 4 to 6 inches above waterline), apply only as a band treatment. Do not treat trees planted in the bottom of irrigation furrows, nor trees grown under flat flood or basin irrigation, as injury to trees may result. Where complete weed control to harvest is desired, additional weed control measures may be required during the growing season.

ARTICHOKE (CALIFORNIA)

Apply 2 to 4 pounds per acre in late fall or early winter after the last cultivation. Apply before weeds germinate or to emerging seedlings. Direct spray to cover the area between the rows and at the base of artichoke plants, keeping contact with crop plants at a minimum.

ASPARAGUS

Apply as a band or broadcast treatment. Do not apply to young plants during the first growing season (except as noted below), nor to newly seeded asparagus, nor on plants with exposed roots, as severe injury may result. Preemergence weed control will be reduced on soils with greater than 5% organic matter.

ESTABLISH PLANTINGS: On light soils and other soils low in clay or organic matter, apply 1 to 2 pounds per acre. On soils high in clay or organic matter, use 2 to 4 pounds per acre. Two applications may be used. The first application should be made before weeds become established but no earlier than 4 weeks before spear emergence and no later than the early cutting period. If weeds are controlled into the cutting period by cultural practices, application may be delayed until immediately after the last cultivation. A second application may be made immediately following completion of harvest provided rainfall is expected. When two applications are used, the second application should be made 3 to 4 weeks before application. In Washington (irrigated crop), apply a single treatment of 4 pounds per acre. If treatment is delayed until late winter or early spring, incorporation of the chemical in the top 1 to 2 inches of soil may substitute for lack of rain to activate the herbicide.

NEWLY PLANTED CROWNS

SAN JOAQUIN DELTA, CALIFORNIA: Make a single treatment of 2 to 4 pounds per acre on soils high in clay or organic matter. Use the lower rate on clay loams and the higher rate on peat soils. Do not use on soils containing more than 2% organic matter. Soil must be settled by rainfall or irrigation prior to treatment. Do not treat crowns planted to a depth of less than 2 inches.

BANANA and PLANTAIN

NEW PLANTINGS: To control annual weeds, apply 1.5 to 3 pounds per acre after planting but before weed or crop emergence. Do not apply to loose soil directly over the planting material.

ESTABLISHED PLANTINGS: For control of annuals and for top-kill of perennials such as bermudagrass, bird-seed grass and guineagrass, apply 3 to 6 pounds per acre plus surfactant. Avoid contact of banana and plantain plants with spray or drift as injury may result. Where dense weed growth is present, remove weed growth before application. If application is made to soil free of weeds, omit surfactant from the spray mixture. Repeat treatment as needed. Apply at 6 week intervals or longer for a maximum of 12 pounds of DuPont™ KARMEX® XP per acre (broadcast basis) in 12 months.

Do not replant treated areas to any crop within 2 years after last application as injury to subsequent crops may result. Exception: sugarcane or pineapple may be planted after 1 year.

BARLEY (WINTER)

WESTERN OREGON AND WESTERN WASHINGTON: For drill planted barley, make a single application of 1.5 to 2 pounds per acre as soon as possible after planting but before emergence of barley.

Do not replant treated areas to any crop within 1 year after last application as injury to subsequent crops may result.

BIRDFOOT TREFOIL (LOTUS)

WESTERN OREGON and UTAH: Treat only stands established for at least 1 year. Do not apply to seedling trefoil as injury may result. Make a single application of 2 pounds per acre when trefoil is dormant (October 15 to December 15).

Do not replant treated areas to any crop within 1 year after last application as injury to subsequent crops may result.

BLUEBERRY, CANEBERRY, GOOSEBERRY

Use only in fields which have been established for at least 1 year. Do not apply to berries interplanted with fruit trees. Do not apply to plants whose roots are exposed as injury may result. Apply as a band treatment at the base of canes or bushes. For spring application, apply before germination and growth of annual weeds.

AR, FL, GA, MS, MO, NH, NC, SC

BLUEBERRY: Apply 1.5 to 2 pounds per acre in the spring and repeat treatment after harvest in the fall. Add a surfactant to improve control of small, emerged weeds.

CALIFORNIA

BLACKBERRY, BOYSENBERRY, DEWBERRY, LOGANBERRY, RASPBERRY: For control of winter annual weeds, apply 2 pounds per acre in October or November. Repeat at the same rate in late spring to control summer annuals. A single application of 3 pounds per acre in January or February will control annual weeds in some areas, but the separate fall and spring schedule is preferred.

INDIANA, MICHIGAN, MINNESOTA, OHIO

BLUEBERRY: Apply 2 to 4 pounds per acre in late spring. Alternatively, apply 2 pounds per acre in the fall and repeat at the same rate in the spring.

RASPBERRY: Apply 3 pounds per acre in late spring.

MAINE, MASSACHUSETTS

BLUEBERRY: Apply 2 pounds per acre in late spring.

MARYLAND, NEW JERSEY

BLUEBERRY: For control of winter annual weeds, apply 2 pounds per acre from October to December, or make a single application of 2.5 pounds per acre in early to mid-spring.

WESTERN OREGON, WESTERN WASHINGTON

NEW YORK AND PENNSYLVANIA

(PERENNIAL GRASSES): Use only in established vineyards (at least 4 years old) for spot control of perennial grasses such as orchard-grass, quackgrass and ryegrass. Apply in the spring as a band treatment to ridged soil (2 to 4 inches high) under trellis at the rate of 5 pounds per acre. Band width should not exceed 30 inches. Do not apply more than once every 4 years. Use only on heavy soils types such as loams, silt loams, clay loams. Do not use in areas where grape roots are shallow or exposed because of high bedrock, poor drainage or erosion, as injury to grapevines may result.

EAST OF THE ROCKY MOUNTAINS: On soils low in clay or organic matter (1 to 2%), apply 2 to 3 pounds per acre. On soils high in clay or organic matter, apply 3 to 5 pounds per acre. Apply in the spring just prior to germination of annual weeds.

WEST OF THE ROCKY MOUNTAINS: For best results, apply during the winter months when weeds are less than 2 inches in height or diameter. Rainfall or overhead sprinkler irrigation sufficient to wet the soil to a depth of 2 inches is necessary to activate the herbicide. Abnormally heavy rainfall following application just before spring growth may move the herbicide into the root zone of grapes which could result in injury. For initial treatment apply 3 to 4 pounds per acre. Subsequent annual applications of 2 pounds per acre will usually give adequate weed control. Do not apply to vines with trunks less than 1.5 inches in diameter as injury may result.

GRASS SEED CROPS

(PERENNIAL EXCEPT WHERE SPECIFICALLY INDICATED)

Except as noted, apply only to established plantings at least 1 year old.

NOTE: Apply a single application per year at up to 3 pounds per acre. May be applied by aerial application in the Pacific Northwest only.

COLORADO, KANSAS, NEW MEXICO, OKLAHOMA: On sand bluestem, side oats grama and switchgrass apply 2 to 3 pounds per acre during the dormant period shortly before weed seedlings emerge. Do not apply after crop begins growth in the spring as crop injury may result. In fields where ash residues have accumulated from burning straw use 3 pounds per acre. Spread unburned chaff or straw with a harrow or chopper before application.

EASTERN OREGON, EASTERN WASHINGTON: On perennial bluegrass and fescue apply 1.0 to 3.0 pounds per acre as broadcast in enough diluent to get even distribution. Apply in spring before rapid growth of the crop begins and when the windgrass is still small (1-4 leaf). DO NOT use on coarse (sax) textured soils.

WESTERN OREGON/WESTERN WASHINGTON: On alta fescue, Astoria bentgrass, Highland bentgrass, Kentucky bluegrass (Kenton bluegrass) and orchardgrass apply 2 to 3 pounds per acre between October 1 and November 15. In fields where ash residues have accumulated from burning straw, use 3 pounds per acre. Spread unburned chaff or straw with a harrow or chopper before application. For best results apply as soon as possible after fall rains start. Established weeds beyond two to four leaf stage should be removed prior to treatment.

Well established vigorous stands of spring planted alta fescue, Kentucky bluegrass and orchardgrass may be treated the following fall provided the crop is planted before April 1 and treatment is not applied before October 15. apply 2 pounds per acre.

OREGON, WASHINGTON: Apply in the fall to perennial ryegrass at the rate of 1 to 2 pounds per acre and to tall fescue at the rate of 2 to 3 pounds per acre. Use a sufficient volume of water, a minimum of 25 gallons per acre, for thorough coverage of weed foliage. For best results, make applications at the onset of the fall rains and before weeds have become established (typically October 1 through November 15). Established weeds beyond the 2-4 leaf stage should be removed prior to treatment. Apply only to well established, vigorous stands. Do not apply to perennial ryegrass stands less than 1 year old. Use mechanical agitation and avoid overlap of spray patterns. Weed control efficacy may be reduced in fields where ash residues have accumulated from burning straw.

ANNUAL RYEGRASS FOR THE CREATION OF ROWS: Apply 1 to 2 pounds per acre as a directed or shielded spray so the intended crop row area is not treated. These applications should be made where excessive populations of annual ryegrass are anticipated to volunteer from previous crops. Applications can be made as a directed/shielded spray during seeding or after emergence of annual ryegrass. These applications generally will occur between October 1 and January 15. DuPont™ KARMEX®XP is most effective when applied before annual ryegrass volunteer plants have been more than 2 leaves. If larger plants are to be treated, addition of a labeled postemergence herbicide, will provide more effective control.

Adjuvants and sprigant to allow the establishment of the desired row width (generally about 3 inches) and spacing (generally 9 to 12 inches). Use of low pressure nozzles, shielded nozzles, or drop nozzles to reduce spray movement into the intended crop row area is recommended.

FINE FESCUE GRASS SEED CROPS (INCLUDING CHEWINGS, CREEPING RED AND HARD FESCUE TYPES): For the suppression of rattail fescue, apply at 1.0 to 2.0 pounds per acre on soils having at least 1% organic matter. Do not use on sand, loamy sand, gravelly soils or exposed subsols.

KARMAE™ XP is for use on sandy soils with other herbicides. **KARME™ XP** can be applied to stands established at least 1 year or to new plantings that have been established for at least 6 months and have a minimum of eight tillers at time of application.

Apply in fall before grass weeds are beyond the one to two leaf stage and before broadleaf weeds are larger than 1 to 2 inches tall or across. Use the high end of the rate range for large weeds or where weed populations are high.

Approximately 1/2 to 1 inch of rainfall or sprinkler irrigation is needed to move **KARME™ XP** into the weed zone before weeds develop an established root system. Weeds larger than the size indicated or those having a well established root system before **KARME™ XP** is properly applied by rainfall/irrigation may not be adequately controlled.

Weed control may be reduced by heavy straw residues or ash from field burning.

TALL FESCUE

MISSOURI, NORTHERN ARKANSAS, EASTERN KANSAS

On soils having at least 1% organic matter, apply **KARME™ XP** at 2.0 pounds per acre in the fall for general broadleaf weed and grass control and the suppression of downy brome in established Tall Fescue grow for seed production. Treat only well established vigorous stands of tall fescue and begin applications at the onset of fall rains. For best results, rainfall of 1/2 to 1 inch is needed within two weeks after application. Do not treat the plants in the fall, but do treat until the following fall; for spring planted tall fescue, do not treat in the fall of the same year, extend the tall fescue establishment interval until the next fall.

Avoid overlapping of spray swaths and shut off spray booms while starting, turning, slowing or stopping as severe crop injury or death may result.

Do not use on sand, loamy sand, gravelly soils or exposed sub-soils.

TANK MIXTURES: **KARME™ XP** can be applied either alone or in a program involving tank mixes with other herbicides and adjuvants on soils with other herbicides. **KARME™ XP** can be applied to stands unless prior experience indicates it is safe to use higher rates. Tank mixes with other herbicides can increase the risk of crop injury. When using certain tank mix for the first time, limit use to a small area to determine safety before treating large areas.

USE PRECAUTIONS

Do not replant treated areas to any crop within 2 years of last application as injury to subsequent crops may result.

Do not apply to snow covered or frozen ground as injury to the crop or poor weed control may result. Do not treat stands lacking in vigor due to poor fertility, environmental stress, insect or disease or damage from other herbicides.

NEW PLANTINGS

OREGON, WASHINGTON: For use in newly planted bentgrass, chewing fescue, Kentucky bluegrass, perennial ryegrass, orchardgrass and tall fescue. During planting operation, spray a suitable band of activated charcoal at a 1 inch band on soil surface at 15 pounds per acre of crop where row spacing is 20 inches (300 pounds per acre) for use on sandy soils with other herbicides. **KARME™ XP** can be applied to stands established with DuPont™ KARMEX®XP as a single broadcast spray at the rate of 2.5 to 3 pounds per acre. Apply as soon as possible after planting but before crops or weeds emerge and before rains or sprinkler irrigation. Fall or spring plantings may be treated. Best results usually occur with early fall plantings. Treatment will not control downy brome or wild oats.

MACADAMIA NUT

HAWAII: Use only under trees established in the orchard for at least 1 year. Apply at 2 to 6 pounds per acre immediately after harvest, preferably before weeds emerge. If weeds have emerged, add surfactant. Retreat as needed but do not exceed 10 pounds per acre per year.

OATS

Do not replant treated areas to any crop within 1 year of last application as injury to subsequent crops may result.

SPRING OATS - DRILL PLANTED

IDAHO, EASTERN OREGON, EASTERN WASHINGTON: Use in areas where average annual rainfall exceeds 16 inches. Make a single application of 1 to 1.5 pounds per acre after planting, either before or after oats emerge but within 6 weeks of planting. Best results are usually obtained when application is made 3 to 4 weeks after planting. Apply before weeds are 3 to 4 inches in height.

WINTER OATS - DRILL PLANTED

MIXED WITH PEAS OR VETCH

WESTERN OREGON, WESTERN WASHINGTON: Make a single application of 1.5 to 2 pounds per acre as soon as possible after planting but before crop emergence.

OLIVE (CALIFORNIA)

Use only under trees established in the grove for at least 1 year. Apply 2 pounds per acre after the grove has been laid-up in final form in late October or November. Repeat at same rate in March or April. Remove weed growth prior to treatment.

PAPAYA

For trees established in the orchard for at least 1 year. Apply 2.5 to 5 pounds per acre, preferably before weeds emerge. If weeds have emerged, add surfactant.

For papaya trees less than one year old, apply 2.5 to 5 pounds per acre of **KARME™ XP** as a post-plant treatment between the rows. Use only in orchards that are lined with mulch paper in the crop row. Treat up to mulch paper only.

Apply preemergence or early postemergence in sufficient spray volume to provide uniform coverage of the weeds and/or soil and to allow proper dispersion and suspension of the product in the spray tank.

Do not apply more than 5 pounds per acre per year.

Do not allow spray to contact papaya foliage or other desirable vegetation.

Do not graze livestock in treated areas.

PEAS (AUSTRIAN FIELD)

WESTERN OREGON: **KARME™ XP** is for selective control of certain weeds in Austrian field peas.

Apply 1.5 to 2 pounds **KARME™ XP** per acre as a broadcast spray with air or ground equipment as soon as possible after planting but before crop emerges for control of weeds such as chickweed, shepherdspurse, wild mustard, fiddleneck, lambsquarters, pigweed and annual bluegrass. Use lower rate on coarse-textured soils and high rates on fine-textured soils. Do not use **KARME™ XP** on sand, sandy loam, gravelly soils or exposed subsols or on soils having less than 1% organic matter as crop injury may result. Do not replant treated area to another crop within one year of application. Crop injury may result if severe winter stress, disease or insect damage to the crop follows application.

PEACH

Use **KARME™ XP** only under trees established in the orchard for at least 3 years. Apply 2 to 2.75 pounds per acre in the early spring before weeds emerge or during the early seedling stage of weed growth. In California, apply 2 to 3.75 pounds per acre.

For peaches grown east of the Rocky Mountains, do not apply within 20 days of harvest.

In the Far West, do not apply within 8 months of harvest.

Where crop is grown under furrow irrigation or under raised-berm flood irrigation (trees 4 to 6 inches above water line), apply only as a band treatment. Do not treat trees planted in the bottom of irrigation furrows, nor trees grown under flat flood or basin irrigation, as injury to trees may result. Where complete weed control to harvest is desired, additional weed control measures may be required during the growing season. **GENERAL PRECAUTIONS:** On trees treated 2 years, apply 2 to 2.75 pounds per acre in the spring. Repeat application in the fall but do not exceed 5.0 pounds per acre per year. Add surfactant to improve control of small, emerged weeds.

DUPONT™ KARMEX®XP PLUS TERBACIL: Use only under trees established in the orchard for at least 2 years. Apply either in the spring or after harvest in the fall before weeds emerge or during early seedling stage of weed growth.

| Soil Texture | RATE POUNDS PER ACRE | | |
|-----------------------|------------------------|-----------------------------|-----|
| | 1 to 2% Organic Matter | More Than 2% Organic Matter | |
| Sandy loam | 1.0 | 1.5 | 2.0 |
| Loam, Silt loam, Silt | 1.5 | 2.0 | 2.0 |
| Clay loam, Clay | 2.0 | 2.0 | 2.0 |

PEAR

Use only under trees established in the orchard for at least 1 year. Do not treat varieties grafted on full-dwarf root stocks. Apply 4 pounds per acre in the spring from March through May. In the Far West, apply 4 pounds per acre to weeds less than 2 inches in height or diameter under dormant trees. Alternatively, apply to small weeds at 2 pounds per acre postharvest followed by 2 pounds per acre prior to budbreak.

PECAN

Make a single band or broadcast application as a directed spray using a minimum of 30 gallons of water per acre. Apply in the spring before weeds emerge or during the early seedling stage of growth.

| Soil Texture | RATE POUNDS PER ACRE | | |
|-----------------------|----------------------|-------------------------|-------------|
| | Karmex™ XP Alone* | OR KARME™ XP + Terbacil | Tank mix ** |
| Sandy loam | 2.0 | 1.5 | 1.5 |
| Loam, Silt loam, Silt | 3.0 | 1.75 | 1.75 |
| Clay loam, Clay | 4.0 | 2.0 | 2.0 |

* Use only under trees established in the grove for at least 3 years, and on soils with at least 0.5% organic matter. ** Use only under trees established in the grove for at least 1 year, and on soils with at least 1% organic matter. **NOTE:** Do not use on eroded areas where subsoil or roots are exposed, nor on trees that are diseased or lacking in vigor or on trees planted in irrigation furrows as injury may occur.

PEPPERMINT, SPEARMINT

Apply **KARME™ XP** at 0.75 to 1.0 pound per acre on soils having 1.0% to 2.0% organic matter. Apply **KARME™ XP** at 1.0 to 2.0 pounds per acre on soils having 2.1% to 3.0% organic matter. Apply **KARME™ XP** at 2.0 to 3.0 pounds per acre on soils having more than 3.0% organic matter.

USE PRECAUTIONS Do not apply to stands of mint suffering from stress due to low fertility, drought, winter injury, insects, disease or damage from other herbicides or other causes.

Do not apply to snow covered or frozen ground as injury to the crop or poor weed control may result. Do not apply to sand, loamy soil, gravelly soils or exposed subsols. Do not apply to soils that have a high salt content and/or high water table or poor drainage that retards mint root development resulting in a shallow root system. Do not apply to soils having less than 1% organic matter.

APPLICATION TIMING: Apply **KARME™ XP** to established (at least one year) stands of mint during the late winter dormant period or after flaming in the spring prior to the emergence of new growth. Do not cultivate after application.

If weeds are present at time of application, the use of a surfactant at 0.25% volume/volume or crop oil concentrate at 1.0% volume/volume may be used to increase the performance of **KARME™ XP** postemergence to weeds.

TANK MIXES AND SEQUENTIAL TREATMENTS: **KARME™ XP** can be applied either alone or in a program involving tank mixes and/or sequential treatments with other herbicides and adjuvants providing **KARME™ XP** is not applied to actively growing mint plant.

When using a tank mix with other herbicides, use the lower end of the **KARME™ XP** use rate range unless prior experience indicates it is safe to use higher rates. Tank mixes and sequential treatments with other herbicides can increase the risk of crop injury. When using a certain tank mix or sequential treatment for the first time, limit use to a small area to determine safety before treating large areas.

PINEAPPLE

HAWAII: Apply 2 to 6 pounds per acre as a broadcast spray just before or immediately after planting but prior to weed emergence. Use 2 to 4 pounds per acre after harvesting the plant crop or ratoon crop (for first ratoon crop as well as subsequent ratoon crops) but before differentiation.

For plant crop only, additional broadcast or interspace applications may be made prior to differentiation at the rate of 2 pounds per acre at intervals of not less than 2 months. Additional applications to plant crop may be made as needed to interspace only using 2 pounds per acre. Do not apply more than 12 pounds per acre as broadcast sprays nor more than 16 pounds total per acre per plant crop. Treated areas may be planted to pineapple or sugarcane 1 year after last application.

FLORIDA: Apply 4 to 8 pounds per acre as a broadcast spray just before or immediately after planting but prior to weed emergence. For ratoon crop use 4 pounds per acre after harvesting plant crop. For plant crop only, a second and third broadcast or interspace application may be made prior to differentiation at the rate of 2 pounds per acre at intervals of not less than 2 months. Additional applications to plant crop may be made as needed to interspace only using 2 pounds per acre. Do not apply more than three broadcast sprays (maximum 12 pounds per acre) prior to differentiation nor more than 16 pounds total per acre per plant crop. Treated areas may be planted to pineapple or sugarcane 1 year after last application.

PUERTO RICO: Apply 3.75 to 6.25 pounds per acre as a broadcast spray before or immediately after planting but prior to weed emergence. Preemergence application controls weeds such as pigweed, crotalaria, morningglory, purslane, crabgrass, foxtail, goosegrass, fall panicum and sourgrass.

PRICKLY PEAR CACTUS

ARIZONA, CALIFORNIA, NEW MEXICO, AND TEXAS: DuPont™ KARMEX® XP herbicide can be used for control of little nelly, shepherd's purse, and common mustard in prickly pear cactus in the States of Arizona, New Mexico and Texas.

Use only on established prickly pear cactus. Apply 4 pounds per acre as directed spray to the lower portion of the cactus plant. Do not spray over the top of the cactus plant. Apply with ground application equipment only. Do not apply within 7 days of harvest. Harvest may include cactus fruit and pads.

RED CLOVER

WESTERN OREGON: Make a single application of 2 pounds per acre on established red clover stands at least 9 months old. Apply when red clover is dormant between October 15 and December 15. Do not apply to seedling red clover. Do not replant treated area to any crop within 1 year after last application as injury to subsequent crops may result.

Treatment will control annual weeds such as bluegrass, chickweed, hawksbeard, rattail fescue, ryegrass and velvetgrass.

SORGHUM (GRAIN)

SOUTHWESTERN STATES: Apply 0.25 to 0.5 pounds per acre plus surfactant. Apply as a directed postemergee spray after sorghum is 15 inches tall to control weeds 2 to 4 inches in height. Use lower rate on broadleaf weeds up to 2 inches tall. Use the higher rate on grasses up to 2 inches and broadleaf weeds up to 4 inches tall. When the lower rate is used, a second application may be made if needed. Do not exceed 0.5 pound per acre. Treatment of weeds under drought stress is usually ineffective.

Do not replant treated areas to crops other than corn within 4 months following band treatment and 6 months following broadcast treatment as injury to subsequent crops may result.

DO NOT SPRAY OVER THE TOP OF SORGHUM.

SUGARCANE

To prevent post-harvest injury on new cane varieties, test tolerance to DuPont™ KARMEX® XP prior to adoption as field practice. Do not treat sugarcane growing in tillers covered sub-soils or rocky areas as crop injury may result. Temporary chlorosis and stunting of the crop may result from application over emerged cane. Application over emerged cane should be made only as directed below, without the addition of a surfactant or crop oil concentrate. To minimize chlorosis and stunting, use directed postemergence sprays.

KARME™ XP may be applied as a directed spray (including hooded and shielded spray) in combination with post-harvest. Consult the label of the tankmix partner for rates and timings of application, restrictions, and precautions.

PREEMERGENCE - FLORIDA: For high organic soils, apply 2 to 4 pounds per acre as a broadcast or band spray prior to weed emergence after planting or after harvesting plant crop (for ratoon crop).

POSTEMERGENCE - FLORIDA: Make one or two applications of 2 pounds per acre as needed by directed spray or broadcast. Alternatively, for panicle control, make up to three applications of 0.5 to 1 pound per acre plus surfactant as a directed spray after cane has emerged but before panicle exceeds 2 inches in height. Adjust nozzles to spray beneath cane plants and between rows to cover weed foliage and to minimize control of cane leaves with spray or drift. Do not apply more than 6 pounds total per acre between planting (or ratooning) and harvest.

HAWAII: Apply 2 to 6 pounds per acre as a broadcast spray prior to weed emergence after planting or after harvesting plant crop or ratoon crop. Sequential applications of 2 to 4 pounds per acre may be made as a broadcast spray over emerged cane or by directed spray inter-row.

If weeds are emerged, add a surfactant and apply as a directed spray.

Do not apply more than three treatments nor more than 12 pounds per acre in Hawaii between planting (or ratooning) and harvest. Treated areas may be replanted to sugarcane or pineapple 1 year after last application.

LOUISIANA, TEXAS: Apply 3 to 5 pounds per acre as a broadcast spray prior to weed emergence.

KARME™ XP may be applied as a broadcast spray after planting and following the harvest of sugarcane. DuPont™ KARMEX® XP may also be applied broadcast in late winter. Application is best when made prior to weed emergence.

KARME™ XP may be applied as a post-directed spray immediately after the last cultivation. Direct the spray application to the base (no more than 1/3 the plant height) of the sugarcane plants. When small weeds (3 inches or less) are present at application, add a surfactant at 0.25% V/V or crop oil concentrate at 1.0% V/V to the spray mix.

USE PRECAUTIONS

Temporary leaf yellowing may occur following application. Do not apply more than 7.5 pounds per acre broadcast per year. Use proportionately less for band applications.

TREE PLANTINGS

CO, MT, NE, ND, SD, WY: Use only under established plantings 1 year or older of American elm, caragana, cottonwood, Douglas fir, green ash, honeysuckle, Ponderosa pine, red cedar, Russian olive and Siberian elm. Use 2.5 to 5 pounds per acre as a band 4 feet wide on each side of row for elm. Use 1 to 2 pounds per acre for other species. 1 ounce **KARME™ XP** treats 135 feet of tree row (2 feet on each side of row) at the rate of 5 pounds per acre. Apply as a directed spray in early spring before weeds emerge and before trees leaf out. Do not apply to foliage of trees, nor under trees growing in low areas as injury may result.

ID, NM, OR, WA: **KARME™ XP** is for control of weeds to aid in the establishment of hybrid poplar plantings. Apply at 1.0 to 3.0 pounds per acre depending upon soil texture and organic matter content. Use 1.0 to 2.0 pounds per acre on coarse textured soils and 2.0 to 3.0 pounds per acre on medium to fine textured soils. Do not use on gravelly soils or on any soil having less than 0.5% organic matter as injury to trees may result. **Injury may result from applications to poplar plantings grown on sandy soil with low organic matter with sprinkler irrigation.** When applied in a band, the application rate will be in proportion to the area banded on a per acre basis.

HAWAII: Apply in late winter or early spring as a uniform broadcast spray before or after planting but prior to bud swell, or as a directed spray after bud swell. Apply before weeds emerge or after emergence while weeds are small. Some rainfall or water is necessary to move **KARME™ XP** into the weed root zone before weeds become well established. If weeds are present at time of treatment, add a surfactant at 1 to 2 quarts per 100 gallons of spray solution.

PLANT: Take precautions to prevent treated soil (usually top 1 inch) from coming into contact with roots of trees during the planting process as injury may result.

POST-PLANT (BROADCAST): It is best to wait until rain or irrigation has settled the soil around the newly planted trees before applying **KARME™ XP**. If trees are dormant, a broadcast application can be made.

POST-PLANT (DIRECTED): If buds have started to swell, use a directed spray pattern that prevents **KARME™ XP** from contact with trees as injury may result. During the growing season (from bud swell to leaf drop) **KARME™ XP** may be applied to the trunk and 2.0 to 3.0 pounds per acre on medium to fine textured soils. **KARME™ XP** can be tank mixed with a registered glyphosate herbicide pre-plant and as a directed spray to broaden the spectrum of weeds controlled and improve post-emergence activity. Use 1.0 to 3.0 pounds **KARME™ XP** plus glyphosate herbicide (according to label instructions) depending upon soil type and weeds to be controlled. Note: There are several formulations of glyphosate herbicide. Check the glyphosate herbicide label to be used. Do not use a pre-plant or post-directed spray on hybrid poplar plantations as injury may result. Avoid contact of glyphosate herbicide with foliage, green stems, trees or other desirable vegetation because severe damage or destruction may result.

WALNUT (ENGLISH)

CALIFORNIA, OREGON, WASHINGTON

Use only under trees established in orchards for at least 1 year. As an initial treatment, apply 2.75 pounds per acre after the orchard has been laid-up in final form (non-tillage program) in late fall or early winter. Retreat annually with 2 to 2.75 pounds per acre. In California apply 2 to 3.75 pounds per acre. Alternatively, apply 2 pounds per acre in October or November and repeat at the same rate in March or April.

Do not use on sand, loamy sand, gravelly soils or exposed sub-soils, nor where organic matter is less than 1%.

Do not make more than two applications per year. Do not apply more than 4 pounds per acre per year. In California, do not apply more than 3.75 pounds per acre per year. When using **KARME™ XP** in a sequential treatment program, allow a minimum of 150 days between applications.

WHEAT (WINTER)

USE PRECAUTIONS

Crop injury may result where severe winter stress, disease or insect damage follows application. Winter-sensitive varieties may be less tolerant of **KARME™ XP** than winter-hardy varieties. Crop injury may result from failure to observe the following: Do not use on sandy or loamy sand soils, nor on gravelly or sandy loams with less than 1% organic matter. Do not use on thin covered or exposed sub-soil areas (gray knolls). Do not treat wheat in areas where there are pre-treat or post-treat herbicide sprays or herbicide plantations as injury may result. Do not treat wheat plants lacking in vigor due to poor emergence, insect damage, disease, high alkalinity or other causes. Do not apply after the wheat has reached the 'boot' stage of maturity. Unless specified otherwise, do not use with surfactants or nitrogen solution. Do not replant treated areas to any other crop within 1 year after last treatment (except as noted) as injury to subsequent crops may result. Do not apply more than one application per year season.

IDAHO, OREGON, WASHINGTON

EAST OF CASCADE RANGE: Where average annual rainfall exceeds 16 inches, make a single application of 1 to 1.5 pounds per acre.

FALL TREATMENT:

For early fall planted wheat (seeded before September 10), apply 3 to 6 weeks after planting but before weeds are 3 to 4 inches tall. Treatment after October 1 has generally given best results. Application should not be made after soil freezes in the fall. Wheat planted in late October should not be treated until the following spring.

SPRING TREATMENT

Apply as soon as wheat starts to grow. Treatment made prior to April 10 will usually give good results, provided winter growth is not less than 4 inches tall. Application later than May 1 may give poor results. Alternatively, make a single application of 0.5 to 1 pound DuPont™ KARMEX®XP plus 0.25 pound bromoxynil per acre as a tank mixture, either in the fall after wheat has emerged but before soil freezes or in the spring as soon as soil thaws. Apply before weeds are more than 2 inches tall or across.

Where average annual rainfall is 10 to 16 inches following fall planting, make a single application of 1 to 1.5 pounds per acre when sufficient moisture is available to germinate wheat seed. Apply before soil freezes and weeds are 2 inches tall. Application later than March 1 may give poor results.

If fall-planted wheat fails to grow due to winter kill or adverse growing conditions after fall treatment, only fields treated before November 1 may be replanted to spring wheat. Spring wheat should not be planted before April 1 and only after deep discing and plowing to a depth of 4 to 6 inches prior to planting. Do not make a second application during the same crop year or injury to the crop may result.

OREGON, WASHINGTON

WEST OF CASCADE RANGE: Make a single application of 1.5 to 2 pounds per acre as soon as possible after planting. If wheat and weeds have emerged, apply before weeds are 3 to 4 inches tall. Alternatively, apply a tank mixture of **KARME™ XP** plus bromoxynil as detailed above for East of Cascade Range.

OTHER AREAS OF OREGON AND WASHINGTON: Make a single application in the spring as soon as wheat (fall-planted) starts to grow and before weeds are 2 inches tall. Application later than May 1 may give poor results.

KANSAS, OKLAHOMA, TEXAS: Do not use on sand or sandy loam soils. Use 1 pound per acre on silt and silt loam soils and 1.5 to 2 pounds per acre on clay, clay loam and silt clay loam soils.