



DuPont™ Velpar® DF CU

HERBICIDE



LICENSED

PERIOD 2015-2017 LIC. NO.

9200.218

RESTRICTED USE PESTICIDE

GROUP 5 HERBICIDE

Dispersible Granules

Active Ingredient

Hexazinone

[3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1H,3H)-dione] 75%

Other Ingredients 25%

EPA Reg. No. 352-911

EPA Est. No. 11773-IA-001

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for medical emergencies involving this product.

See back panel for additional Precautionary Statements.

Net: 4 lb Nonrefillable Container

DuPont™ Velpar® DF CU

HERBICIDE



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive, causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.
- Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product and as soon as possible wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

The active ingredient, hexazinone, in this product is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

See Directions for Use in Supplemental Labeling attached.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

APPLY DIRECTIONS FOR USE
FOLDER HERE

Sold by: E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805

A01789390

Made in U.S.A.



DuPont™ Velpar® DF CU

HERBICIDE RESTRICTED USE PESTICIDE

GROUP **5** HERBICIDE

A01789407
(SL-1948 060415 06-11-15)

Dispersible Granules

Active Ingredient	By Weight
Hexazinone	
[3-cyclohexyl-6-(dimethylamino)-1-methyl-1,3,5-triazine-2,4(1H,3H)-dione]	75%
Other Ingredients	25%
	TOTAL 100%

EPA Reg. No. 352-911

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for medical emergencies involving this product.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive, causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:
Long-sleeved shirt and long pants.
Shoes plus socks.
Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product and as soon as possible wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

The active ingredient, hexazinone, in this product is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. DuPont™ Velpar® DF CU herbicide, referred to below as DuPont™ Velpar® DF CU, Velpar® DF CU herbicide or Velpar® DF CU, must be used only in accordance with instructions on this label, or in supplemental DuPont labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

The correct use rates by crop and geographical area, specified on the label, and proper mixing/loading site considerations and application procedures must be followed to minimize potential for hexazinone movement into ground water. Users are encouraged to consult with their state Department of Agriculture, Extension Service, or other pesticide lead agency for information regarding soil permeability, aquifer vulnerability, and best management practices for their area.

PRODUCT INFORMATION

Velpar® DF CU herbicide is a water-dispersible granule that is mixed in water and applied as a spray for weed control in certain crops. Velpar® DF CU is an effective general herbicide providing both contact and residual control of many annual and biennial weeds and woody plants. It is also effective for control of most perennial weeds.

Velpar® DF CU is noncorrosive to equipment. Care must be exercised when applying Velpar® DF CU near desirable trees or shrubs as they can absorb Velpar® DF CU through roots extending in to treated areas.

This product may be applied on agricultural sites that contain areas of temporary surface water caused by collection of water between planting beds, in equipment ruts, or in other depressions created by management activities. It is permissible to treat intermittent drainage, intermittently flooded low lying sites, seasonally dry flood plains and transitional areas between upland and lowland sites when no water is present. DO NOT make applications to natural or man-made bodies of water such as lakes, reservoirs, ponds, streams and canals.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

Velpar® DF CU is absorbed through the roots and foliage. Moisture is required to activate Velpar® DF CU in the soil. Best results are obtained when the soil is moist at the time of application and 1/4-1/2 inches of rainfall occurs within 2 weeks after application.

For best results, apply Velpar® DF CU preemergence or postemergence when weeds are less than 2 inches in height or diameter. Herbicidal activity is most effective under conditions of high temperature (above 80 °F), high humidity, and good soil moisture. Herbicidal activity may be reduced when vegetation is dormant, semi-dormant, or under stress (e.g. temperature or moisture).

Herbicidal activity will usually appear within 2 weeks after application to susceptible plants under warm, humid conditions; while 4-6 weeks may be required when weather is cool or dry, or when susceptible plants are under stress. If rainfall after application is inadequate to activate Velpar® DF CU in the soil, plants may recover from contact effects and continue to grow.

On woody plants, symptoms usually appear within 3-6 weeks after sufficient rainfall has carried the herbicide into the root zone during periods of active growth. Defoliation and subsequent refoliation may occur, but susceptible plants are killed.

The degree and duration of control will depend on the following:

- Use rate
- Weed spectrum and size at time of application
- Environmental conditions at and following treatment

Where a rate range is shown, use the higher levels of the dosage range on hard-to-control species, fine-textured soils, or soils containing greater than 5% organic matter or carbon. Use the lower levels of the dosage range on coarse-textured soils and/or on soils low in organic matter. Refer to specific uses for rate ranges.

APPLICATION INFORMATION

DuPont™ Velpar® DF CU may be applied by ground equipment and, where permitted, aerial equipment. Use rates, minimum spray gallonage, and other application information are described for various uses.

Dispose of the equipment washwater by applying it to a use-site listed on this label or in accordance with directions given in the "Storage and Disposal" section of this label.

Before spraying, calibrate equipment to determine the quantity of water necessary to uniformly and thoroughly cover the vegetation and soil in a measured area to be treated. Make sure the volume of water is sufficient to completely suspend the Velpar® DF CU.

TANK MIXTURES

Velpar® DF CU herbicide may be tank mixed with other herbicides and /or adjuvants registered for the crops specified in the label. Refer to the label of the tank mix partner(s) for any additional use instructions or restrictions. The most restrictive label provisions apply. If other label instructions conflict with this avel do not tank mix the herbicide and/or adjuvant with Velpar® DF CU herbicide.

RESISTANCE

Velpar® DF CU, which contains the active ingredient hexazinone, is a Group 5 herbicide based on the mode of action classification system of the Weed Science Society of America.

When herbicides with mode of action classifications that affect the same biological sites of action are used repeatedly over several years to control the same weed species in the same treatment area, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that area. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different biological site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide instructions available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

AGRICULTURAL USES

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

ALFALFA

DuPont™ Velpar® DF CU is labeled for control of certain weeds in established alfalfa grown for hay or seed production.

- Do not apply within 30 days of harvest (cutting for hay), or feeding of forage or grazing.
- Do not exceed 2 pounds per acre per application.
- Do not exceed 2 pounds (1.5 pounds active ingredient hexazinone) per acre per year.

APPLICATION INFORMATION

NON-DORMANT AND SEMI-DORMANT VARIETIES
In the following states, make a single application of Velpar® DF CU during the winter months when alfalfa plants are in the least active stage of growth.

Arizona	Montana	Oklahoma	Washington
California	Nebraska	Oregon	Wyoming
Colorado	Nevada	South Dakota	
Idaho	New Mexico	Texas	
Kansas	North Dakota	Utah	

In the following states, make a single application of Velpar® DF CU either in the spring before new growth exceeds 2 inches in height or to alfalfa stubble after cutting, following hay removal and before regrowth exceeds 2 inches in height.

Arkansas	Maine	New Jersey	Vermont
Connecticut	Maryland	New York	Virginia
Delaware	Massachusetts	North Carolina	West Virginia
Illinois	Michigan	Ohio	Wisconsin
Indiana	Minnesota	Pennsylvania	
Iowa	Missouri	Rhode Island	
Kentucky	New Hampshire	Tennessee	

NOTE: Severe alfalfa injury may result following application, if after cutting the regrowth is more than 2 inches high, or there is significant stubble left after cutting or grazing, or the air temperature is above 90 °F.

DORMANT VARIETIES
Make a single application of Velpar® DF CU after alfalfa becomes dormant and before new growth exceeds 2 inches in height in the spring. Where weeds have emerged, use a surfactant.

USE RATES

Use higher rates on hard-to-control species, (see **Weeds Controlled** section below) fine textured soils, soils containing greater than 5% organic matter, or under adverse environmental conditions such as temperature extremes or when weeds are stressed due to low rainfall.

For dormant alfalfa, use a surfactant approved for crops at the rate of 0.25% v/v (1 quart per 100 gallons of spray solution). Select the appropriate rate for soil texture and organic matter content as follows:

Soils	DuPont™ Velpar® DF CU (Lb/Acre) Percent Organic Matter in Soil		
	<1%	1-5%	>5%
Coarse Texture			
Loamy sand, sandy loam	2/3 - 1	2/3 - 1	1 1/3 - 2
Medium Texture			
Loam, silt loam, silt, clay loam, sandy clay loam	2/3 - 1	1 - 2	1 1/3 - 2
Fine Texture			
Silty clay loam, sandy clay, silty clay, clay	1 - 2	1 - 2	1 1/3 - 2

- NOTE:**
- In the states of MT, ND, SD, and WY, do not exceed a use rate of 1 1/3 pounds per acre on medium and fine textured soils.
 - In the state of Montana (MT), do not apply to soils with less than 1.5% organic matter.
 - In the state of Wyoming (WY):
Do not apply to soils with less than 0.5% organic matter.
Apply to irrigated alfalfa only.

WEEDS CONTROLLED

Velpar® DF CU, when applied preemergence or early postemergence at the following rates, is labeled for the control or suppression of the following species in alfalfa:

1/3 - 2/3 Lb/Acre			
Tansymustard	<i>Descurainia pinnata</i>		
2/3 - 1 1/3 Lb/Acre			
Bluegrass, annual	<i>Poa annua</i>	Mustard, Jim Hill (tumble)	<i>Sisymbrium altissimum</i>
Brome, downy (cheatgrass)	<i>Bromus tectorum</i>	Mustard, wild	<i>Brassica kaber</i>
Buckwheat, wild	<i>Polygonum convolvulus</i>	Orchardgrass (seeding)	<i>Dactylis glomerata</i>
Catchfly, English	<i>Silene gallica</i>	Pennycress, field	<i>Thlaspi arvense</i>
Chamomile, mayweed (dogfennel)	<i>Anthemis cotula</i>	Pigweed, redroot	<i>Amaranthus retroflexus</i>
Chickweed, common	<i>Stellaria media</i>	Radish, wild	<i>Raphanus raphanistrum</i>
Fiddleneck, tarweed	<i>Amsinckia lycopsoides</i>	Rocket, London	<i>Sisymbrium irio</i>
Filaree	<i>Erodium sp.</i>	Rocket, common yellow	<i>Barbarea vulgaris</i>
Flkweed	<i>Descurainia Sophia</i>	Salsify	<i>Tragopogon spp.</i>
Groundsel, common	<i>Senecio vulgaris</i>	Shepherdspurse	<i>Capsella bursa-pastoris</i>
Henbit*	<i>Lamium amplexicaule</i>	Speedwell, purslane	<i>Veronica peregrina</i>
Lettuce, Miner's	<i>Montia perfoliata</i>	Spurry, corn	<i>Spergularia arvensis</i>
Mustard, blue	<i>Chorispora tenella</i>		

1 1/3 - 2 Lb/Acre			
Alfalfa* (seeding)	<i>Medicago sativa</i>	Lettuce, prickly*	<i>Lactuca serriola</i>
Barley, foxtail (seeding)	<i>Hordeum jubatum</i>	Malow, common	<i>Malva neglecta</i>
Bluegrass, perennial* (spring only)	<i>Poa spp</i>	Ryegrass, Italian (annual)	<i>Lolium multiflorum</i>
Cockle, white*	<i>Melandrium album</i>	Quackgrass*	<i>Elytrigia repens</i>
Dandelion, common*	<i>Taraxacum officinale</i>	Speedwell, ivyleaf	<i>Veronica hederifolia</i>
Dandelion, false* (spotted catsear)	<i>Hypochaeris radicata</i>	Tea, Mexican*	<i>Chenopodium ambrosioides</i>
Foxtail*	<i>Setaria spp</i>	Thistle, Canada (seeding)	<i>Cirsium arvense</i>
Kochia	<i>Kochia scoparia</i>	Thistle, Russian	<i>Salsola iberica</i>
Lambsquarters, common	<i>Chenopodium album</i>		

* Suppression - a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control. DuPont™ Velpar® DF CU, when applied in late spring or after cutting at the following rates, will control these species listed below:

2/3 - 2 Lb/Acre			
Crabgrass	<i>Digitaria spp</i>	Jimsonweed	<i>Datura stramonium</i>
Feabane	<i>Cyniza spp</i>	Lambsquarters, common	<i>Chenopodium album</i>
Foxtail	<i>Setaria spp.</i>	Pigweed, redroot	<i>Amaranthus retroflexus</i>

SEED ALFALFA (CA, ID, MT, NV, OR, UT, WA)

Velpar® DF CU may be used for general broadleaf weed and grass control in established alfalfa grown for seed.

DORMANT VARIETIES
Make a single application of Velpar® DF CU after alfalfa becomes dormant and before new growth exceeds 2 inches in height in the spring. Where weeds have emerged, use a surfactant.

NON-DORMANT AND SEMI-DORMANT VARIETIES
In the following states, make a single application of Velpar® DF CU during the winter months when alfalfa plants are in the least active stage of growth.

WEEDS CONTROLLED
Refer to the Alfalfa - Weeds Controlled section for specific use rates and weeds controlled.

USE RESTRICTIONS FOR SEED ALFALFA

- Do not apply within 30 days of harvest (cutting for hay), or feeding of forage or grazing.
- Do not use Velpar® DF CU on fields with sandy loam or loamy sand soils having less than 1% organic matter.
- Do not exceed 2/3 pound per acre on fields with sandy loam or loamy sand soils having 1-2% organic matter.
- Do not exceed 2/3 pound per acre on seed alfalfa that has been established for only one growing season.

SEED ALFALFA - WALLA WALLA COUNTY, WASHINGTON

Velpar® DF CU herbicide may be used for the suppression of prickly lettuce and quackgrass and control of Canada thistle (seeding), kochia, and certain other weeds in established alfalfa grown for seed.

Use Rates: 1 1/3 to 2 pounds per acre

Kochia	<i>Kochia scoparia</i>		
Lettuce, prickly*	<i>Lactuca serriola</i>		
Quackgrass*	<i>Elytrigia repens</i>		
Thistle, Canada (seeding)	<i>Cirsium arvense</i>		

* Suppression
USE RESTRICTIONS FOR SEED ALFALFA - WALLA WALLA COUNTY WASHINGTON

Do not apply within 30 days of harvest (cutting for hay), or feeding of forage or grazing. Do not exceed 2 pounds DuPont™ Velpar® DF CU herbicide per acre per application. Do not exceed 2 pounds (1.5 pounds active ingredient hexazinone) per acre per year.

SPRAY EQUIPMENT

Apply Velpar® DF CU using a fixed boom power sprayer or aerial equipment. For ground applications apply in a minimum of 20 gallons of spray solution per acre and by air in a minimum of 5 gallons.

CHEMIGATION FOR ALFALFA

Apply this product only through center pivot or linear-move sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. Severe alfalfa injury may result following application after cutting if either the regrowth is more than 2" high or significant stubble is left after alfalfa cutting. If you have questions about calibration, you may contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments when needed.

DORMANT APPLICATIONS
Select the appropriate rate (see **Use Rates** section) for soil texture and organic matter content using 0.25" to 0.75" of sprinkler irrigation as a continuous injection during the application. Best results are obtained when soil is moist at time of application, and when weeds have not germinated or are less than 2" tall or across.

APPLICATION AFTER CUTTING
Apply Velpar® DF CU at 5.3 ounces per acre to stubble after cutting, following hay removal, and before regrowth exceeds 2" in height. Apply Velpar® DF CU using 0.25" to 0.75" of sprinkler irrigation as a continuous injection during the application. Best results are obtained when soil is moist at time of application and when weeds have not germinated or are less than 2" tall or across.

NOTE: Making an application when daily temperatures are forecast to be in the mid-to-high 90 degree range within 3 to 5 days after treatment may increase the potential for crop injury.

SPRINKLER CHEMIGATION
The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

MIXING INSTRUCTIONS

1. Fill the supply tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of Velpar® DF CU and continue agitation until the Velpar® DF CU is fully dispersed, at least 5 minutes.
3. Once the Velpar® DF CU is fully dispersed, maintain agitation and continue filling tank with water. Velpar® DF CU must be thoroughly mixed with water before adding any other material.

4. As the tank is filling, add tank mix partners (if desired). Follow use precautions and directions on the tank mix partner label.
5. After thorough mixing, the agitation system can be stopped to prevent excessive foaming in the tank. Once thoroughly mixed the solution in the supply tank does not require additional agitation unless specified on the companion products label. If foaming occurs in the injection supply tank, a defoaming agent (defoamer) may be added.

6. Apply Velpar® DF CU spray mixture within 48 hours of mixing to avoid product degradation.

USE PRECAUTIONS - CHEMIGATION
• Distributing treated water in an uneven manner can result in crop injury, lack of effectiveness, or over-tolerance pesticide residues in the crop. Therefore, to ensure that the mixture is applied evenly at the labeled rate, use sufficient water, apply the mixture for the proper length of time and ensure sprinkler produces a uniform water pattern.

USE RESTRICTIONS - CHEMIGATION
• Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
• Do not permit run-off during chemigation.

POSTING OF AREAS TO BE TREATED
Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, daycare centers, hospitals, in-patient clinics, nursing homes, or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to all the following requirements:

- Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas.
- The printed side of the sign must face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

- All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words "KEEP OUT", followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word "STOP". Below the symbol shall be the words "PESTICIDE IN IRRIGATION WATER".
- Posting required for chemigation does not replace other posting and reentry requirements for farm worker safety.

REPLANTING (FOLLOWING ALFALFA)

- Do not replant treated areas to any crop except corn, root crops or sugarcane within two years after treatment, as crop injury may result.
- Corn may be planted 12 months after the last treatment in areas of moderate to high rainfall (greater than 20 inches), provided the use rate did not exceed 1 lb per acre.

- Root crops such as potatoes, sugarbeets, radish and carrots may be planted 12 months after last treatment, provided the use rate does not exceed 2/3 lb per acre. Sites with use rates higher than 2/3 lb per acre must not be replanted to any root crop within 2 years after application of DuPont™ Velpar® DF CU, or unacceptable crop injury may result.

In areas where irrigation is needed to produce the crop, the crop rotation intervals listed may need to be extended if the normal irrigation amount is reduced for any reason.

- Sugarcane may be planted any time following treatment.
- In California, do not replant seed alfalfa areas to any crop within two years after treatment, as crop injury may result.

CROP ROTATION

USE RESTRICTIONS - ALFALFA

- Do not apply to snow-covered or frozen ground.
- Crop injury to alfalfa can be influenced by several factors including alfalfa variety, soil conditions, uniformity of application and environmental conditions, etc. If no prior use history for the site or variety, treat only a small area when first using VELPAR® DF CU.
- If abnormally dry conditions exist following application, restrict the first irrigation to no more than 1/2 acre inch of water.
- Temporary yellowing of alfalfa may occur following VELPAR® DF CU applications.
- Treat only stands of alfalfa established for one year or for one growing season (except in California), provided:
 - The alfalfa stand has a well developed tap root structure that is at least 10 inches in length (0.25 inch diameter below the crown) throughout the field and the crop is healthy, vigorous, and not under stress from weather conditions, low fertility, insects or disease damage.
 - In areas with shorter growing seasons, such as, higher elevations, adequate alfalfa tap root growth may not occur and especially when alfalfa is grown together with a cover or nurse crop. If an adequate tap root is not present, delay application of DuPont™ VELPAR® DF CU until the alfalfa has gone through a minimum of two growing seasons.
- In California, fall planted alfalfa may be treated in the following winter months with VELPAR® DF CU at 1/3 to 2/3 pounds per acre (use higher rate for fine textured soils) provided:
 - alfalfa root growth exceeds 6 inches in length
 - vegetative top growth of alfalfa has lateral development of secondary growth
 - alfalfa is healthy and vigorous, not growing under stress from insect, disease, winter injury or other types of stress.
- Injury may result to alfalfa plants that fail to meet these growth criterion listed above.
- Do not use VELPAR® DF CU on seedling alfalfa, alfalfa-grass mixtures, or other mixed stands as injury may result to the seedling alfalfa or companion crop.
- Do not use a surfactant with VELPAR® DF CU when treating non-dormant alfalfa.
- Do not use VELPAR® DF CU on gravelly or rocky soils, exposed subsoils, hardpan, sand, poorly drained soil, or alkali soils.

BLUEBERRY

HIGH BUSH BLUEBERRIES

VELPAR® DF CU is labeled for control of certain herbaceous and woody weeds in established high bush blueberry fields.

APPLICATION INFORMATION

VELPAR® DF CU may be applied to high bush blueberries that have been established for 3 or more years. Apply VELPAR® DF CU in the spring before the lower leaves of the blueberry plant have fully expanded. Avoid contact of the leaves with the spray solution. Using calibrated ground spray equipment, make the application in sufficient water to provide thorough and uniform coverage to the treated area (usually 20 gallons per acre). Shut off spray booms when starting, turning, slowing or stopping, or injury to the crop may result.

USE PRECAUTIONS FOR HIGH BUSH BLUEBERRIES

- Application to blueberry foliage will result in crop injury.
- Since the effect of VELPAR® DF CU on blueberries varies with soil type, plant vigor, uniformity of applications and amount of rainfall, it is suggested that growers limit their first use to small areas.

USE RESTRICTIONS FOR HIGH BUSH BLUEBERRIES

- Do not apply through any type of irrigation system.
- Do not apply within 90 days of harvest.
- Do not apply to flooded field with standing water.

USE RATES (Lbs/Acre)

Soil texture	less than or equal to 3% organic matter	greater than 3% organic matter
Coarse loamy sand, sandy loam (50-85% sand)	1.3	1.6
Medium loam, silt loam, silt, clay loam, sandy clay loam		2.6
Fine silty clay loam, clay loam, sandy clay, silty clay, clay	1.3 - 2*	2.6

*Use the higher rate as the soil organic matter approaches 3%.

LOW BUSH BLUEBERRIES

DuPont™ VELPAR® DF CU may be used for the control of certain weeds in low bush blueberries.

APPLICATION INFORMATION

VELPAR® DF CU may only be applied to pruned blueberry fields in the spring before leaf emergence. Using calibrated ground spray equipment, make the application in sufficient water to provide thorough and uniform coverage to the treated area (usually 20 gallons per acre). Shut off spray booms when starting, turning, slowing or stopping, or injury to the crop may result.

USE PRECAUTIONS FOR LOWBUSH BLUEBERRIES

- Application to blueberry foliage will result in crop injury.
- Since the effect of VELPAR® DF CU on blueberries varies with soil type, plant vigor, uniformity of applications and amount of rainfall, it is suggested that growers limit their first use to small areas. If excessive leaf drop is observed after treatment, reduce rate in future applications.
- Maintain a 50 foot buffer from any well head or water reservoir.

USE RESTRICTIONS FOR LOWBUSH BLUEBERRIES

- Do not apply through any type of irrigation system.
- Do not apply to flooded field with standing water.
- Do not apply within 450 days of harvest.
- Do not exceed 2.4 pounds per acre if field has been treated with hexazinone within the past 8 years.

Soil texture	less than or equal to 3% organic matter	greater than 3% organic matter
Coarse loamy sand, sandy loam (50-85% sand)	1.2	1.6
Medium loam, silt loam, silt, clay loam, sandy clay loam		2
Fine silty clay loam, clay loam, sandy clay, silty clay, clay	1.2 - 2.4*	2.4 - 3.6**

*Use the higher rate as the soil organic matter approaches 3%.

**Use the higher rate for harder to control species.

IMPREGNATION ON DRY BULK FERTILIZER

Dry bulk fertilizer may be impregnated or coated with VELPAR® DF CU for application to established high bush or low bush blueberries. All instructions on this label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating and labeling. If fertilizer materials are excessively dusty, use a suitable additive to reduce dust prior to impregnation, as dusty fertilizer will result in poor distribution during application. The dry fertilizer must be properly impregnated and uniformly applied to the alfalfa to avoid crop injury and/or poor weed control. To impregnate the fertilizer, use a system consisting of a conveyor or closed drum used to blend dry bulk fertilizer. Any commonly used fertilizer can be impregnated with VELPAR® DF CU, except potassium nitrate or sodium nitrate. Do not use VELPAR® DF CU on limestone. Use a minimum of 250 lb dry bulk fertilizer per acre and up to a maximum of 450 lb per acre. To impregnate or coat the dry bulk fertilizer with VELPAR® DF CU, mix 2 2/3 pounds of VELPAR® DF CU with sufficient water to make one gallon of suspension and thoroughly agitate. Direct the nozzles to deliver a fine spray of this suspension toward the fertilizer for thorough coverage while avoiding spray contact with mixing equipment. Uniform impregnation of VELPAR® DF CU to dry bulk fertilizer will vary, and if the absorptivity is not adequate, the use of an absorptive powder may be required to produce a dry, free-flowing mixture. "Microcel E" is the absorbent powder of choice. When another herbicide is used with VELPAR® DF CU, mix and impregnate the fertilizer immediately.

Apply impregnated fertilizer as soon as possible after impregnation for optimum performance.

Select the rate of DuPont™ VELPAR® DF CU to apply per acre from the appropriate section of this label. Then refer to the rate chart below to determine the amount of VELPAR® DF CU that is to be impregnated on a ton of dry bulk fertilizer, based on the amount of fertilizer to be distributed in one acre.

Rate Chart for Impregnating Fertilizer with VELPAR® DF CU

Fertilizer Rate/Acre	VELPAR® DF CU Rate Per Acre	
	2/3 Lbs	1 Lbs
250 lbs	5.3 lbs/ton	8.0 lbs/ton
300 lbs	4.4 lbs/ton	6.6 lbs/ton
350 lbs	3.7 lbs/ton	5.7 lbs/ton
400 lbs	3.3 lbs/ton	5.0 lbs/ton
450 lbs	2.9 lbs/ton	4.4 lbs/ton

For rates other than those listed, use the following formula to calculate the amounts of VELPAR® DF CU to be impregnated per ton of dry fertilizer.

Lbs VELPAR® DF CU Per Acre	X	1 Ton Fertilizer	=	Lbs VELPAR® DF CU per Ton of Fertilizer
----------------------------	---	------------------	---	---

APPLICATION

Uniform application of VELPAR® DF CU-impregnated dry fertilizer is essential for satisfactory weed control. Accurate calibration of the application equipment is essential for uniform distribution to the surface. The customary method of application is to apply 1/2 the labeled rate and overlap 50%. This results in the best distribution pattern.

WEEDS CONTROLLED

DuPont™ VELPAR® DF CU is labeled for the control or suppression of the following species in high and low bush blueberry crops:

Aster, heath*	<i>Aster ericoides</i>	Lettuce, Miner's	<i>Montia perfoliata</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>	Lettuce, prickly*	<i>Lactuca serriola</i>
Blackberry** (briar)	<i>Rubus spp</i>	Mustard, blue	<i>Chorispora tenella</i>
Bluegrass, Kentucky (perennial)*	<i>Poa pratensis</i>	Mustard, Hill Jim (tumble)	<i>Sisymbrium altissimum</i>
Brome, downy (cheatgrass)	<i>Bromus tectorum</i>	Orchardgrass *	<i>Dactylis glomerata</i>
Broomsedge*	<i>Andropogon virginicus</i>	Orchardgrass (seedling)	<i>Dactylis glomerata</i>
Carrot, wild*	<i>Daucus carota</i>	Panicgrass (witchgrass)	<i>Panicum capillare</i>
Catchfly, English	<i>Silene gallica</i>	Panicum, fall	<i>Panicum dichotomiflorum</i>
Chamomile, mayweed	<i>Anthemis cotula</i>	Pearly everlasting	<i>Anaphalis margaritacea</i>
Cherry, wild	<i>Prunus serotia</i>	Pennycress, field	<i>Thlaspi arvense</i>
Chickweed, common	<i>Stellaria media</i>	Pigweed, redroot	<i>Amaranthus retroflexus</i>
Cinquefoil	<i>Potentilla spp</i>	Quackgrass	<i>Agropyron repens</i>
Cockle, white*	<i>Melanium album</i>	Radish, wild	<i>Raphanus raphanistrum</i>
Dandelion, common*	<i>Taraxacum officinale</i>	Ragweed, common	<i>Ambrosia elatior</i>
Dandelion, false* (spotted catsear)	<i>Hypochoeris radicata</i>	Raspberr** (briar)	<i>Rubus spp</i>
Daisy, oxeye	<i>Chrysanthemum leucanthemum</i>	Rocket, London	<i>Sisymbrium irio</i>
Doak, curly*	<i>Rumex crispus</i>	Rocket, common yellow	<i>Barbarea vulgaris</i>
Dogfennel	<i>Eupatorium capillifolium</i>	Ryegrass, Italian (annual)	<i>Lolium multiflorum</i>
Fescue*	<i>Festuca spp</i>	Ryegrass, perennial*	<i>Lolium perenne</i>
Fiddleneck, tarweed	<i>Amsinckia hycopsoides</i>	Salsify	<i>Tragopogon spp</i>
Filaree	<i>Erodium spp</i>	Shepherdspurse	<i>Capsella bursa-pastoris</i>
Fireweed* (willowweed)	<i>Eupilobium angustifolium</i>	Smartweed, Pennsylvania	<i>Polygonum persicivanicum</i>
Flabiate, flax-leaved	<i>Coryza bonariensis</i>	Sorrel, red	<i>Rumex acetosella</i>
Flixede	<i>Descurainia Sophia</i>	Sorrel, sheep	<i>Rumex angiocarpus</i>
Foxtail, yellow	<i>Setaria lutescens</i>	Spurry, corn	<i>Spergula arvensis</i>
Goldenrod	<i>Solidago spp</i>	Strawberry, wild	<i>Fragaria virginiana</i>
Groundsel, common	<i>Senecio vulgaris</i>	Tansymustard (pinnaet)	<i>Descurainia pinnata</i>
Hawkweed	<i>Hieracium spp</i>	Tea, Mexican*	<i>Chenopodium ambrosioides</i>
Horseweed/marestail	<i>Coryza canadensis</i>	Velvetgrass	<i>Holcus lanatus</i>
Jimsonweed	<i>Datura stramonium</i>	Yarrow	<i>Achillea spp</i>
Lambsquarters, common	<i>Chenopodium album</i>		

2.4 to 3.6 Lbs/acre

Dogbane**	<i>Apocynum spp</i>	Laurel, sheep	<i>Kalmia angustifolia</i>
Meadow-sweet	<i>Filipendula ulmaria</i>	Rose, wild**	<i>Rosa spp</i>
Blackberry, trailing	<i>Rubus ursinus</i>		

- Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.
- ** Harder to control species.

PINEAPPLE

DuPont™ VELPAR® DF CU is labeled for control of certain weeds in pineapple.

APPLICATION INFORMATION

Mix the proper amount of VELPAR® DF CU in water. Add a surfactant at the rate of 0.25% VV.

Use the lower rates on coarse-textured soils or in areas where rainfall exceeds 65 inches per year. Use the higher rates on fine-textured soils or in areas where rainfall is less than 65 inches per year.

Intercrop period - Apply VELPAR® DF CU as a broadcast spray in 100–400 gallons of water per acre at the rate of 1/3 - 2 1/3 pounds per acre. For aerial application, use at least 10 gal water per acre.

Post mulch, preplant - Apply VELPAR® DF CU as a broadcast spray in 100–400 gallons of water per acre at the rate of 1/3 - 2 1/3 pounds per acre.

Post plant, before planted cuttings start active growth - Apply VELPAR® DF CU as a broadcast spray 3–10 months after planting in 100–400 gallons of water per acre at the rate of 1/3 - 2 1/3 pounds per acre. When weed growth has escaped control by other herbicide applications, a post-planting application may be made after the planted cuttings start to grow.

Prior to forcing first ratoon - Apply VELPAR® DF CU as a broadcast spray in 100–400 gallons of water per acre at the rate of 1/3 - 2 1/3 pounds per acre.

Directed postemergence (appleand weeds) inter-space application - Apply VELPAR® DF CU as a directed spray 3–10 months after planting in 50–200 gallons of water per acre (broadcast basis) at the rate of 1/3 - 2 1/3 pounds per acre (broadcast basis) using a stroller boom or knapsack.

Directed spot treatments for perennial grasses before floral induction - Spray perennial grasses postemergence to weed (50–200 gallons per acre depending on size) with 1 1/3 - 2 1/3 pounds per 100 gallons of water as a spot treatment.

Treatments to field edges and roadsides - Apply VELPAR® DF CU at 2 1/3 - 4 8/10 pounds per acre in 100–400 gallons of water.

WEEDS CONTROLLED

VELPAR® DF CU is labeled for the control or suppression of the following species in pineapple crops:

Ageratum, tropic	<i>Ageratum conyzoides</i>	Kao haele*	<i>Leucaena glauca</i>
Balsamapple	<i>Momordica charantia</i>	Moana loa vine*	<i>Canavalia cathartica</i>
Castorbean	<i>Ricinus communis</i>	Morningglory	<i>Ipomoea spp</i>
Crabgrass	<i>Digitaria spp</i>	Oxalis	<i>Oxalis spp</i>
Crotalaria spp	<i>Crotalaria spp</i>	Papoyo	<i>Solanum sandwicense</i>
Dallisgrass	<i>Paspalum dilatatum</i>	Richardsonium	<i>Richardsonia spp</i>
Guineagrass	<i>Panicum maximum</i>	Vaseygrass	<i>Paspalum urvillei</i>
Junglerice	<i>Echinochloa colonum</i>		

- Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

USE RESTRICTIONS - PINEAPPLE

- Do not exceed 4.8 lb VELPAR® DF CU per acre per crop.
- Do not apply VELPAR® DF CU within 181 days of harvest.

SUGARCANE

VELPAR® DF CU is labeled for selective weed control in sugarcane except in the State of Florida.

APPLICATION INFORMATION

Apply a single treatment of VELPAR® DF CU per year using a fixed-boom sprayer and a minimum of 25 gallons per acre unless otherwise directed.

HAWAII

Apply DuPont™ VELPAR® DF CU pre- or postemergence at the following rates for the indicated soil texture:

Soils	VELPAR® DF CU (Lb/Acre) (Plus surfactant 0.25% by volume)
Coarse Texture	
Sand, loamy sand, sandy loam	2/3 - 1 2/10
Medium Texture	
Loam, silt loam, silty clay loam	2/3 - 2 1/3
Fine Texture	
Clay, gray hydromorphic clay	2 1/3 - 4 8/10

Use the higher levels of the labeled rate ranges on soils high in organic matter. Do not apply more than twice the highest labeled rate for the indicated soil texture per crop (18–24 months).

Add an adjuvant for all uses.

For preemergence use only, VELPAR® DF CU may be applied with aerial equipment using at least 10 gallons of spray per acre.

Apply VELPAR® DF CU herbicide as a spot spray application for emerged weeds in sugarcane. Mix 1 to 4 pounds of VELPAR® DF CU per 100 gallons of water. Apply a sufficient volume of spray solution to thoroughly wet weed foliage but do not exceed a use rate of 4.8 pounds per acre. Use the lower concentrations on coarse-textured soils that are low in organic matter, and use the higher concentrations on fine-textured soils that are high in organic matter.

LOUISIANA

Apply 2/3 - 1 2/10 pound of VELPAR® DF CU per acre broadcast in the fall before sugarcane emerges or in the spring before active cane tillering begins. Fall treatments of 2/3 - 1 2/10 pound per acre may be followed by a spring treatment of 2/3 - 1 2/10 pound per acre. Do not apply more than 2 pound per year. Use the lower rates on coarse textured soils and the higher rates on fine textured soils.

PUERTO RICO

For preemergence treatments, apply 1/3 - 2/3 pound of VELPAR® DF CU per acre.

For postemergence treatments, apply 1/3 - 2/3 pound of VELPAR® DF CU per acre to weeds after they have emerged. Use the lower rates on coarse-textured soils and the higher rates on fine-textured soils (high in clay or organic matter). Each raton may receive up to 2/3 pound of VELPAR® DF CU per acre.

For spot treatment of emerged weeds, VELPAR® DF CU may be applied with a knapsack sprayer in concentrations of 1/3 - 2/3 pound per 100 gallons of water. Apply a sufficient spray volume to wet the weed foliage. Do not exceed 100 gallons of spray per treated acre. Use the lower concentration on coarse-textured soils and the higher concentration on fine-textured soils.

Note: Since it is difficult to calibrate "spot" knapsack applications, extra care must be taken not to exceed the rate equivalent of the maximum of 2/3 pound VELPAR® DF CU per acre.

Do not apply more than 1 1/3 pound of VELPAR® DF CU per acre per crop season.

TEXAS

Apply 2/3 - 2 1/3 pound of VELPAR® DF CU per acre. On plant cane, apply the herbicide before the cane emerges or as a directed layby treatment. On stubble cane, apply VELPAR® DF CU preemergence to early postemergence (up to the 3-leaf stage) or as a directed layby treatment. A pre- or early postemergence treatment may be followed by a layby treatment, provided at least 60 days have elapsed and 3 inches of rainfall or sprinkler irrigation have occurred since the first treatment.

Do not apply more than 2 1/3 pound of VELPAR® DF CU per acre per season.

Use the following rates according to the different soil textures:

Soils	Preemergence	VELPAR® DF CU (Lb/Acre)	Layby
Coarse Texture*			
Sandy loam	1/3		1/3
Medium texture			
Loam, silt loam	9/10		9/10
Fine texture			
Clay loam	1 1/3		1 1/3

* With at least 2% organic matter

On dormant cane, a surfactant may be added to the spray mixture to increase control of emerged weeds.

WEEDS CONTROLLED

DuPont™ VELPAR® DF CU is labeled for the control or suppression of the following species in sugarcane crops:

Ageratum, tropic*	<i>Ageratum conyzoides</i>	Millet, Texas	<i>Panicum texanum</i>
Alexandergrass	<i>Brachiaria plantaginea</i>	Morningglory, hairy	<i>Ipomoea pentaphylla</i>
Balsamapple	<i>Momordica charantia</i>	Morningglory, threeleaf	<i>Lonoclea triloba</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>	Mustard, wild	<i>Sinapis arvensis</i>
Bernudgrass*	<i>Cynodon dactylon</i>	Oxalis	<i>Oxalis spp</i>
Burnweed, American (fireweed)	<i>Erechtites hieracifolius</i>	Paintbrush, Flora's	<i>Emilia sonchifolia</i>
Chickweed, common	<i>Stellaria media</i>	Panicum, browntop	<i>Panicum fasciculatum</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>	Paspalum, ricegrass	<i>Paspalum orbiculare</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>	Paspalum, sour	<i>Paspalum conjugatum</i>
Crotalaria, fuzzy	<i>Crotalaria incana</i>	Pigweed, redroot	<i>Amaranthus retroflexus</i>
Crotalaria, showy	<i>Crotalaria spectabilis</i>	Pigweed, slender (green)	<i>Amaranthus viridis</i>
Cuphea, tarweed	<i>Cuphea carthagenensis</i>	Pigweed, smooth	<i>Amaranthus chlorostachys</i>
Dallisgrass	<i>Paspalum dilatatum</i>	Popolo	<i>Solanum sandwicense</i>
Fingergrass, radiate	<i>Chloris radiata</i>	Purslane, common	<i>Portulaca oleracea</i>
Fingergrass, swollen	<i>Chloris barbata</i>	Sandbur	<i>Cenchrus spp</i>
Foxtail, bristly	<i>Setaria verticillata</i>	Sensitive plant (hila hila)	<i>Mimosa spp</i>
Foxtail, yellow	<i>Setaria lutescens</i>	Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Geranium, Carolina	<i>Geranium carolinianum</i>	Southistle, common	<i>Sonchus oleraceus</i>
Goosegrass	<i>Elysiue indica</i>	Spanishneedles	<i>Bidens bipinnata</i>
Guineagrass	<i>Panicum maximum</i>	Sprangletop	<i>Leptochloa spp</i>
Henbit*	<i>Lamium amplexicaule</i>	Spurge, prostrate	<i>Euphorbia humistrata</i>
Itchgrass*	<i>Rottboellia cochinchinensis</i>	Spurge, graceful	<i>Chamaesyce hypericifolia</i>
Job's-tears	<i>Cox lacynia</i>	Sunflower	<i>Helianthus spp</i>
Johnsongrass (seedling)	<i>Sorghum halepense</i>	Vaseygrass	<i>Paspalum urvillei</i>
Junglerice	<i>Echinochloa colonum</i>	Waltheria (hia loa)	<i>Waltheria spp</i>
Lambsquarters, common	<i>Chenopodium album</i>		

- Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

USE RESTRICTIONS - SUGARCANE

- Do not plant any crop other than sugarcane following an application of VELPAR® DF CU.
- Do not plant sugarcane forage livestock.
- Do not apply VELPAR® DF CU:
 - Within 180 days of harvest in Hawaii.
 - Within 234 days of harvest in Louisiana.
 - Within 288 days of harvest in Puerto Rico.
 - Within 234 days of harvest in Texas.
- To avoid injury to sugarcane, observe the following:
 - Do not use VELPAR® DF CU on cane that shows poor vigor because of insect damage, disease, or winter injury, or shows symptoms of other stress conditions such as drought stress.
 - Do not use a surfactant in applications unless otherwise specified.
- Do not use VELPAR® DF CU on gravelly or rocky soils, thinly covered subsoils, or coarse-textured soils (sands to sandy loams) with less than 1% organic matter.
- Temporary chlorosis of the crop may result from application over emerged cane. Applications during active cane growth must be directed to cover the weeds and soil while minimizing crop contact.
- Do not use DuPont™ VELPAR® DF CU on varieties known to be susceptible to herbicides.
- Extremely heavy rainfall after application may result in poor weed control and/or crop injury, especially if the application is made to dry soil.

ADDITIONAL INSTRUCTIONS, PRECAUTIONS, AND RESTRICTIONS FOR AGRICULTURAL USES

SPRAY TANK CLEAN OUT

Thoroughly clean all traces of VELPAR® DF CU from application equipment immediately after use. Flush the tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens