



LICENSED

PERIOD 2015-2017 LIC. NO.

9203.420



LICENSED

PERIOD 2012-2014 LIC. NO.

53226

Net Contents: 50 U.S. gal/189.3 L

# OCTAVE<sup>®</sup> FS

## Sanitizer and Disinfectant

Acid Liquid Sanitizer for Food Processing Equipment in Dairies, Dairy Farms, Breweries, Wineries, Beverage and Food Processing Plants  
Disinfectant for Farms, Livestock Quarters, Poultry Premises, Poultry Hatcheries, Veterinary Clinics, Animal Life Science Laboratories, Animal Care Facilities and Industrial Facilities

**This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.**

### ACTIVE INGREDIENTS:

Hydrogen Peroxide.....	7.52%
Peroxyoctanoic Acid.....	1.00%
Octanoic Acid.....	2.85%
OTHER INGREDIENTS:.....	88.63%
TOTAL:.....	100.00%

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER. CORROSIVE.** Causes irreversible eye damage and skin burns. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield or safety glasses), coveralls worn over long-sleeved shirt and long pants, socks, chemical-resistant gloves, and chemical resistant footwear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash clothing before reuse. Prolonged or frequently repeated skin contact may cause an allergic reaction in some individuals.

**FOR COMMERCIAL USE  
STRONG OXIDIZING AGENT**

### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or going for treatment.

**FOR EMERGENCY MEDICAL INFORMATION IN USA OR CANADA, CALL: 1-800-328-0026.**

**FOR EMERGENCY MEDICAL INFORMATION WORLDWIDE, CALL: 1-651-222-5352 (IN THE USA).**

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

**PHYSICAL AND CHEMICAL HAZARDS:** Strong oxidizing agent. Corrosive. Do not use in concentrated form. Mix only with water according to label instructions. Never bring concentrate in contact with other sanitizers, cleaners or organic substances. This product is not to be used around electrical equipment.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to birds, fish, and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

### STORAGE & DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

**PESTICIDE STORAGE:** Product should be kept cool and in a vented container to avoid any explosion hazard.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill or by incineration.

EPA Reg. No. 1677-209

EPA Est. 1677-MN-1 (P), 61056-IL-1 (SI), 1677-CA-2 (R), 1677-TX-1 (D), 1677-IL-2 (J), 1677-GA-1 (M), 1677-CA-1 (S), 1677-WV-1 (V).

Superscript refers to first letter of date code.



# OCTAVE™ FS

# 53226

## Net Contents: 50 U.S. gal/189.3 L

See front label for Precautionary Statements, First Aid instructions, and Storage & Disposal.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### SANITIZATION

*Octave FS* acid sanitizer is recommended for use on pre-cleaned surfaces such as equipment, pipelines, tanks, vats, fillers, evaporators, pasteurizers and aseptic equipment in dairies, dairy farms, breweries, wineries, beverage and food processing plants. This product is effective as a sanitizer when solution is prepared in water of up to 500 ppm hardness as CaCO<sub>3</sub>.

**NOTE:** FOR MECHANICAL OPERATIONS, prepared use solution may not be reused for sanitizing but may be reused for other purposes such as cleaning.

FOR MANUAL OPERATIONS, fresh sanitizing solution should be prepared at least daily or more often if the solution becomes diluted or soiled.

### SANITIZING FOOD CONTACT SURFACES

Prior to sanitizing, remove gross food particles, and then wash with a detergent solution, followed by a potable water rinse. Sanitize with a concentration of 1 oz *Octave FS* concentrate per 6 to 8 gallons of water (0.1% - 0.13% v/v concentration). At this dilution *Octave FS* is an effective food contact surface sanitizer against the following organisms: *Staphylococcus aureus*, *Escherichia coli*, *Escherichia coli O157:H7*, *Pseudomonas aeruginosa*, *Listeria monocytogenes*, *Salmonella typhimurium*, *Enterobacter sakazakii*, *Campylobacter jejuni* and *Candida albicans*. At the same dilution, *Octave FS* is also effective for sanitization against the common spoilage organisms *Pediococcus damnosus*, *Lactobacillus malefermentans* and *Geotrichum candidum*. Use immersion, coarse spray or circulation techniques as appropriate to the equipment. All surfaces should be exposed to the sanitizing solution for a period of not less than one minute unless a longer time is specified by the governing sanitary code. Drain thoroughly and allow to air dry. Do not rinse.

### FOAM SANITIZING FOOD CONTACT SURFACES

*Octave FS* is an effective foam sanitizer of pre-cleaned food contact surfaces. Foam sanitize with a concentration of 1 oz *Octave FS* concentrate per 6 to 8 gallons of water (0.1% - 0.13% v/v concentration). At this dilution *Octave FS* is an effective foaming food contact surface sanitizer against the following organisms: *Staphylococcus aureus*, *Escherichia coli*, *Escherichia coli O157:H7*, *Listeria monocytogenes*, *Salmonella typhimurium*, and *Enterobacter sakazakii*. At the same dilution, *Octave FS* is also effective for foam sanitization against the common spoilage organisms *Pediococcus damnosus* and *Lactobacillus malefermentans*. Use an appropriate device to generate foam intended to cover the food contact surfaces. Contact your Ecolab representative for information on a recommended foamer. All surfaces should be exposed to the sanitizing solution for a period of not less than one minute unless a longer time is specified by the governing sanitary code. Drain thoroughly and allow to air dry. Do not rinse.

### CONTINUOUS TREATMENT OF CONVEYORS

Wash, rinse and sanitize conveyor equipment. During processing, apply solution containing 1 oz *Octave FS* concentrate per 6 to 8 gallons of water (0.1% - 0.13% v/v concentration) to a conveyor with *Mikro Master* or other suitable dispensing equipment. At this dilution *Octave FS* is effective against *Staphylococcus aureus*, *Escherichia coli*, *Escherichia coli O157:H7*, *Pseudomonas aeruginosa*, *Listeria monocytogenes*, *Salmonella typhimurium*, *Enterobacter sakazakii*, *Campylobacter jejuni* and *Candida albicans*. At the same dilution *Octave FS* is effective at reducing the common spoilage microorganisms *Pediococcus damnosus*, *Lactobacillus malefermentans* and *Geotrichum candidum*. Controlled volumes of *Octave FS* are applied to return portion of conveyor through nozzles so located as to permit maximum drainage of *Octave FS* from equipment and to prevent puddles on top of belt. During interruptions in operations, coarse spray the processing equipment with *Octave FS* solution at not more than 0.13% v/v concentration. Conveyor equipment must be free of product when applying coarse spray. Conveyor surface should be exposed to the sanitizing solution for a period of not less than one minute.

### SANITIZING NON-FOOD CONTACT SURFACES

Pre-clean surfaces as directed above. Sanitize non-food contact surfaces such as floors, walls, tables, chairs, benches, troughs, and drip pans with 1 to 5 oz *Octave FS* concentrate per 8 gallons of water (0.1% - 0.5% v/v concentration). At this concentration the product is effective as a non-food contact surface sanitizer against the following organisms: *Staphylococcus aureus*, *Enterobacter aerogenes*, *Listeria monocytogenes*, *Salmonella typhimurium*, *Pseudomonas aeruginosa*, *Enterobacter sakazakii* and *Geotrichum candidum*. All surfaces should be exposed to the sanitizing solution for a period of not less than 5 minutes. Drain thoroughly and allow to air dry. No rinse necessary.

### FOAM SANITIZING NON-FOOD CONTACT SURFACES

*Octave FS* is an effective foam sanitizer of pre-cleaned non-food contact surfaces, such as boots, floors, walls and associated equipment. Sanitize with a concentration of 1 to 5 oz *Octave FS* concentrate per 8 gallons of water (0.1% - 0.5% v/v concentration). At this concentration the product is effective as a non-food contact surface sanitizer against *Staphylococcus aureus*, *Enterobacter aerogenes*, *Listeria monocytogenes*, *Salmonella typhimurium*, *Enterobacter sakazakii* and *Campylobacter jejuni*. Use an appropriate device to generate foam intended to cover non-food contact surfaces. Wet surfaces thoroughly. Surfaces should be exposed to the sanitizing foam for a period of not less than 5 minutes. No rinse is necessary. Contact your Ecolab representative for information on a recommended foamer.

### FOAM SANITIZING NON-FOOD CONTACT SURFACES

*Octave FS* is an effective foam sanitizer of pre-cleaned non-food contact surfaces, such as boots, floors, walls and associated equipment. To foam sanitize, prepare a solution of 1 to 5 oz *Octave FS* concentrate per 8 gallons of water (0.1% - 0.5% v/v concentration) and 1.4 oz *Liquid K* per 8 gallons (0.13% v/v concentration). *Liquid K* is the only approved foam generator. At this concentration the product is effective as a non-food contact surface sanitizer against *Staphylococcus aureus*, *Enterobacter aerogenes*, *Listeria monocytogenes*, *Salmonella typhimurium*, *Enterobacter sakazakii* and *Campylobacter jejuni*. Use an appropriate device to generate foam intended to cover non-food contact surfaces. Wet surfaces thoroughly. Surfaces should be exposed to the sanitizing foam for a period of not less than 5 minutes. No rinse is necessary. Contact your Ecolab representative for information on a recommended foamer.

### ENTRYWAY FOAM SANITIZING SYSTEMS:

To prevent cross contamination from area to area, set the system to deliver sanitizing solution at 1 oz *Octave FS* per 1 to 8 gallons of water (0.1% - 0.78% v/v).

Alternatively, *Octave FS* can be used along with *Liquid K* to apply the foam by setting the system to deliver 1 oz *Octave FS* per 1 to 8 gallons of water (0.1% - 0.78% v/v) and 1.4 oz *Liquid K* per 8 gallons of water (0.13% v/v concentration).

The foam (or spray) should cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor.

### SHOE BATH SANITIZER DIRECTIONS:

To prevent tracking into animal areas and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made solution should be placed at all entrances to buildings and hatcheries. Scrape waterproof work boots (shoes) and place in solution of 1 oz *Octave FS* per 6 to 8 gallons of water (0.1% - 0.13% v/v) for one minute prior to entering area. (If there is a heavy soil load or excessive traffic place work boots (shoes) in a use-solution of 1 oz per 1 to 8 gallons of water (0.1 - 0.78% v/v) (or equivalent dilution) for one minute prior to entering area.) Change the solution in the bath daily or more often if solution appears diluted or soiled.

### SHOE FOAM SANITIZER DIRECTIONS:

*Octave FS* can be used to prevent tracking into processing, packaging and storage areas of food and beverage plants. Apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 1 oz per 1 to 8 gallons of water (0.1% - 0.78% v/v) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacturer of the foam generator/aerator. Scrape waterproof shoes. Stand and/or walk through foamed area for one minute prior to entering area. Foam area should be washed and replaced daily or when it appears dirty.

### DISINFECTION

*Octave FS* disinfects as it cleans in one operation. *Octave FS* can be used to disinfect floors, walls and other hard nonporous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, shelves, racks, carts, exterior surfaces of refrigerators and coolers, tile, linoleum, vinyl, glazed porcelain, plastic (such as polypropylene and polyethylene), stainless steel, or glass. Areas of use include dairies, dairy farms, breweries, wineries, beverage and food processing plants, poultry farms, veterinary facilities, animal care facilities and industrial facilities.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging material must be removed from the room or carefully protected.

### COMBINATION DISINFECTION AND CLEANING

*Octave FS* is effective as a disinfectant at a concentration of 1 - 2 oz *Octave FS* concentrate per 2 gallons (0.39% - 0.78% v/v concentration) of hard water (500 ppm as CaCO<sub>3</sub>), in the presence of 5% blood serum. At this dilution, *Octave FS* is effective against *Staphylococcus aureus*, *Salmonella enterica*, *Pseudomonas aeruginosa* and *Candida albicans*. For heavily soiled areas a pre-cleaning step is required. Prepare a disinfecting and cleaning solution by diluting 1 - 2 oz *Octave FS* concentrate per 2 gallons of water (0.39% - 0.78% v/v concentration). Apply use solution with mop, cloth, sponge, brush, scrubber, or coarse spray device or by soaking or foaming so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, and then remove solution and entrapped soil with a clean wet mop, cloth, or wet vacuum pickup. Prepare a fresh solution daily or when it becomes soiled or diluted.

### FOAM DISINFECTION

*Octave FS* is an effective foam disinfectant and cleaner on hard, non-porous surfaces, such as boots, floors, walls, drains, and associated equipment. For this application, prepare a solution of 1 - 2 oz *Octave FS* concentrate per 2 gallons of water (0.39% - 0.78% v/v concentration). Apply solution as a foam using recommended equipment. Wet surfaces thoroughly. At this concentration, the product is effective against the following: *Staphylococcus aureus*, *Salmonella enterica*, and *Pseudomonas aeruginosa*. Surfaces should be exposed to the disinfecting foam for a period of not less than 10 minutes. No rinse is necessary. Contact your Ecolab representative for information on a recommended foamer.

### VRUCIDAL

*Octave FS* is an effective virucide at a concentration of 1 - 2 oz per 2 gallons (0.39%-0.78% v/v concentration) of hard water (500 ppm as CaCO<sub>3</sub>), in the presence of 5% blood serum. At this dilution, *Octave FS* is effective against Influenza A (H1N1), Influenza A (H3N2), Influenza B, Reovirus, Influenza A, Norovirus, Feline calicivirus and Avian Influenza when used at 20°C with a 10 minute contact time. Apply as directed under Disinfection.

### DIRECTIONS FOR FOGGING

To sanitize hard surfaces as an adjunct to acceptable manual cleaning and disinfecting of room surfaces: Prior to fogging, food products and packaging materials must be removed from the room or carefully protected. Fog desired areas using one quart of a 0.39% - 0.78% v/v (3900 - 7800 ppm product) *Octave FS* solution (1 - 2 oz. per 2 gallons of water) per 1000 cu. ft. of room volume. Vacate the area of all personnel during fogging and until the hydrogen peroxide air concentration is below 0.5 ppm. Allow surfaces to drain thoroughly before operations are resumed. Solutions above 0.5% may be corrosive and are not to be used on all surfaces. Test solutions on surfaces prior to use. All hard non-porous food contact surfaces treated with the fog must be rinsed thoroughly with a potable water rinse.

### DISINFECTION OF POULTRY PREMISES, TRUCKS, COOPS AND CRATES

1. Remove all poultry and feed from premises, trucks, coops and crates.
2. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with a detergent and rinse with water.
5. Saturate surfaces with a 1 - 2 oz per 2 gallons (0.39%-0.78% v/v concentration) solution of *Octave FS* for a period of 10 minutes.
6. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
7. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with a detergent and rinse with potable water before reuse.

See your Ecolab Representative for specific recommendations for all cleaning and rinsing requirements.

### POULTRY HATCHERY DISINFECTION

Clean out any remaining eggs and chicks. Remove gross soils, such as litter, down, shell fragments or other hatching related debris. Empty all racks and other equipment. Thoroughly wash all surfaces, including floors, walls, conveyors, trays and water systems with a recommended detergent. Rinse thoroughly with water. Apply a 1 - 2 oz per 2 gallons (0.39%-0.78% v/v concentration) solution of *Octave FS* with a mop, cloth, brush, foam or coarse spray. Wet all surfaces and allow to remain wet for 10 minutes. Ventilate buildings and other closed spaces. Allow to dry before reintroducing eggs.

### DISINFECTION AND DEODORIZING OF ANIMAL HOUSING FACILITIES (BARN, KENNELS, HUTCHES)

Remove animals and feed from facilities. Remove litter, waste matter and gross soils from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, rack and other feeding and watering equipment. Wash surfaces with a recommended alkaline detergent, by manual, foam, or spray application. Rinse with water. Apply a 1 - 2 oz per 2 gallons (0.39%-0.78% v/v concentration) solution of *Octave FS* with a mop, cloth, brush, foam or coarse spray. Wet all surfaces and allow to remain wet for 10 minutes. Ventilate buildings and other closed spaces. Allow to air dry before reintroducing animals.

### FARM PREMISE DISINFECTION DIRECTIONS:

1. Remove all animals and feed from premises, vehicles, and enclosures.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean surfaces with soap or detergent by manual, foam, or spray application and rinse with water.
5. Saturate surfaces with a use-solution of 1 - 2 oz *Octave FS* per 2 gallons of water (0.39% - 0.78% v/v concentration or 3900 - 7800 ppm product), (or equivalent dilution) and allow to remain wet for a period of 10 minutes.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure. Wipe or allow to air dry.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

Ecolab Inc., Food & Beverage Division  
370 Wabasha Street N.  
St. Paul, Minnesota 55102-1390 U.S.A.

© 2007 Ecolab Inc. Made in U.S.A. 712250/5301/0607  
All Rights Reserved