

FABRISTAT

HEALTH	3
FLAMMABILITY	2
REACTIVITY	0

UN2920

Corrosive liquids, flammable, n.o.s.
(Dialkyldimethylammonium chloride)



@TD_L

PROD. No.

LOT No.

Net Weight:



LICENSED

PERIOD 2015-2017 LIC. NO.

9203.502



FABRISTAT

Twin-Chain Quaternary
Ammonium Compound Concentrate
Residual Bacteriostat and Self-Sanitizing Agent
Under Humid or Wet Contamination Conditions

KEEP OUT OF REACH OF CHILDREN
DANGER

Manufactured for:
Ecolab Inc.,
370 Wabasha Street, N.,
St. Paul, MN 55102-1390 USA

HEALTH	3
FLAMMABILITY	2
REACTIVITY	0

Active Ingredients:

Octyl decyl dimethyl ammonium chloride	25.0%
Dioctyl dimethyl ammonium chloride	10.0%
Didecyl dimethyl ammonium chloride	15.0%
Other Ingredients	50.0%
TOTAL:	100.0%

See side panel for Precautionary Statements and First Aid

6200101

PROD. No.

LOT No.

Net Weight:

EPA Est. No:

8-27-14
EPA Reg. No. 6836-63-1677

FABRISTAT is based on a highly active mixture of three Twin Chain quaternary ammonium compounds. Soiled and contaminated fabrics such as diapers, hospital and institutional linen, and athletic apparel is of major housekeeping concern, not only in hospitals, but in institutions, hotels, restaurants and schools. Used as directed, FABRISTAT provides effective residual bacteriostatic and self-sanitizing properties for laundered items such as diapers, hospital and institutional linens, and athletic apparel. FABRISTAT is designed for commercial and institutional laundry use by addition to the final rinse step. For recommended solution concentration, see "Directions for Use." FABRISTAT is economical to use because it is concentrated. It should be handled with care. FABRISTAT is an effective laundry sanitizer when used as directed in waters up to 200 ppm hardness against:

- Corynebacterium ammoniagenes*
- Escherichia coli O157:H7 [E. coli O157:H7]*
- Klebsiella pneumoniae*
- Listeria monocytogenes*
- Salmonella enterica [Salmonella]*
- Staphylococcus aureus [Staph]*
- Staphylococcus aureus – Community Associated Methicillin-Resistant [CA-MRSA]*
- Staphylococcus aureus – Hospital Associated Methicillin resistant [HA-MRSA]*

FABRISTAT provides residual self-sanitizing efficacy in the laundry rinse cycle when used as directed in waters up to 200 ppm hardness against:

- Klebsiella pneumoniae*
- Staphylococcus aureus*

For residual bacteriostatic or self-sanitizing activity, conditions of high relative humidity or wet contamination are required. FABRISTAT provides residual bacteriostatic efficacy in the laundry rinse cycle when used as directed against gram negative and/or gram positive odor causing bacteria.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER. Corrosive: Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses), protective clothing and protective gloves (rubber or chemical resistant). May be fatal if swallowed or inhaled. Do not breathe vapor. Wear a MSHA/NIOSH-approved dust/mist filtering respirator number TC-21C or a NIOSH approved respirator with any N, R, P, or HE filter. Harmful if absorbed through the skin. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

FIRST AID
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

ENVIRONMENTAL HAZARDS
This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

Dilution Chart
Laundry Sanitizer3 oz. per 100 pounds dry fabric(781 ppm active quat)
Residual Bacteriostat0.56 oz. per 100 pounds dry fabric(175 ppm active quat)
Residual Self Sanitization ...0.6 oz. per 100 pounds dry fabric(195 ppm active quat)

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

LAUNDRY SANITIZER
Add 3 oz. of FABRISTAT in high efficiency applications per a maximum of 100 pounds of fabric [dry weight] to sanitize laundered fabrics in the rinse cycle. A minimum cycle time of 5 minutes is required.

RESIDUAL BACTERIOSTATIC
Initial Treatment: Use 0.56 oz. (3.5 teaspoons) of FABRISTAT for a minimum 24 hours holding in commercial or industrial applications per a maximum of 100 pounds of fabric (dry weight). Dilute the appropriate amount of FABRISTAT in one to two gallons of water and then add this solution to the wash wheel at the beginning of the final rinse cycle. A minimum rinse cycle time of 10 minutes is required.
Repeat treatment: Follow Initial Treatment directions and re-treat fabric after each washing or if odor persists.

RESIDUAL SELF-SANITIZING
Use 0.6 oz. of FABRISTAT in commercial or industrial applications per a maximum of 100 pounds of fabric (dry weight). Dilute the appropriate amount of FABRISTAT in one to two gallons of water and then add this solution to the wash wheel at the beginning of the final rinse cycle. A minimum rinse cycle time of 8 minutes is required.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage: Open dumping is prohibited. Store in original container in areas inaccessible to children.
Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
Container Disposal: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying.
(Plastic and Metal Containers): Triple rinse as follows: Fill container ¼ full with water and reclose the container. Agitate vigorously, and dispose of rinsate consistent with pesticide disposal instructions. Repeat two more times. Then offer for recycling if available or puncture and dispose in sanitary landfill or by other procedures approved by state and local authorities. Follow pesticide disposal instructions for rinsate. If not triple rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state, and federal regulations. **(For metal containers only):** DO NOT cut or weld metal containers. **(For Bag in Box Containers):** Completely empty bag into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities