DISC.

12/16/2004

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO. 9205.43

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2008-2010 LIC. NO.

**FOR INDOOR USE ONLY**

JT EATON™

# ANSWER® for RATS

FEEDER BOX

# KILLS RATS and Mice

CONVENIENT FEEDER BOX

**REMOVE TAB**

CONVENIENT FEEDER BOX

**REMOVE TAB**

Manufactured in the U.S.A.

**ACTIVE INGREDIENT,**
Chlorophacinone
CAS [3691-35-8] ..... 0.005%
**OTHER INGREDIENTS** ... 99.995%
**TOTAL:** 100.000%

EPA REG. NO. 56-69
EPA EST. NO. 56-OH-1

**KEEP OUT OF REACH OF CHILDREN**
**CAUTION**
SEE BACK FOR ADDITIONAL PRECAUTIONARY STATEMENTS

T.M.

**NET CONTENTS: 4 oz. of bait**

**Item #326**

JT EATON™

# ANSWER® FOR RATS

FEEDER BOX

0 76706 32600 3

JT EATON™ is a trademark and ANSWER® is a registered trademark of J.T. Eaton & Co., Inc. • Mfg. by J.T. Eaton & Co., Inc, Twinsburg, OH 44087 • U.S.A. • Made in U.S.A. © 2004

**ITEM NO. 326**
**FNW32604OJTG**

THIS BOX MADE FROM RECYCLED MATERIAL.

## PRECAUTIONARY STATEMENTS - HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Keep away from humans, domestic animals and pets. Keep away from feed or foodstuff. Avoid contact with skin, eyes or clothing. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of the blood and cause bleeding. Wear gloves when handling this product. Wash thoroughly with soap and water after handling bait.

**ENVIRONMENTAL HAZARD:** This product is toxic to fish and wildlife. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

## FIRST AID

**HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE, INCLUDING 1-800-858-7378**

| | |
|---|---|
| **IF SWALLOWED** | • Call a Poison Control Center, doctor or **1-800-664-9042** or **1-800-858-7378** immediately for treatment advice.<br>• Have person sip a glass of water if able to swallow.<br>• Do not induce vomiting, unless told to by a Poison Control Center or doctor. |
| **IF ON SKIN OR CLOTHING** | • Take off contaminated clothing.<br>• Rinse skin immediately with plenty of water for 15 to 20 minutes.<br>• Call a Poison Control Center, doctor or **1-800-664-9042** or **1-800-858-7378** immediately for treatment advice. |

**NOTE TO PHYSICIAN**: This material may reduce the clotting ability of blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

## STORAGE AND DISPOSAL

**STORAGE:** Store only in original container in a cool, dry place inaccessible to children and pets.
**DISPOSAL: IF FEEDER BOX IS EMPTY:** Do not reuse empty container. Place in trash or offer for recycling if available.
**IF PARTLY FILLED:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place any unused product down any indoor or outdoor drain.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. The efficacy of the product may be reduced under high moisture conditions.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:
1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in ***tamper-resistant bait stations***. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** THIS IS NOT A TAMPER-PROOF OR TAMPER-RESISTANT BAIT STATION. DO NOT USE THIS PRODUCT IN AREAS ACCESSIBLE TO CHILDREN, PETS, DOMESTIC ANIMALS OR NONTARGET WILDLIFE. DO NOT USE THIS PRODUCT OUTDOORS. This product may be used to control Norway Rats (Rattus norvegicus), Roof Rats (Rattus rattus), and House Mice (Mus musculus) in homes and in industrial, commercial or agricultural buildings. Do not place feeder boxes in areas where there is a possibility of contaminating food, water, or surfaces that come in direct contact with either.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, beside burrows, inside buildings, or in corners and concealed places.

**APPLICATION DIRECTIONS:** The amount of bait in one 4 oz. feeder box of this product is unlikely to be sufficient to kill more than 1 or 2 rats.
•**For Rats:** Apply 1 to 4 feeder boxes (4 to 16 oz. of bait) per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.
•**For Mice:** Open feeder box, while wearing gloves, and apply 1/4 to 1/2 ounce of bait at 8- to 12- foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.
•**For Rats and Mice:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and any leftover bait properly.
To prevent reinfestation, limit as many sources of food, water and harborages as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish with fresh bait as often as needed.