# Equi-Spot™

SPOT-ON FLY CONTROL FOR HORSES

**Protects horses from flies, gnats, mosquitoes and ticks**

- Kills & repels house flies, stable flies, face flies, horn flies, eye gnats, and ticks on horses
- Aids in the control of horse flies, deer flies, mosquitoes and black flies on horses

Includes 1 FREE Application!

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No.: 270-279 • EPA Est. No.: 270-NB-003

Manufactured for

FARNAM

Farnam Companies, Inc.

| ACTIVE INGREDIENTS: | |
|---|---|
| *Permethrin: (CAS #52645-53-1) | 45.0% |
| OTHER INGREDIENTS: | 55.0% |
| TOTAL | 100.0% |

*cis/trans ratio: Max 55% (+) cis and min 45% (+) trans

Net Contents: 3 [0.34 fl.oz.(10.0 ml)] Applicators

CM K 369

# Equi-Spot™

SPOT-ON FLY CONTROL FOR HORSES

**READ ENTIRE LABEL BEFORE EACH USE**
**USE ONLY ON HORSES OVER 12 WEEKS OF AGE**

**PRECAUTIONARY STATEMENTS:**
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS:**
**CAUTION**

**Hazards to Humans:** Harmful if swallowed. Avoid contact with eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling.

**Hazards to Animals:** ***FOR EXTERNAL USE ON HORSES ONLY. Do not use on foals less than 12 weeks of age.*** Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, nursing animals or horses. Consult a veterinarian before using this product with known drug interactions. ***DO NOT USE ON HORSES OTHER THAN HORSES.*** Certain individuals may develop mild irritation or other adverse reactions. Consult a veterinarian before using on a horse with any open wounds or cuts.

**FIRST AID:**

**IF SWALLOWED:**
- Call a physician or poison control center immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**IF IN EYES:**
- Hold eyes open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

In case of medical emergencies or health and safety inquiries, or in case of fire, leaking or damaged containers, or for product use information, call (602) 285-1660.

Have the product container or label with you when calling a poison control center or doctor, for treatment.

**Adverse Reactions:** Some animals may be sensitive to ingredients in this product. Reactions in horses may include skin sensitivity. Horses may show lethargy, increased pruritis (itchiness), erythema (redness), rash and hair discoloration or hair loss at the application site. Observe the horse following treatment. Sensitivity may occur after using ANY pesticide product on animals. If signs of sensitivity occur, bathe your horse with a mild, non-insecticidal shampoo and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**PHYSICAL OR CHEMICAL HAZARDS:**
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAMES
**ENVIRONMENTAL HAZARDS:**
This product is extremely toxic to fish. Do not add directly to water. Do not contaminate water when disposing of product or packaging.

**DIRECTIONS FOR USE:**
***It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.***
**HORSES:**
***DO NOT USE ON HORSES INTENDED FOR FOOD USE.***
***DO NOT USE ON FOALS UNDER 12 WEEKS (3 MONTHS) OF AGE.***

Remove product tubes from package. Fold along perforation between tubes and tear off one tube. Return unused tubes to package. Holding tube with notched end pointing up and away from face and body, snap off narrow end at notches.

**Apply one tube (10.0 ml) of product solution as follows:**
- 5 ml along the back line starting at the poll (top of head behind the ears) along the base of the mane, down the dorsal midline to the dock (base of the tail).
- 1 ml to the top of the forehead under the forelock.
- 1 ml as a stripe to the back of each foreleg from the ankle up to the knee (2 ml total).
- 1 ml as a stripe to the back of each hindleg from the ankle up to the hock (2 ml total).
- Do not allow the product to contact horse's eyes or mucous membranes.
- Wrap tube and put in trash.
- Do not reapply for 14 days (2 weeks).



**STORAGE AND DISPOSAL:**
**Do not contaminate water, food or feed by storage or disposal.**

**Storage:** Store in cool, dry place. Protect from freezing.

**Disposal:** If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or call 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**NOTICE OF WARRANTY**

FARNAM COMPANIES, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS, IN ACCORDANCE WITH THE STATEMENTS MADE ON THIS LABEL.


FARNAM COMPANIES, INC.
Horse Products Division
P.O. Box 34820
Phoenix, AZ 85067-4820
www.farnam.com

In case of emergency or for product use information, call 602-285-1660
7 AM - 3:45 PM MST M-F


©2002 Farnam Companies, Inc.

Product Code: 3002606 RM #: 2003486 01-2031

0 86621 02606 3

CM K 369

PMS

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2011-2013 LIC. NO. 9212.343

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2014-2016 LIC. NO.

2002-2004
2005-2007

9212.343

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2008-2010 LIC. NO.