

LICENSED
HAWAII
DEPT. OF AGRICULTURE

Permit No. 2003-2005 9212.360



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2009-2011 LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2012-2014 LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO.

9212.360
Disc.

READ ENTIRE LABEL
BEFORE EACH USE.
USE ONLY ON
DOGS/PUPPIES.

SCRATCHEX
Color-Full

Formula 5 FLEA & TICK COLLAR FOR DOGS

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. Buckle collar around dog's/puppy's neck. Fit collar snug enough to prevent placement of the dog's/puppy's leg or lower jaw through the collar, yet loose enough to be rotated about the neck. Generally, if used on large dogs, two or three fingers should fit between the collar and dog's/puppy's neck. 2. Leave 2-3 extra inches on the collar for adjustment, cut off excess length and dispose of in trash. 3. Check periodically and adjust fit if necessary, especially when puppies are rapidly growing. 4. May be used in addition to regular collar. Replace collar when effectiveness diminishes.

WHAT TO EXPECT:

For Fleas: 1. This product is designed to kill adult fleas. 2. This collar starts killing fleas as soon as it is placed around dog's/puppy's neck. 3. Protection against fleas will be achieved within a few days, providing continuous protection for up to 5 months. 4. Replace collar every 5 months to help prevent flea infestations on your pet and in your home.

For Ticks: 1. Ticks are tougher to kill than fleas and are killed more slowly. 2. Full protection against adult and immature ticks will be achieved within a few days, providing continuous protection for up to 5 months. 3. Adult ticks will be killed over the entire pet. Dead ticks will fall off or may be easily removed. 4. Replace collar every 5 months for continuous protection from ticks. If your dog/puppy goes swimming or is out in the rain, it is not necessary to remove the collar. The continuous action collar will rapidly replace any flea and tick killing material removed.

STORAGE and DISPOSAL: Storage: Store in original unopened container away from children. Disposal: Do not reuse container or used collar. Dispose of in trash.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Do not get dust or collar in mouth. Harmful if swallowed. Do not get dust in eyes. Will cause temporary pupillary constriction. * The dust released by this collar is a cholinesterase inhibitor.

Humans: Do not open package until ready to use. Do not allow children to handle this collar. Dust will form on this collar during storage. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling collar.

Animals: Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. Do not use on puppies under 12 weeks of age. Do not use this product on dogs simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea and tick collars may be immediately replaced. When collar is first worn, observe risk area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water.

FIRST AID

FIRST AID	
Call a poison control center or doctor for treatment advice.	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-234-2269 for emergency medical treatment information.	
NOTE TO PHYSICIAN/VETERINARIAN: *Atropine and *oximastropine are antidotal only if symptoms of cholinesterase inhibition are present.	
Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label.	
Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.	
See top panel of box for details of unconditional guarantee.	

FARNAM
PET PRODUCTS
For healthier, happier pets

Distributed by Farnam Companies, Inc.
Pet Products Division
P.O. Box 34820 • Phoenix, AZ 85067-4820

For information, or in an emergency,
call 1-800-234-2269.
Visit our Web site at www.farnampet.com

EPA Reg. No. 2724-254-210
EPA Est. No. 2724-TX-1
Product Code: 3003151 02-1500
©2003 Farnam Companies, Inc.

GUARANTEED OR YOUR MONEY BACK


If not satisfied, send used collar and original receipt with purchase price circled to Farnam Companies, Inc., Pet Products Division
P.O. Box 34820 • Phoenix, AZ 85067-4820

SCRATCHEX
Color-Full

Formula 5 FLEA & TICK COLLAR

for dogs

Works From Head To Tail to
Kill Fleas
and Lyme Disease-
Carrying Ticks
up to 5 months



Crystal Clear
DO NOT ALLOW CHILDREN
TO PLAY WITH THIS COLLAR
CAUTION: See Precautions on back.

ACTIVE INGREDIENT:
Propoxur (CAS#114-26-1).....10%
OTHER INGREDIENTS.....90%
TOTAL 100%

contents: 1 collar
NET WT. 1 OZ. (28 g)

SCRATCHEX® Color-Full® Formula 5 FLEA & TICK COLLAR FOR DOGS

SCRATCHEX® Color-Full® Formula 5 FLEA & TICK COLLAR FOR DOGS

7171660



SCRATCHEX® Color-Full®
Formula 5
FLEA & TICK COLLAR
FOR DOGS