



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO. **9212.420**

**Disc.**

# ADAMS™ FLEA & TICK COLLAR FOR CATS & KITTENS

PS

ADAMS™ FLEA & TICK COLLAR  
FOR CATS & KITTENS

# ADAMS™

## FLEA & TICK COLLAR

for CATS & KITTENS



Maintenance STEP

- ◆ **Kills fleas & ticks**  
(including those carrying Lyme disease)
- ◆ **Kills flea eggs**  
(to prevent reinfestation)

**ACTIVE INGREDIENTS:**

Propoxur (CAS #114-26-1)..... 10.0%

(S)-Methoprene (CAS #65733-16-6)..... 2.1%

**OTHER INGREDIENTS:**..... 87.9%

**TOTAL**..... 100.0%

**DO NOT ALLOW CHILDREN  
TO PLAY WITH THIS COLLAR**

**CAUTION**

See Precautions on Back

Net Weight: .37 OZ (10.5 g)

**8**  
MONTH FLEA  
PROTECTION




0 39079 06047 9

ADAMS™ FLEA & TICK COLLAR  
FOR CATS & KITTENS

# ADAMS™

## FLEA & TICK COLLAR FOR CATS & KITTENS



**READ ENTIRE LABEL BEFORE EACH USE.  
USE ONLY ON CATS AND KITTENS 12 WEEKS  
OF AGE AND OLDER.**

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**1.** Buckle collar around cat's neck. Fit collar snug enough to prevent placement of the cat's leg or lower jaw through the collar, yet loose enough to be rotated about the neck. **2.** Leave 1-2 extra inches on the length of the collar for adjustment, cut off excess, and dispose of in trash. **3.** Check periodically and adjust fit, if necessary, especially when kittens are rapidly growing. **4.** Replace in 8 months for continuous protection from adult fleas and flea egg hatch. Replace in 6 months for continuous protection from ticks. Collar does not need replacement if cat becomes wet.

To minimize flea infestation, use appropriate products to: Treat the pet; Treat the indoor environment, especially where pets spend time resting; Treat the outdoor environment when necessary.

**STORAGE and DISPOSAL:** Collar should be left in place once fitted on cat. If collar is temporarily removed, do not set it on wood or other surfaces.

**Storage:** Store in original, unopened package away from children. **Disposal:** Do not reuse container or used collar. Dispose of in trash.

### WHAT TO EXPECT

**FOR FLEAS:** **1.** This ADAMS™ Flea & Tick Collar for Cats & Kittens is designed to kill adult fleas and flea eggs. **2.** This collar begins working to kill fleas and flea eggs as soon as it is placed around the cat's neck. **3.** Protection against fleas and flea eggs will be achieved within 48 hours, providing continuous protection for 8 months. **4.** Replace collar every 8 months to help control adult flea infestations on your pet.

**FOR TICKS:** **1.** Full protection against adult and immature ticks will be achieved within a few days, providing continuous protection for up to 6 months. **2.** Adult ticks will be killed over the entire pet. Dead ticks will fall off or may be easily removed. **3.** Replace collar every 6 months for continuous protection from ticks.

**PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION:** Do not get dust or collar in mouth. Harmful if swallowed or absorbed through skin. Do not get dust in eyes, will cause temporary constriction of the pupils.\* The dust released by this collar is a cholinesterase inhibitor. Avoid contact with skin, eyes or clothing.

**HUMANS:** Do not open package until ready to use. Do not allow children to play with this collar. Pesticidal dust will form on this collar during storage. Wash hands thoroughly with soap and water after handling collar. **ANIMALS:** Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing cats. Do not use on kittens under 12 weeks of age. Do not use this product on cats simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting\*\* drugs or pesticides (e.g. carbaryl, etc.). However, flea and tick collars may be immediately replaced. When collar is first worn, observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your cat with mild soap and rinse with large amounts of water.

### FIRST AID

**If swallowed**

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**If on skin or clothing**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call poison control center or doctor for treatment advice.

**If in eyes**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

**IN CASE OF EMERGENCY, CALL 1-800-234-2269 OR A POISON CONTROL CENTER.** Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**NOTE TO PHYSICIAN/VETERINARIAN:** \*Atropine is antidotal only if symptoms of cholinesterase inhibition are present.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

Satisfaction guaranteed or your money back. For product use information, call 1-800-234-2269 or visit our Web site [adamsfleacontrol.com](http://adamsfleacontrol.com).

\*\*For information on cholinesterase inhibition, please call our 800 number or consult your veterinarian.

EPA Reg. No. 2724-491-270  
EPA Est. No. 2724-TX-1



Manufactured for:  
**FARNAM PET PRODUCTS**  
A Division of Farnam Companies, Inc.  
P.O. Box 34820, Phoenix, AZ 85067-4820

Product Code: 3006047 RM#: 107047 06-1880  
©2006 Farnam Companies, Inc.

