STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO. 9212.464

ADAMS™

Flea & Tick Control Collar for Small Dogs

CENTRAL Garden & Pet

READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
Contains an organophosphate that inhibits cholinesterase.	
If swallowed	• Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.
Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-800-234-2269 for emergency medical treatment information.	
Note to Physicians and Veterinarians: This product contains tetrachlorvinphos (an organophosphate), which is a cholinesterase inhibitor. Cholinergic symptoms may include salivation, miosis, incoordination, muscle fasciculation and/or weakness, vomiting and diarrhea. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine. Call 1-800-234-2269 or your local Poison Control Center for further information.	

DIRECTIONS FOR USE:
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REMOVE COLLAR FROM PACKAGE, UNROLL AND STRETCH TO ACTIVATE INSECTICIDE GENERATOR. As the collar begins to work, a fine white powder will appear on surface.

Do not use on puppies under 6 weeks old. Do not unroll collar until ready to use. Place the Adams™ Flea & Tick Control Collar for Small Dogs around the dog's neck, adjust for proper fit, and buckle in place. The collar must be worn loosely to allow for growth of the dog and to permit the collar to move around the neck. A properly fitted collar is one that, when fastened, will snugly slide over the pet's head. Leave 2 or 3 inches on the collar for extra adjustment and cut off and dispose of the extra length.

Consult a veterinarian before using this product on debilitated, aged, or medicated animals. If used on pregnant or nursing dogs, do not replace collar until the puppies are at least 6 weeks old.

Sensitivity may occur after using ANY pesticide product for pets. Some animals may become irritated by any collar if it is applied too tightly. If this occurs, loosen the collar. If irritation continues, remove the collar from the animal. If signs of sensitivity occur, remove the collar and bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or similar chemicals. However, flea and tick collars may be immediately replaced with new collars.

This collar is intended for use only as an insecticide/IGR generator.

The collar will begin to kill fleas, ticks and flea eggs and larvae immediately. Its full protection against harder to kill ticks will be built up within a few days after being placed on the pet. For continuous flea and tick protection under normal conditions replace the collar every 7 months. Under conditions where pets are exposed to severe flea and tick infestations, it may be necessary to replace the collar more frequently.

Wetting will not impair the collar's effectiveness or the pet's protection. If the pet goes swimming or is out in the rain it is not necessary to remove the collar. The Adams™ Flea & Tick Control Collar for Small Dogs may be worn with a regular collar.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage and disposal.
Pesticide Storage: Store this product in a cool, dry place that is inaccessible to children.
Pesticide Disposal: If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured for:
Farnam Pet Products
A Division of Farnam Companies, Inc.
P.O. Box 34820 • Phoenix, AZ 85067-4820

For information or comments, call 1-800-234-2269
Visit our Web site at adamsfleacontrol.com

Product Code: 100503443 • RM #: 300505573 • 09-1506
Printed in U.S.A.

EPA REG. No. 2596-139-270
EPA Est. No. 2724-TX-1

©2009 Farnam Companies, Inc.

Adams is a licensed trademark and Farnam and the Farnam logo are trademarks of Farnam Companies, Inc.

Central Garden & Pet and its logo are trademarks of Central Garden & Pet Company

RECEIVED
DEC - 2 2009
Pesticides Branch

ADAMS™ Flea & Tick Control Collar for Small Dogs
SMALL DOGS 7 Month Protection

ADAMS™

Flea & Tick Control Collar for Small Dogs

SMALL DOGS
7 Month Protection

SMALL DOGS
7 Month Protection

ADAMS™

Flea & Tick Control Collar for Small Dogs

KILLS:
- fleas
- ticks

KILLS & PREVENTS
- flea eggs
- larvae

ACTIVE INGREDIENTS:

Tetrachlorvinphos	14.55%
(S)-Methoprene	1.02%
OTHER INGREDIENTS:	84.43%
TOTAL:	100.00%

DO NOT LET CHILDREN PLAY WITH THIS COLLAR

CAUTION

See Back Panel for Additional Precautionary Statements

Net Weight: .69 OZ (19.8 g)

0 39079 09006 3

SMALL DOGS
7 Month Protection
ADAMS™ Flea & Tick Control Collar for Small Dogs