



FLEA & TICK COLLAR FOR SMALL DOGS



READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS/PUPPIES.

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. Buckle collar around dog's/puppy's neck. Fit collar snug enough to prevent placement of the dog's/puppy's leg or lower jaw through the collar, yet loose enough to be rotated about the neck. Generally, 2 or 3 fingers should fit between the collar and dog's/puppy's neck. **2.** Leave 2-3 extra inches on the collar for adjustment, cut off excess length, and dispose of in trash. **3.** Check periodically and adjust fit if necessary, especially when puppies are rapidly growing. **4.** May be used in addition to regular collar. Replace collar when effectiveness diminishes.

WHAT TO EXPECT:

For Fleas: **1.** This Bio Spot® Flea & Tick Collar for Small Dogs is designed to kill adult fleas. **2.** Protection against fleas will be achieved within a few days, providing continuous protection for up to 5 months. **3.** Replace collar every 5 months or sooner if needed.

For Ticks: **1.** Ticks are tougher to kill than fleas and are killed more slowly. **2.** Protection against adult and immature ticks will be achieved within a few days, providing continuous protection for up to 5 months. **3.** Adult ticks will be killed over the entire pet. Dead ticks will fall off or may be easily removed. **4.** Replace collar every 5 months or sooner if needed.

STORAGE and DISPOSAL: **Storage:** Store in original unopened package away from children. **Disposal:** Do not reuse container or collar. Dispose of in trash. If collar is temporarily removed, wrap in paper or plastic; do not allow children to have access to collar.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION: Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction*. The dust released by this collar is a cholinesterase inhibitor. **NOTICE: This product contains propoxur, a chemical known to the State of California to cause cancer.**

Humans: Do not open package until ready to use. Do not allow children to handle this collar. Dust will form on this collar during storage. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling collar.

Animals: Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. **Do not use on puppies under 12 weeks of age.** Do not use this product on dogs simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea and tick collars may be immediately replaced. When collar is first worn, observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water.

FIRST AID

Call a poison control center or doctor immediately for treatment advice.

If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If in eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 800-234-2269 for emergency medical treatment information.

NOTE TO PHYSICIAN/VETERINARIAN: Contains an N-methyl Carbamate that inhibits cholinesterase. *Atropine and "homatropine are antidotal only if symptoms of cholinesterase inhibition are present.

To the extent consistent with applicable law, Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

Satisfaction guaranteed or your money back. In case of emergency or for product use information, call 800-234-2269.

Manufactured for: Farnam Companies, Inc.
P.O. Box 34820, Phoenix, AZ 85067-4820
EPA Reg. No. 2724-254-270 • EPA Est. No. 2724-TX-1
Item: 100516474 • Art: 201310 • RM#: 300515660
©2006-2013 Farnam Companies, Inc.

Bio Spot and Bio Spot with design are registered trademarks of Farnam Companies, Inc. Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.



FLEA & TICK COLLAR FOR SMALL DOGS



FLEA & TICK COLLAR FOR SMALL DOGS



UP TO 5 MONTHS PROTECTION!

FLEA & TICK COLLAR FOR SMALL DOGS



KILLS:



fleas



ticks

For dogs with necks up to 15"

DO NOT ALLOW CHILDREN TO PLAY WITH THIS COLLAR

CAUTION

See Precautions on back

ACTIVE INGREDIENT:

Propoxur (CAS # 114-26-1) 10%

OTHER INGREDIENTS: 90%

TOTAL: 100%

NET WEIGHT: 0.69 OZ (19.8 g)



FLEA & TICK COLLAR FOR SMALL DOGS



0 39079 00106 9



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

9212.525