
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 889A-S4 0413



LICENSED

PERIOD **2013-2015** LIC. NO.

9226.252

Mertect® 340-F

Fungicide

Flowable (Water-Dispersible Suspension)

This supplemental label expires on 04/30/2016 and must not be used or distributed after that date.

Active Ingredient:

Thiabendazole: 2-(4-thiazolyl)benzimidazole* 42.3%

Other Ingredients: 57.7%

Total: 100.0%

*4.1 lbs. active ingredient/gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-889

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Mertect 340-F as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

SEED TREATMENT

Wheat, Winter and Spring

Control of seed- and soilborne Common Bunt (Stinking Smut):

For ready mix or slurry seed treaters, disperse 1.3 fl. oz. of Mertect 340-F for seedborne Common Bunt, or 2.6 fl. oz. per 100 lbs. of seed for soilborne Common Bunt, in an equal amount of water (1:1 ratio) and apply this dispersion per bushel. Keep dispersion under constant agitation.

For wheat seed having high infection levels of seedborne Fusarium scab, a lower rate of .17 fl. oz. per 100 lbs. of seed is recommended. Mertect 340-F should be used as a co-fungicide combined with Raxil- or Vitavax-based fungicides to include seed and seedling protection against the broad spectrum of diseases associated with the wheat crop, as listed on the product labels.

Winter Wheat Only:

Mertect 340-F aids in control of Dwarf Bunt in Colorado, Idaho, Montana, Oregon, Washington, Wyoming, and Utah: For ready mix or slurry seed treaters, disperse 2.6 – 3.9 fl. oz. of Mertect 340-F in an equal amount of water (1:1 ratio) and apply this selected dispersion per 100 lbs. of seed. Keep dispersion under constant agitation. For best results against Dwarf Bunt, plant winter wheat treated with Mertect 340-F as late in the growing season as is practical. Use only on normal or late winter wheat plantings. Use the lower rate for late plantings or winter wheat. The level of product effectiveness will vary with the weather conditions and amount of plant growth which occurs prior to the disease infection period.

Mertect® and the Syngenta logo are trademarks of a Syngenta Group Company

©2013 Syngenta

SUPPLEMENTAL LABELING

Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 889A-S3 0510



LICENSED

PERIOD **2013-2015** LIC. NO. **9226.252**



LICENSED

PERIOD **2010-2012** LIC. NO.

Mertect® 340-F

Fungicide

Flowable (Water-Dispersible Suspension)

Active Ingredient:

Thiabendazole: 2-(4-thiazolyl)benzimidazole* 42.3%

Other Ingredients: 57.7%

Total: 100.0%

*4.1 lbs. active ingredient/gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-889

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Mertect 340-F as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

CROP USE DIRECTIONS

Mushrooms (*Agaricus* spp.) - Dry Bubble (*Verticillium fungicola*), Wet Bubble (*Mycogone perniciosus*), Cobweb (*Dactylium dendroides*) and Green Mold (*Trichoderma* spp.):

FOR USE AS AN ADMIX WITH MUSHROOM SPAWN GRAINS, MUSHROOM GROWING SUPPLEMENT, AND/OR AS A DRENCH DURING NORMAL WATERING OPERATIONS FOR MUSHROOM GROWING BEDS AT CASING OR BETWEEN BREAKS

NOTE: Do not exceed a combined total of 20 fl. oz. per 1,000 sq. ft. of product, between the various applications to spawn grains, supplement, and mushrooms. This amount may be used in various treatment regimes. Please refer to the tables below for the rates for the various usage so that the 20 fl. oz. per 1,000 sq. ft. is not exceeded.

Spawn Grains: Apply a maximum of 5.4 fl. oz. of Mertect 340-F per 1,000 sq. ft. Follow the table below to determine the fl. oz. of Mertect 340-F that can be mixed with the spawn to deliver 5.4 fl. oz. per 1,000 sq. ft.

Dosage Table for Use on Spawn Grains

Do not apply Mertect 340-F to gypsum, limestone, or chalk. Admix Mertect 340-F directly onto spawn grain.

Pounds spawn per 8000 sq. ft.	Max. fl. oz. Mertect per 100 pounds spawn
960	4.5
1040	4.1
1120	3.8
1200	3.6
1280	3.4
1360	3.2
1440	3.0

Mushroom Supplements: A finished batch of mushroom supplement can contain up to 0.1% thiabendazole (by weight). The thiabendazole added to the supplement must be included as part of the total of the 20 fl. oz. of Mertect 340-F allowed per 1,000 sq. ft. per crop. The total amount of Mertect 340-F that can be applied from all treatments per crop is dependent on the amount of supplement applied (use the table below to determine the remaining volume of Mertect 340-F that can be applied per crop).

Lbs supplement* per 8,000 sq. ft.	Fl. oz. Mertect 340-F remaining that may be applied to mushrooms, supplement, casing, or other timing	Fl. oz. Mertect 340-F remaining that may be applied to mushrooms, supplement, casing, or other timing if a spawn treatment was also applied
800	17	11.6
1200	15	9.6
1400	14.5	9.1
1800	13	7.6
2200	11	5.6
2400	9	3.6

*Assumes at maximum rate of 0.1% thiabendazole.

The amount added for supplement treatment must be deducted from the total remaining amount that can be applied to mushrooms so as not to exceed 20 fl. oz. per 1,000 sq. ft. per crop.

Note: If you are unsure if the mushroom supplement that you are using contains thiabendazole, contact the manufacturer of the mushroom supplement.

Applying to Mushrooms: Refer to the tables above for the amount of Mertect 340-F used in any spawn grain or supplement treatments before applying Mertect 340-F to the mushrooms to ensure that 20 fl. oz. per 1,000 sq. ft. is not exceeded.

Casing Applications: Apply no more than 8 fl. oz. Mertect 340-F per 1000 sq. ft. Mertect 340-F may be applied at casing as a drench during normal watering operations.

Other Timings: Apply no more than 4 fl. oz. Mertect 340-F per 1000 sq. ft.

Do not apply within 12 hours prior to harvest.

Mertect® and the Syngenta logo are trademarks of a Syngenta Group Company

©2010 Syngenta



LICENSED

PERIOD 2013-2015 LIC. NO. 9226.252

PULL HERE TO OPEN ►



Mertect[®]340-F



Fungicide

Flowable (Water-Dispersible Suspension)

Active Ingredient:

Thiabendazole: 2-(4-thiazolyl)benzimidazole* 42.3%

Other Ingredients: 57.7%

Total: 100.0%

*4.1 lbs. active ingredient/gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-889 EPA Est. 39578-TX-1

Product of India
Formulated in the USA

SCP 889A-L1Q 0413
4025273

2 gallons
Net Contents



TM

FIRST AID	
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-888-8372 for emergency medical treatment information.</p>	
<p>HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if inhaled. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes plus socks

In addition to the above, handlers making application to mushroom houses using hand held sprayers must wear chemical resistant gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

continued...

PRECAUTIONARY STATEMENTS (continued)

Environmental Hazards

This pesticide is toxic to fish. Do not apply when weather conditions favor runoff or drift from the target area. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of waste.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

NOT FOR USE IN FORMULATING AGRICULTURAL PESTICIDES.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Manual seed treatments are prohibited.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the REI of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

GENERAL INFORMATION

Mertect 340-F is a systemic fungicide which aids in the control of many important plant diseases. If treatment is not effective following use of Mertect 340-F as recommended, a tolerant strain of fungus may be present. Consideration should be given to prompt use of other suitable non-benzimidazole fungicides. Mertect 340-F is not for use in formulating agricultural products.

CROP USE DIRECTIONS

GENERAL - SHAKE OR STIR WELL BEFORE USE.

Clean equipment before using Mertect 340-F. **When Mertect 340-F is dispersed in water, the resulting suspension must be constantly agitated.** Avoid foaming.

Use the higher recommended rates under conditions of heavier disease pressure.

Compatibility information for Mertect 340-F with other chemicals is limited. Consult your supplier or agricultural experiment station for specific recommendations.

Do not apply this product through any type of irrigation system.

Carrots - *Botrytis Cinerea* Gray Mold and *Sclerotinia sclerotiorum* (Sclerotinia Rot): Dip carrots, before storage, for 5-10 seconds in a suspension containing 41 fl. oz. of Mertect 340-F per 100 gals. of water. Replenish the suspension when the volume is too low or when it becomes dirty.

The runoff and wastes from the dipping operation should not be discarded in a drainage which could contaminate public water systems.

Mushrooms (*Agaricus* spp.) - Dry Bubble (*Verticillium fungicola*), Wet Bubble (*Mycogone perniciosa*), Cobweb (*Dactylium dendroides*) and Green Mold (*Trichoderma* spp.):

FOR USE AS AN ADMIX WITH MUSHROOM SPAWN GRAINS, MUSHROOM GROWING SUPPLEMENT, AND/OR AS A DRENCH DURING NORMAL WATERING OPERATIONS FOR MUSHROOM GROWING BEDS AT CASING OR BETWEEN BREAKS

NOTE: Do not exceed a combined total of 20 fl. oz. per 1,000 sq. ft. of product, between the various applications to spawn grains, supplement, and mushrooms. This amount may be used in various treatment regimes. Please refer to the tables below for the rates for the various usage so that the 20 fl. oz. per 1,000 sq. ft. is not exceeded.

Spawn Grains: Apply a maximum of 5.4 fl. oz. of Mertect 340-F per 1,000 sq. ft. Follow the table below to determine the fl. oz. of Mertect 340-F that can be mixed with the spawn to deliver 5.4 fl. oz. per 1,000 sq. ft.

Dosage Table for Use on Spawn Grains

Do not apply Mertect 340-F to gypsum, limestone, or chalk. Admix Mertect 340-F directly onto spawn grain.

Pounds spawn per 8000 sq. ft.	Max. fl. oz. Mertect per 100 pounds spawn
960	4.5
1040	4.1
1120	3.8
1200	3.6
1280	3.4
1360	3.2
1440	3.0

Mushroom Supplements: A finished batch of mushroom supplement can contain up to 0.1% thiabendazole (by weight). The thiabendazole added to the supplement must be included as part of the total of the 20 fl. oz. of Mertect 340-F allowed per 1,000 sq. ft. per crop. The total amount of Mertect 340-F that can be applied from all treatments per crop is dependent on the amount of supplement applied (use the table below to determine the remaining volume of Mertect 340-F that can be applied per crop).

Lbs supplement* per 8,000 sq. ft.	Fl. oz. Mertect 340-F remaining that may be applied to mushrooms, supplement, casing, or other timing	Fl. oz. Mertect 340-F remaining that may be applied to mushrooms, supplement, casing, or other timing if a spawn treatment was also applied
800	17	11.6
1200	15	9.6
1400	14.5	9.1
1800	13	7.6
2200	11	5.6
2400	9	3.6

*Assumes at maximum rate of 0.1% thiabendazole.

The amount added for supplement treatment must be deducted from the total remaining amount that can be applied to mushrooms so as not to exceed 20 fl. oz. per 1,000 sq. ft. per crop.

Note: If you are unsure if the mushroom supplement that you are using contains thiabendazole, contact the manufacturer of the mushroom supplement.

Applying to Mushrooms: Refer to the tables above for the amount of Mertect 340-F used in any spawn grain or supplement treatments before applying Mertect 340-F to the mushrooms to ensure that 20 fl. oz. per 1,000 sq. ft. is not exceeded.

Casing Applications: Apply no more than 8 fl. oz. Mertect 340-F per 1000 sq. ft. Mertect 340-F may be applied at casing as a drench during normal watering operations.

Other Timings: Apply no more than 4 fl. oz. Mertect 340-F per 1000 sq. ft.

Do not apply within 12 hours prior to harvest.

Ornamental Bulbs and Corms (One Application Only) - *Fusarium* Basal Rot and *Penicillium* Blue Mold: Clean and treat bulbs or corm within 24-48 hours of digging. Prepare suspension of 30 fl. oz. of Mertect 340-F per 100 gals. water. Submerge stock completely in dipping suspension. Suspension temperature should be 55-75°F (13-24°C). Discard suspension (1) when it becomes dirty, (2) after using five times, or (3) after 24 hours, whichever occurs first.

Fusarium (Bulbs and Corms): Dip bulbs 15-30 minutes, corms 15 minutes.

Blue Mold (Bulbs): Dip 10-15 minutes.

After treatment, dry bulbs or corms in a shaded, well-ventilated area. Curing or retarding may precede or follow treatment.

Potato - Fusarium Tuber Rot: Mist unwashed tubers on a conveyor line, with tumbling action, entering storage with 0.42 fl. oz. of Mertect 340-F to each 2,000 lbs. of tubers in sufficient water for complete coverage. If an additional treatment is necessary before shipping, mist the tubers at the same rate or dip the tubers for 20 seconds in a solution containing 0.42 fl. oz. of Mertect 340-F per gallon of water. Do not treat seed potatoes after cutting.

Mertect 340-F for potatoes should be used as part of an Integrated Management Program, which includes prevention of bruising during harvest and maintaining proper storage temperature and relative humidity. Neither cultural practices nor chemical measures will prevent Fusarium infection if potatoes are damaged or improperly stored.

Pome Fruit (Apples and Pears) - Blue Mold Rot, Bull's Eye Rot, and Gray Mold (Nest Rot, Cluster Rot), Stem End and Neck Rot: Dip, flood, or spray harvested fruit with a suspension of 16 fl. oz. of Mertect 340-F in 100 gals. of water. Do not treat for over 3 minutes. Treat apples only before and after storage for maximum decay control. Treat pears only once.

Sweet Potato (One Application Only) "Seed Roots" - Black Rot, Scurf, and Foot Rot: Dip the seed roots in a suspension containing 8 fl. oz. of Mertect 340-F per 7.5 gals. of water (107 fl. oz. per 100 gals.). Treat the seed roots for 1-2 minutes and plant immediately. Discard the suspension when the volume is too low or when it becomes dirty. Do not use the treated roots for food or feed.

SEED TREATMENT

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

Seed treatment with Mertect 340-F fungicide will not provide postemergence protection of plants grown in or near infected fields.

Mixing Procedures

Apply Mertect 340-F as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the recommended amount of Mertect 340-F into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow seed to dry before bagging.

Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with thiabendazole fungicide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed."

Mixing with other pesticides is limited to adequate physical compatibility and satisfactory phytotox test results.

Soybean (One Application Only): Mertect 340-F is recommended for protection against pod and stem blight (*Phomopsis* spp.) on soybean seed to improve germination and early season seedling health. For protection against other seed and soil-borne diseases, it is recommended that a co-fungicide such as Syngenta Apron® XL + Maxim® 4FS, or an Apron MAXX® brand fungicide, be used at EPA registered label rates and tank mixed in combination with Mertect 340-F. Consult the label for appropriate use rates and follow all use instructions for the products. Utilized as a seed treatment for protection against seed borne and early season seedling pod and stem blight on soybean: Apply Mertect 340-F at the rate of 0.08 to 0.16 fl. oz. per 100 lbs. of seed.

Peas (Dry)

Chickpea (garbanzo beans)
(*Cicer arietinum*)

Lentil
(*Lens esculenta*)

Peas
(*Pisum* spp.)

Pigeon peas
(*Cajanus cajan*)

Mertect 340-F fungicide is used to effectively suppress seed borne *Ascochyta* blight caused by the pathogen *Ascochyta rabiei*. Mertect 340-F seed treatment must also be used to minimize the threat of highly infectious and limit spread of residue-borne and wind-borne infection of chickpeas and lentil crops by *Ascochyta* blight.

Chickpea Seed Treatment: For slurry solutions in water, mix 2.04 fl. oz. of Mertect 340-F fungicide in an equal amount of water (1:1) to provide adequate coverage and apply to 100 lbs. of seed. Keep dispersion under constant agitation. It is essential that thorough coverage of seed be obtained.

Lentil Seed Treatment: For slurry solutions in water, mix 1.05 fl. oz. of Mertect 340-F fungicide in an equal amount of water (1:1) to provide adequate coverage and apply to 100 lbs. of seed. Keep dispersion under constant agitation. It is essential that thorough coverage of seed be obtained.

Peas Seed Treatment: For a ready mix or slurry treaters, mix 1.02 fl. oz. of Mertect 340-F fungicide in adequate water to provide complete coverage and apply to 100 lbs. of seed. Keep dispersion under constant agitation. It is essential that thorough coverage of seed be obtained.

For Export Use on Peas: Mertect 340-F may be mixed with other registered seed treatment products in accordance with the importing Nation's Phytosanitary requirements. Treated seed must be labeled "for Export Only"- Not to be Sold or Offered for Sale in the United States.

For best results against *Ascochyta* blight, plant chickpeas, peas or lentil seed treated with Mertect 340-F fungicide as late in the spring as possible.

Restrictions and Limitations

- Maximum one application per season.
- The product must not be tank mixed with any pesticide unless physical compatibility and phytotoxicity tests have been completed and proved to be satisfactory.
- Do not feed vines or hay grown from treated seed to livestock.

Wheat, Winter and Spring

Control of seed- and soilborne Common Bunt (Stinking Smut):

For ready mix or slurry seed treaters, disperse 1.3 fl. oz. of Mertect 340-F for seedborne Common Bunt, or 2.6 fl. oz. per 100 lbs. of seed for soilborne Common Bunt, in an equal amount of water (1:1 ratio) and apply this dispersion per bushel. Keep dispersion under constant agitation.

For wheat seed having high infection levels of seedborne *Fusarium* scab, a lower rate of .17 fl. oz. per 100 lbs. of seed is recommended. Mertect 340-F should be used as a co-fungicide combined with Raxil- or Vitavax-based fungicides to include seed and seedling protection against the broad spectrum of diseases associated with the wheat crop, as listed on the product labels.

Winter Wheat Only:

Mertect 340-F aids in control of Dwarf Bunt in Colorado, Idaho, Montana, Oregon, Washington, Wyoming, and Utah: For ready mix or slurry seed treaters, disperse 2.6 – 3.9 fl. oz. of Mertect 340-F in an equal amount of water (1:1 ratio) and apply this selected dispersion per 100 lbs. of seed. Keep dispersion under constant agitation. For best results against Dwarf Bunt, plant winter wheat treated with Mertect 340-F as late in the growing season as is practical. Use only on normal or late winter wheat plantings. Use the lower rate for late plantings or winter wheat. The level of product effectiveness will vary with the weather conditions and amount of plant growth which occurs prior to the disease infection period.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store Mertect 340-F in a cool, dry place away from food, drink, and animal feeding stuffs. Do not store below 32°F (0°C). Repeated product freezing and thawing should be avoided. Shake or stir product before use.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility in accordance with Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available.

Residue Removal [capacities equal to or less than 5 gallons]

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

For Bulk and Minibulk Containers:

Residue Removal [capacities greater than 5 gallons]

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Container Disposal

Refillable container. Refill this container with Mertect 340-F only. Do not reuse this container for any other purpose.

Residue Removal

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Apron®, Apron MAXX®, Maxim®, Mertect®, the ALLIANCE FRAME 
the SYNGENTA Logo and the PURPOSE ICON 
are Trademarks of a Syngenta Group Company

©2013 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for
Syngenta Crop Protection, LLC.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 889A-L1Q 0413
4025273



KEEP OUT OF REACH OF CHILDREN. CAUTION

See directions for use inside booklet.

Fungicide

Flowable (Water-Dispersible Suspension)

Active Ingredient:

Thiabendazole:

2-(4-thiazolyl)benzimidazole* 42.3%

Other Ingredients: 57.7%

Total: 100.0%

*4.1 lbs. active ingredient/gallon

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-889

EPA Est. 39578-TX-1

Mertect® and the Syngenta logo are Trademarks of a Syngenta Group Company

©2013 Syngenta

Manufactured for
Syngenta Crop Protection, LLC.
P.O. Box 18300
Greensboro, North Carolina 27419-8300

**SCP 889A-L1Q 0413
4025273**

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-888-8372 for emergency medical treatment information.

HOT LINE NUMBER: For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call **1-800-888-8372**.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Harmful if inhaled. Avoid breathing spray mist.

Environmental Hazards

This pesticide is toxic to fish. Do not apply when weather conditions favor runoff or drift from the target area. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of waste.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

NOT FOR USE IN FORMULATING AGRICULTURAL PESTICIDES.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store Mertect 340-F in a cool, dry place away from food, drink, and animal feeding stuffs. Do not store below 32°F (0°C). Repeated product freezing and thawing should be avoided. Shake or stir product before use.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility in accordance with Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available.

Residue Removal [capacities equal to or less than 5 gallons]

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

2 gallons
Net Contents



BAR CODE # IS
(01) 0 07 02941 41942
LAST DIGIT IS CHECK DIGIT
UCC/EAN 128