



STATE OF HAWAII
 Department of Agriculture
ACCEPTED
 LICENSE NO. **9226.262**

PULL HERE ►
 TO OPEN



syngenta.

Fungicide

A seed treatment product for protection against certain seed and seedling diseases of crop plants

1 quart

Net Contents

<i>Active Ingredient:</i>	
Fludioxonil (CAS No. 131341-86-1)	40.3%
<i>Other Ingredients:</i>	59.7%
<i>Total:</i>	100.0%

Maxim 4FS contains 4 lb active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-758 EPA Est. 100-NE-001

**Product of Switzerland
 Formulated in the USA**

**SCP 758A-L7P 0811
 4046778**

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly. • Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance for Incidents Involving Human or Animal Exposure or for Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if inhaled. Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or Viton®. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.
- Shoes plus socks

continued...

PRECAUTIONARY STATEMENTS (*continued*)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsates.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Maxim 4FS must be used only in accordance with directions on this label or in separately published Syngenta supplemental labeling approved by EPA for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

continued...

AGRICULTURAL USE REQUIREMENTS (continued)

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or Viton
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

GENERAL INFORMATION

Maxim 4FS is a seed treatment fungicide which provides protection against certain soil-borne and seed-borne diseases of crop plants. Maxim 4FS is active against *Fusarium*, *Rhizoctonia*, *Helminthosporium*, and weakly pathogenic fungi such as *Aspergillus* and *Penicillium*. When rate ranges are given, use higher rate when disease pressure is expected to be severe.

MIXING PROCEDURES

Apply Maxim 4FS as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of Maxim 4FS into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer's application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow seed to dry before bagging. Treated seed bagged for later use must contain an EPA-approved dye or colorant that imparts an unnatural color to the seed.

Federal Law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with fludioxonil fungicide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs. Forage may not be grazed until 30 days after planting."

Maxim 4FS does not control diseases caused by *Pythium* spp. or *Phytophthora* spp. If these diseases are expected to be a problem, apply Maxim 4FS tank mixed with Apron XL®. If Apron XL is not registered on that crop, apply Maxim as a tank mix with another labeled seed treatment product that has activity against these diseases. Consult the label for appropriate use rate and follow all use instructions for the products.

Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing.

BRASSICA (COLE) LEAFY VEGETABLES GROUP

Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa, and Chinese mustard), cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

BULB VEGETABLES

Garlic (including great headed), leek, onion (dry bulb and green), welch onion, and shallot

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed or propagating material.

CEREAL GRAINS (INCLUDES FORAGE AND SILAGE)

Barley, buckwheat, corn (field corn, popcorn, and sweet corn), millet (pearl and proso), oats, rice, rye, sorghum, teosinte, triticale, wheat, and wild rice

Corn (Field Corn and Popcorn): For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, and for control of seed-borne head smut (*Sphacelotheca reiliana*), apply Maxim 4FS as shown below. For *Pythium* spp. control, use Maxim 4FS in combination with labeled rates of Apron XL (100-799) and Dynasty® (100-1159). Observe all precautions, limitations, rates, and directions for use on the respective labels before applying.

Maxim 4FS Use Rate		
fl oz/ 100 lb seed	fl oz/80,000 kernel count	mg active ingredient per kernel
0.08	0.036	0.0064
0.16	0.072	0.0128

Maxim 4FS contains 14.18 grams active ingredient per ounce.

Grain Sorghum: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed (0.0008-0.0016 mg fludioxonil per seed*).

Apply Apron XL in combination with Maxim 4FS for protection against damping-off due to *Pythium* spp. Apply 0.32-0.64 fl oz of Apron XL per 100 lb of seed (0.0023-0.0047 mg mefenoxam per seed*) tank mixed with labeled rates of Maxim 4FS.

For systemic downy mildew control on cultivars which possess sorghum downy mildew resistance, apply 0.64 fl oz of Apron XL per 100 lb of seed (0.0047 mg mefenoxam per seed*) as a tank mix with recommended rates of Maxim 4FS. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew grown in areas with recent history of severe disease, apply 1.28 fl oz of Apron XL (0.0094 mg mefenoxam per seed*) tank mixed with recommended rates of Maxim 4FS. Follow all application instructions found on the Apron XL label.

*Based upon 14,500 seeds per pound of sorghum seed.

Note: Seed treated with the 1.28 fl oz of Apron XL should be planted and not stored or carried over to the following year. Only high quality sorghum seed ($\geq 90\%$ germ) should be treated with Apron products.

Rice: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.02 to 0.08 fl oz of Maxim 4FS per 100 lb of seed.

Sweet Corn: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08 fl oz of Maxim 4FS per 100 lb of seed.

Apply Apron XL seed treatment product in combination with Maxim 4FS for protection against damping-off due to *Pythium* spp. Apply Apron XL at 0.32 fl oz per 100 lb of seed tank mixed with the label rates of Maxim 4FS. For protection against systemic downy mildew of sweet corn, apply Apron XL at 1.28 fl oz per 100 lb of seed tank mixed with label rates of Maxim 4FS. Follow application and other instructions found on the Apron XL product label.

For all other cereal crops: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

Important: (1) Do not use treated seed for feed, food, or oil. (2) Cereal forage may not be grazed until 30 days after planting.

CUCURBIT VEGETABLES

Chayote, Chinese waxgourd, citron melon, cucumber, gherkin, gourd edible (includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp., all muskmelon, pumpkin, all squash, and watermelon

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

FLAX, COTTON, CRAMBE, JAJOBA, LESQUERELLA, RAPE, SAFFLOWER, SUNFLOWER

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

For sunflower, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed or 0.0025-0.005 mg active ingredient per seed.

FOLIAGE OF LEGUME VEGETABLES

Any cultivar of bean (*Phaseolus* spp.), pea (*Pisum* spp.), and soybean

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

FRUITING VEGETABLES EXCEPT CUCURBITS

Eggplant, groundcherry, pepino, pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

GRASS FORAGE, FODDER, AND HAY

All grasses (including turfgrass, bermudagrass, bluegrass, and fescue or bromegrass), Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed or propagating root or tuber material.

Important: Forage may not be grazed until 30 days after planting.

HERBS AND SPICES

Allspice, angelica, anise, anise (star), annatto, balm, basil, borage, burnet, camomile, caper buds, caraway, caraway (black), cardamom, cassia bark, cassia buds, catnip, celery seed, chervil, chive, Chinese chive, cinnamon, clary, clove buds, coriander (Chinese parsley or cilantro), costmary, culantro (seed), cumin, curry (leaf), dill (seed) and dillweed, common fennel and Florence fennel, fenugreek, grains of paradise, horehound, hyssop, juniper berry, lavender, lemongrass, lovage, lovage (seed), mace, marigold, marjoram, mustard (seed), nasturtium, nutmeg, parsley (dried), pennyroyal, pepper (black and white), poppy (seed), rosemary, rue, saffron, sage, savory (summer and winter), sweet bay, tansy, tarragon, thyme, vanilla, wintergreen, woodruff, and wormwood

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed or propagating root or tuber material.

LEAFY VEGETABLES EXCEPT BRASSICA

Amaranth (leafy, Chinese spinach, tampala), arugula, cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (finocchio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach (including New Zealand and vine), and Swiss chard

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

LEAVES OF ROOT AND TUBER VEGETABLES

Beet (garden and sugar), burdock (edible), carrot, cassava (bitter and sweet), chervil (turnip rooted), chicory, dasheen (taro), parsnip, radish (including oriental), rutabaga, salsify (black), sweet potato, tanier, turnip, and true yam

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed or propagating root or tuber material.

LEGUME VEGETABLES (SUCCULENT OR DRY) (EXCEPT SOYBEANS)

Bean (*Lupinus* spp. – includes grain lupin, sweet lupin, white lupin), bean (*Phaseolus* spp. – includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), bean (*Vigna* spp. – includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, and yardlong bean), broad bean, chickpea, guar, jackbean, lablab bean, lentil, pea (dwarf, edible-pod, English, field, garden, green, snow, sugar snap), pigeon pea, soybean (immature), and sword bean

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed or propagating root or tuber material.

NONGRASS ANIMAL FEEDS (FORAGE, FODDER, STRAW, AND HAY)

Alfalfa, bean (velvet), clover, kudzu, lespedeza, lupin, sainfoin, trefoil, and vetch (including crown vetch and milk vetch)

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

Important: Forage may not be grazed until 30 days after planting.

PEANUTS

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed.

ROOT AND TUBER VEGETABLES

Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley, parsnip, potato (0.08 fl oz only), radish (including oriental), rutabaga, salsify (oyster, black, and spanish), skirret, sweet potato, taniel, turmeric, turnip, yam bean (jicama, manioc pea), and true yam

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl oz of Maxim 4FS per 100 lb of seed or propagating root or tuber material.

Note for Application to Potato: Maxim 4FS applications to potato seed pieces must be made only through specific types of equipment. Please contact your Syngenta seed treatment representative to obtain information on proper equipment for application.

Note for Potatoes Intended for Seed Production: As a resistance management strategy, if Maxim 4FS is used on potatoes intended for seed production, a labeled rate of mancozeb seed treatment dust must be applied to seed tubers after the Maxim 4FS treatment. If the mancozeb dust is not used, an in-furrow application of Quadris® at 0.40 fl oz /1000 ft. of row or Amistar® at 0.13 fl oz/1000 ft. of row must be applied over the Maxim treated seed tubers.

SOYBEANS

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply Maxim 4FS as shown here:

Maxim 4FS Use Rate*			
fl oz/100 lb seed	fl oz/1,000 seeds	fl oz/140,000 seeds	mg active ingredient per seed
0.08	0.00027	0.0373	0.0038
0.16	0.00053	0.0747	0.0076

*Maxim 4FS contains 14.175 grams active fludioxonil per fluid ounce of Maxim 4FS. Based on 3,000 soybean seeds per pound.

SEED TREATMENT - EXPORT USE

FOR TREATMENT OF THE SEED FOR EXPORT ONLY TO COUNTRIES REQUIRING TREATMENT WITH MAXIM 4FS AND APRON XL BEFORE IMPORTATION: User should verify actual registered rate in country of destination and/or use.

Seed treated in accordance with the export use pattern must be labeled as follows: TREATED SEED: FOR EXPORT ONLY – NOT FOR DOMESTIC SALE OR USE. TREATED WITH THE FUNGICIDE, FLUDIOXONIL

Use Restrictions

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed package or containers in accordance with local requirements.

For seeds that are treated and bagged for later use, the following must appear on the bag.

- This seed has been treated with fludioxonil seed treatment.
- Do not use treated seed for feed, food or oil purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry, secure place.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance in proper disposal methods.

Container Handling [less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

continued...

STORAGE AND DISPOSAL (continued)

Container Handling [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Maxim®, Amistar®, Apron XL®, Dynasty®, Quadris®, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



Viton® trademark of E. I. duPont de Nemours & Company.

©2014 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 758A-L7P 0811
4046778



syngenta.

Fungicide

A seed treatment product for protection against certain seed and seedling diseases of crop plants

1 quart

Net Contents

Maxim[®] 4FS Fungicide

A seed treatment product for protection against certain seed and seedling diseases of crop plants

Active Ingredient:
Fludioxonil (CAS No. 131341-86-1) . . . 40.3%
Other Ingredients: . . . 59.7%
Total: . . . 100.0%

Maxim 4FS contains 4 lb active ingredient per gallon.

See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-758

EPA Est. 100-NE-001

Maxim[®] and the Syngenta logo are trademarks of a Syngenta Group Company

©2014 Syngenta

Manufactured for:

Syngenta Crop Protection, LLC

P.O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 758A-L7P 0811 4046778

KEEP OUT OF REACH OF
CHILDREN.

CAUTION

FIRST AID

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER: For 24-Hour Medical Emergency Assistance for Incidents Involving Human or Animal Exposure or for Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call 1-800-888-8372.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if inhaled. Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsates.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry, secure place.

Pesticide Disposal


Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance in proper disposal methods.

Container Handling [less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

syngenta[®]



00110702941812369