

STYLE A  
PK2792

Department of Agriculture  
STATE OF HAWAII  
LICENSED  
PERIOD 2011-2013 LIC. NO.

Department of Agriculture  
STATE OF HAWAII  
DISC. LICENSED  
PERIOD 2014-2016 LIC. NO.  
9227.285

**Directions for Use**

- How to Open**
1. Remove product tubes from the package.
  2. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.
- How to Apply**
1. Invert tube over dog and use open end to part dog's hair.
  2. Squeeze tube firmly to apply all of the solution to the dog's skin as a spot to the dog's back between the shoulder blades.



**6 Step Topical Checklist**

- Read the label completely and follow directions
- Do NOT treat with more than one pesticide product at a time
- Weigh your pet
- Separate animals after treatment to avoid chance of ingestion
- Do NOT use dog products on cats
- Do NOT split tubes between pets

**3 Month Application Tracker**

**First Application**  
DATE: mm / dd / yy

**Second Application**  
DATE: mm / dd / yy

**Third Application**  
DATE: mm / dd / yy

For year-round protection, apply PRONYL OTC™ PLUS monthly.

**PRONYL OTC™ PLUS for dogs**

**PRONYL OTC™ PLUS for dogs**  
For use ONLY on dogs over 8 weeks of age  
Up to 22 lbs



READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.  
PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS. CAUTION.** Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

**HAZARDS TO DOMESTIC ANIMALS.** For external use only. Do not use on puppies under 8 weeks of age. Individual sensitivities, while rare, may occur after using any pesticide product. Some dogs may experience temporary startle effects when any product is applied. Dogs may experience some temporary irritation at the site of product application. If signs persist, or become more severe within a few days of application, consult a veterinarian immediately. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating treatment. Certain medications can interact with pesticides. Consult a veterinarian before using on medicated, debilitated, or aged dogs. Call 1-800-781-4738 for 24-hour assistance.

FIRST AID	
If swallowed:	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	Flush eyes with plenty of water. Call a doctor if irritation persists.
If on skin:	Wash with plenty of soap and water. Get medical attention if irritation persists.

**HOTLINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-781-4738 for emergency medical treatment information.

**DIRECTIONS FOR USE**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT ALLOW CHILDREN TO APPLY PRODUCT.** TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. FOR EXTERNAL USE ON DOGS ONLY. DO NOT USE ON RABBITS. DO NOT USE ON OTHER ANIMALS.

Do not use on puppies under 8 weeks of age. Do not split one tube between dogs. Do not use multiple tubes on one dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Consult a veterinarian before using this product on debilitated, aged, or medicated dogs. Can be used on breeding, pregnant, or lactating dogs. Consult a veterinarian before using on dogs with known organ dysfunction. Dogs may experience some temporary irritation at the site of product application. Certain medications can interact with pesticides. Sensitivities may occur after using ANY pesticide product on dogs. If signs of sensitivity occur bathe your dog with a mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**How to open:** Remove product tubes from the package. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

**How to apply:** Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Do not apply more often than once every 4 weeks.

**FREQUENCY OF APPLICATION**  
Use PRONYL OTC™ PLUS monthly for complete control of flea, tick, and chewing lice infestations. Studies show that PRONYL OTC™ PLUS kills adult fleas for up to three months. Apply monthly if your dog is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. PRONYL OTC™ PLUS remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 4 weeks.

**For Dogs Weighing up to 22 lbs:** Apply one tube (0.67 mL)(0.023 fl oz) as a spot on the dog's back between the shoulder blades.

In Case of Emergency, call: 1-800-781-4738  
Non-Emergency, call: 1-800-224-PETS (7387)  
Made in USA

If you have questions or comments about this product, please write: Sergeant's Consumer Response, P.O. Box 540399, Omaha, NE 68154-0399.

Distributed by: Sergeant's Pet Care Products, Inc. P.O. Box 540399, Omaha, NE 68154-0399  
EPA Reg. No. 2517-134  
EPA Est. No. 2517-TN-1 P003060 AIRR



**PRONYL OTC™ PLUS for dogs**  
For use ONLY on dogs  
Up to 22 lbs  
over 8 weeks of age

**3 MONTH SUPPLY**  
For use ONLY on dogs  
Up to 22 lbs  
over 8 weeks of age

Sergeant's  
**PRONYL OTC™ PLUS for dogs**

Contains Fipronil and (S)-methoprene, the active ingredients used in Frontline Plus® for dogs\*



- KILLS**
- fleas
  - ticks
  - chewing lice
  - flea eggs
  - flea larvae

- Controls flea reinfestation for up to 3 months
- Controls mites that may cause sarcoptic mange
- Waterproof

\* PRONYL OTC™ PLUS for dogs is not manufactured by or distributed by Merial. Frontline Plus® is a registered trademark of Merial.

**KEEP OUT OF REACH OF CHILDREN CAUTION**  
See Back Label Panel for Additional Precautionary Statements  
Net Contents: Three 0.023 fl oz (0.67 mL) applicators


**ACTIVE INGREDIENTS:**

Fipronil	9.80%
(S)-methoprene	8.80%
<b>OTHER INGREDIENT**:</b>	<b>81.40%</b>
<b>TOTAL</b>	<b>100.00%</b>

\*\*diethylene glycol monoethyl ether


**LIFT HERE FOR MORE INFORMATION**

**PRONYL OTC™ PLUS for dogs**



- ▶ Kills adult fleas, flea eggs, flea larvae, and prevents the development of pupae for up to 3 months
- ▶ Kills all stages of deer ticks
- ▶ Kills ticks including those that may transmit Lyme disease
- ▶ Kills brown dog ticks, American dog ticks, lone star ticks, and deer ticks
- ▶ Controls mites that may cause sarcoptic mange
- ▶ Can be used on breeding, pregnant, and lactating dogs
- ▶ Fast acting
- ▶ Long lasting

The active ingredients in PRONYL OTC™ PLUS feature a sustained release formula developed for fast acting and long lasting control of fleas, ticks, and chewing lice for up to 4 weeks on dogs.



PROJECT NAME & PROJECT NUMBER	DATE	DRAFT#	DIMENSIONS	UPC CODE	PART#
Pronyl OTC Plus	7FEB11	25		UPC 0 73091 02965 6	P003060
<b>ART INSTRUCTION FOR THE PRINTER</b>			Printer/Vendor:	Date sent:	
	4-COLOR PROCESS (CMYK)	Pantone 2627	Pantone 165		
<b>Additional Notes:</b>					
Please see the "Emboss" layer on this file. The logo "PRONYL OTC PLUS" will be embossed.			Emboss		
			<b>Sergeant's</b> Pet Care Products, Inc.		