STYLE A

PK2792

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

DISC.

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO. 9227.302

Directions for Use

How to Open:

1. Remove product tubes from the package.
2. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

How to Apply:

1. Invert tube over cat and use open end to part cat's hair.
2. Apply contents of one tube of PRONYL OTC™ PLUS as a spot between the shoulder blades. Squeeze tube firmly to apply all of the solution to the cat's skin.

6 Step Topical Checklist

- ☐ Read the label completely and follow directions
- ☐ Weigh your pet
- ☐ Do NOT use dog products on cats
- ☐ Do NOT treat with more than one pesticide product at a time
- ☐ Separate animals after treatment to avoid chance of ingestion
- ☐ Do NOT split tubes between pets

3 Month Application Tracker

First Application
DATE: ___ mm / ___ dd / ___ yy

Second Application
DATE: ___ mm / ___ dd / ___ yy

Third Application
DATE: ___ mm / ___ dd / ___ yy

For year-round protection, apply PRONYL OTC™ PLUS monthly.

PRONYL OTC™ PLUS for cats

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.

CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRONYL OTC™ PLUS for cats

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON CATS AND KITTENS OVER 8 WEEKS OF AGE.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use only. Do not use on kittens under 8 weeks of age. Individual sensitivities, while rare, may occur after using any pesticide product. Some cats may experience temporary startle effects when any product is applied. Cats may experience some temporary irritation at the site of product application. If signs persist, or become more severe within a few days of application, consult a veterinarian immediately. Certain medications can interact with pesticides. Consult a veterinarian before using on medicated, debilitated, or aged cats. Call 1-800-781-4738 for 24-hour assistance.

FIRST AID	
If swallowed:	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	Flush eyes with plenty of water. Call a doctor if irritation persists.
If on skin:	Wash with plenty of soap and water. Get medical attention if irritation persists.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-781-4738 for emergency medical treatment information.	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR CAT, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. FOR EXTERNAL USE ON CATS ONLY. DO NOT USE ON RABBITS. DO NOT USE ON OTHER ANIMALS.

Do not use on kittens under 8 weeks of age. Do not split one tube between cats. Do not use multiple tubes on one cat. Consult a veterinarian before using this product on debilitated, aged, or medicated cats. Can be used on breeding, pregnant, or lactating cats. Consult a veterinarian before using on cats with known organ dysfunction. Cats may experience some temporary irritation at the site of product application. Certain medications can interact with pesticides. Sensitivities may occur after using ANY pesticide product on cats. If signs of sensitivity occur bathe your cat with a mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

How to open: Remove product tubes from the package. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

How to apply: While holding the cat with one hand, use the other hand to apply the solution. Invert the tube and use the narrow end to part the cat's hair. Apply contents of one tube of PRONYL OTC™ PLUS as a spot between the shoulder blades. Hold the cat for a few seconds to give the solution time to be absorbed into the cat's coat. When finished, wrap the tube in paper and put into the trash. Only one applicator per treatment is needed.

FREQUENCY OF APPLICATION

Use PRONYL OTC™ PLUS monthly for complete control of flea, tick, and chewing lice infestations. Studies show that PRONYL OTC™ PLUS for cats kills adult fleas for up to six weeks. Apply monthly if your cat is at high risk for flea reinfestation. Apply monthly to control ticks, and chewing lice. PRONYL OTC™ PLUS remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.

In Case of Emergency, call: 1-800-781-4738
Non-Emergency, call: 1-800-224-PETS (7387)

Made in USA

If you have questions or comments about this product, please write: Sergeant's Consumer Response, P.O. Box 540399, Omaha, NE 68154-0399.

Distributed by: Sergeant's Pet Care Products, Inc.
P.O. Box 540399, Omaha, NE 68154-0399
EPA Reg. No. 2517-135
EPA Est. No. 2517-TN-1

P003064 AIRR

EPA Reg. No. 2517-135

0 73091 02969 4

PRONYL OTC™ PLUS for cats

For use ONLY on cats of ALL weights over 8 weeks of age

PRONYL OTC™ PLUS for cats

For use ONLY on cats of ALL weights over 8 weeks of age

3 MONTH SUPPLY

Sergeant's

PRONYL OTC™ PLUS for cats

Contains Fipronil and (S)-methoprene, the active ingredients used in Frontline Plus® for cats*

KILLS

- fleas
- ticks
- chewing lice
- flea eggs
- flea larvae

Controls flea reinfestation for up to 6 weeks
Waterproof

* PRONYL OTC™ PLUS for cats is not manufactured by or distributed by Merial. Frontline Plus® is a registered trademark of Merial.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Precautionary Statements

Net Contents: Three 0.017 fl oz (0.50 mL) applicators

ACTIVE INGREDIENTS:

Fipronil	9.80%
(S)-methoprene	11.80%
OTHER INGREDIENT:**	**78.40%**
TOTAL	**100.00%**

**diethylene glycol monoethyl ether

LIFT HERE FOR MORE INFORMATION

PRONYL OTC™ PLUS for cats

Flea Life Cycle



- Kills adult fleas, flea eggs, flea larvae, and prevents the development of flea pupae for up to 6 weeks
- Kills deer ticks for up to 4 weeks
- Kills ticks including those that may transmit Lyme disease
- Kills brown dog ticks, American dog ticks, and lone star ticks
- Rapid elimination of chewing lice infestation
- Can be used on breeding, pregnant, and lactating cats
- Fast acting
- Long lasting

The active ingredients in PRONYL OTC™ PLUS feature a sustained release formula developed for fast acting and long lasting control of fleas, ticks, and chewing lice for 4 weeks on cats.