

Directions for Use



How to Open:

1. Remove product tubes from the package.
2. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.



How to Apply:

1. Invert tube over cat and use open end to part cat's hair.
2. Apply contents of one tube of PRONYL OTC™ MAX for cats as a stripe, starting high on the back of the cat's neck to in front of the shoulder blades. Squeeze tube firmly to apply all of the solution to the cat's skin.

Monthly Application Reminder

First Treatment

DATE: mm / dd / yy

Second Treatment

DATE: mm / dd / yy

Third Treatment

DATE: mm / dd / yy

Apply PRONYL OTC™ MAX monthly for continuous protection against fleas, ticks and chewing lice.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.
CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRONYL OTC™ MAX for cats

PRONYL OTC™ MAX for cats



READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON CATS AND KITTENS OVER 12 WEEKS OF AGE.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Harmful if swallowed. Harmful if absorbed through skin.

HAZARDS TO DOMESTIC ANIMALS. For external use only. Do not use on kittens under 12 weeks of age. Individual sensitivities, while rare, may occur after using any pesticide product. If signs persist, or become more severe within a few days of application, consult a veterinarian immediately. If your pet has an unusual reaction to the initial application, consult a veterinarian before repeating treatment. Certain medications can interact with pesticides. Consult a veterinarian before using on medicated, debilitated or aged animals. Call 1-800-224-PETS (7387) for 24-hour assistance.

FIRST AID

If In Eyes	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with water for 15-20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. •Call a poison control center or doctor for further treatment advice.
If On Skin Or Clothing	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes. •Call a poison control center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none"> •Call a poison control center or doctor immediately for treatment advice. •Have person sip a glass of water if able to swallow. •Do not induce vomiting unless told to do so by a poison control center or doctor. •Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-781-4738.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE EACH USE. FOR EXTERNAL USE ON CATS ONLY. Do not use on kittens under 12 weeks of age. Do not treat your cat with more than one pesticide product at a time. Over dosing your cat can result in serious illness and even death. **Best if used year round.** Consult a veterinarian before using this product on debilitated, aged, medicated cats. Consult a veterinarian before using on cats with known organ dysfunction. Separate the treated cat from all other cats and dogs for 24 hours after treatment has been applied. Cats and dogs that actively groom or engage in close physical contact with treated cat may be at risk of serious harmful effects. The most common signs of ingestion are excessive salivation and foaming at the mouth. If these symptoms occur immediately feed your cat and continue to monitor your pet for the next 24 hours.

Sensitivities may occur after using ANY pesticide product on cats. If signs of sensitivity occur bathe your cat with a mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately by calling 1-800-781-4738.

How to open: Remove product tubes from the package. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

How to apply: While holding the cat with one hand, use the other hand to apply the solution. Invert the tube and use the narrow end to part the cat's hair. Apply contents of one (0.017 fl oz) (0.50 ml) tube of PRONYL OTC™ MAX for Cats as a stripe, starting high on the back of the cat's neck to in front of the shoulder blades. Hold the cat for a few seconds to give the solution time to be absorbed into the cat's coat. When finished, wrap the tube in paper and put into the trash. Only one applicator per treatment is needed.

FREQUENCY OF APPLICATION

Use PRONYL OTC™ MAX for Cats monthly for complete control of flea, tick and chewing lice infestations. PRONYL OTC™ MAX for Cats kills fleas for up to six weeks. Apply monthly, if your pet is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. PRONYL OTC™ MAX for Cats remains effective, even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.

If you have questions or comments about this product, please write: Sergeant's Consumer Response, P.O. Box 540399, Omaha, NE 68154-0399.

Distributed by: Sergeant's Pet Care Products, Inc. P.O. Box 540399, Omaha, NE 68154-0399
 EPA Reg. No. 2517-148
 EPA Est. No. 2517-TN-1*, EPA Est. No. 2517-NE-02*
 Superscript is the first letter in the lot code

In Case of Emergency, call: 1-800-781-4738
 Non-Emergency, call: 1-800-224-PETS (7387)

P004098 RARR

Patents Pending

EPA Reg. No. 2517-148



PK2792



DISC.



LICENSED

9227.325

LICENSED

Product 2012-2014 Lic. No.

Product 2015-2017 Lic. No.


PRONYL OTC™ MAX for cats

3 MONTH SUPPLY

For use ONLY on cats & kittens of ALL weights over 12 weeks of age

Sergeant's
PRONYL OTC™ MAX for cats

Starts **KILLING FLEAS & TICKS** in as little as **ONE HOUR!**



Kills fleas

Kills ticks

Kills chewing lice

Waterproof

Prevents and controls reinfestations

KEEP OUT OF REACH OF CHILDREN CAUTION
 See Back Panel for Additional Precautionary Statements
 Net Contents: Three 0.017 fl oz (0.50 mL) applicators

ACTIVE INGREDIENTS:
 Fipronil 9.80%
 Etofenprox 15.00%
OTHER INGREDIENT*: 75.20%
TOTAL 100.00%
 *Diethylene glycol monoethyl ether

LIFT HERE FOR MORE INFORMATION

- ▶ Starts to kill fleas and ticks in an hour, continues to kill for a month
- ▶ Prevents and controls reinfestations
- ▶ Controls flea reinfestation for up to six weeks
- ▶ Kills fleas that may cause flea allergy dermatitis, flea bite anemia, and tapeworm infestation
- ▶ Kills Deer Ticks (which may transmit Lyme disease) for up to 4 weeks
- ▶ Kills brown, American and lone star dog ticks for up to 4 weeks
- ▶ Fast action against chewing lice infestations
- ▶ Fast acting, long lasting

Sergeant's® PRONYL OTC™ MAX for cats contains Fipronil and Etofenprox to effectively kill fleas, ticks and chewing lice. The active ingredients in Sergeant's® PRONYL OTC™ MAX for Cats feature a sustained-release formula developed for fast-acting and long-lasting control of fleas, ticks and chewing lice for 30 days on cats.

6 Step Topical Checklist

- Read the label completely and follow directions
- Weigh your pet
- Do NOT use dog products on cats
- Do NOT treat with more than one pesticide product at a time
- Separate animals after treatment to avoid chance of ingestion
- Do NOT split tubes between pets

PROJECT NAME & PROJECT NUMBER	DATE	DRAFT#	DIMENSIONS	UPC CODE	PART#
Pronyl OTC MAX	17OCT11	9		UPC 0 73091 02249 7	P004098