

GRANT'S KILLS ANTS ANT & SPIDER KILLER Granules



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

9230.20

VALUE SIZE!

25% more than
14 oz.
size



DISC. GRANT'S KILLS ANTS®

ANT & SPIDER KILLER

Granules

KILLS ANTS AND SPIDERS



ANTS

SPIDERS

KILLS OTHER LISTED INSECTS



CHINCH BUGS



FLEAS



SILVERFISH



COCKROACHES



CRICKETS



SOWBUGS



GRASSHOPPERS



TICKS



SCORPIONS

ACTIVE INGREDIENT:

*Permethrin..... 0.25%

INERT INGREDIENTS:..... 99.75%

TOTAL 100.00%

*Cis/trans ratio: Maximum 55% (±) cis and minimum 45% (±) trans

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for Additional
Precautionary Statements.

NET WEIGHT 17.5 OZ. (496 grams)

FIRST AID	
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Avoid contamination of food or feed. Utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticides.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water. Drifts and runoffs from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is recommended for the control of lawn insects, fleas, and ticks outdoors in residential areas. Repeated applications are made by broadcast treatment with a spreader to lawns, under shrubbery, and other ornamental and turf grass areas, band treatments are recommended around house foundation and spot applications are made to ant mounds and wasp nests.

STORAGE AND DISPOSAL

STORAGE: Store product in its original container in a cool, dry locked place out of reach of children.

DISPOSAL: If empty – Do not reuse this container. Place in trash or offer for recycling if available. If partly filled – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL USE DIRECTIONS

- Determine size of area to be treated – measure and multiply length times width.
- Treat as directed – thorough coverage is important. Water in where indicated.

Keep children and pets off treated areas until this material is washed into turf and grass is dry.

Treatment Areas	Pests	Use Rate	How to Use
Overall Lawn Treatment For Grass and Dichondra Lawns	Ants, Armyworms (Inc. Fall, Yellow-striped, etc.), Bermudagrass mites, Chiggers, Chinch Bugs, Clover Mites, Cutworms, Darkling Ground Beetles, Digger wasps, European Crane Fly Larvae, Flea Beetle (In dichondra), Grasshopper, Hyperodes (Turfgrass Weevil), Lawn Moths, Leafhoppers, Silverfish, Vegetable Weevils (In dichondra)	Treat when insects first appear using granules at the rate of 17.5 oz. per 547 sq. ft.	Apply this product uniformly to home lawns and under shrubbery. Immediately after application, water the lawn lightly. European Crane fly larvae (Leather-jackets) – the best time to apply is in May or June when the insects are young. Chinch Bugs – Start treatments when temperatures reach 80°F or above. A second application may be necessary in 6 weeks if insect activity continues.

(continued)

Treatment Areas	Pests	Use Rate	How to Use
Spot Treatment	Ants nesting outdoors including: Acrobat, Argentine, Big Headed, Cornfield, Field, Harvester, Leafcutter, Little Black, Pavement and Pyramid Ants	Product should be applied to thoroughly cover the nest (1 to 2 teaspoons) and sprinkled gently with water.	If possible, it is best to try to locate nest and treat this first. Nests will usually be found in the ground but may be concealed under stones, boards, slabs or in stumps or logs. In addition to treating ant nests or mounds, entry points of buildings (used by ants) and the soil surface along the perimeter of the building should be treated.
	Digger Wasps	Apply 1 to 2 teaspoons of granules in and around each individual ground nest opening.	Make the application at dusk when the wasps are not active.
	Imported Fire Ants	Apply one cup over, and 2 feet around each fire ant mound.	Immediately after application, water the granules into the soil gently with a light sprinkling to avoid disturbing the colony. Use at least 1 gallon of water per mound, but no more than 2 gallons. Use of high pressure watering equipment may disturb the colony and cause migration of the ants, thus reducing the effectiveness of the treatment. Apply in cool weather (65-80°F), or in early morning or late evening for best results. Treat new mounds as they appear.
Band (Perimeter) Treatment Around House Foundation	Ants, Boxelder Bugs, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Field Crickets, Fleas, Scorpions, Sowbugs, Spiders (Inc. Black Widow), Springtails (Collembola) and Ticks	Treat at a rate of 17.5 oz. per 547 sq. ft.	To help prevent infestation of buildings, apply a 5 foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly after application.

FLEA AND TICK CONTROL (OUTDOORS)

In summer months, fleas and ticks frequently become established in lawns and pet resting areas surrounding the home. For best control of Fleas and Ticks (including but not limited to Bear Tick, Black-legged Tick, Brown Dog Tick, Deer Tick (Ixodes dammini), Lone Star Tick, Western Black-legged Tick, Wood Tick): a regular treatment program should be followed. Treat once a month when pests first appear in late spring (May-June) and continue until late summer (September-October). Fall temperatures below 50°F decrease pest activity. Eliminating weeds around the home and mowing lawns frequently reduces the hiding places for these pests. Note: Dogs, cats, their kennel or bedding area should also be treated with a registered product. In addition, if fleas and ticks are indoors on carpets or pet resting areas, use a registered product. It is very important that all three sites – outdoors, indoors and also the pet be treated with appropriate labeled products to achieve satisfactory control. If any of these areas is not treated, it could serve as a source of reinfestation to the entire premises.

Warranty and Disclaimer: This product is guaranteed to the extent of the purchase price only when used according to label directions.

Manufactured for
GRANT LABORATORIES, INC.
San Leandro, Ca 94578
Form 228-02
EPA Reg. No. 28293-328-1663
EPA Est. No. 59144-CA-1, 62478-FL-1a,
44616-MO-1, 67572-GA-1
Subscript used is first letter of
run code on container

