

Hartz <sup>7/8"</sup>  
InControl™  
Flea & Tick Collar for Dogs

[0.875"]

Hartz <sup>5/8"</sup>  
InControl™  
[0.625"]



Front  
Panel

Flea & Tick Collar  
for Dogs

**KILLS:**  
FLEAS • TICKS  
FLEA EGGS

7 Months Protection

Water Resistant



**ACTIVE INGREDIENTS:**  
Tetrachlorvinphos ..... 14.55%  
(S)-Methoprene ..... 1.02%  
**OTHER INGREDIENTS:** ..... 84.43%  
**TOTAL** ..... 100.0%

**CAUTION: DO NOT LET CHILDREN  
PLAY WITH THIS COLLAR.**  
SEE BACK PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS.

**NET CONTENTS**  
1 COLLAR  
**NET WT. 1.16 oz. (33 g)**

Hartz <sup>7/8"</sup>  
InControl™  
Flea & Tick Collar for Dogs

[0.875"]

[0.625"]



STATE OF HAWAII  
Department of Agriculture  
**ACCEPTED**  
LICENSE NO.

9235.371

**READ ENTIRE LABEL  
BEFORE EACH USE.**  
**USE ONLY ON DOGS.**  
**PRECAUTIONARY  
STATEMENTS**  
**HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS**

**CAUTION:**  
Harmful if swallowed or  
absorbed through skin. Causes  
moderate eye irritation. Avoid  
contact with eyes, skin, or  
clothing. Wash thoroughly with  
soap and water after handling.

**FIRST AID:**  
Contains an organophosphate that inhibits cholinesterase.  
**IF SWALLOWED:**  
Call a Poison Control Center or doctor immediately for treatment  
advice. Have person sip a glass of water if able to swallow.  
Do not induce vomiting unless told to do so by a Poison Control  
Center or doctor. Do not give anything by mouth to an  
unconscious person.

**IF ON SKIN OR CLOTHING:**  
Take off contaminated clothing. Rinse skin immediately with  
plenty of water for 15-20 minutes. Call a Poison Control Center  
or doctor for treatment advice.

**IF IN EYES:**  
Hold eye open and rinse slowly and gently with water for  
15-20 minutes. Remove contact lenses, if present, after the  
first 5 minutes, then continue rinsing eye. Call a Poison Control  
Center or doctor for treatment advice.

Have the product container or label with you when calling a  
Poison Control Center or doctor, or going for treatment.  
**Note to physicians and veterinarians:**  
This product contains an organophosphate and may cause  
cholinesterase inhibition. Cholinergic symptoms may include  
salivation, miosis, incoordination, muscle fasciculation and/or  
weakness, vomiting, and diarrhea. Atropine is antidotal. 2-PAM  
may be effective as an adjunct to atropine. Call 1-800-275-1414  
or your local Poison Control Center for further information.

For more information, call our experts at 1-800-275-1414.

Made by The Hartz Mountain Corporation,  
400 Plaza Drive, Secaucus, NJ 07094  
Hartz® and other trademarks are trademarks of  
The Hartz Mountain Corporation.

EPA Reg. No. 2596-139 • EPA Est. No. 2596-NJ-1



**DIRECTIONS FOR USE:**  
It is a violation of Federal Law  
to use this product in a manner  
inconsistent with its labeling.

REMOVE COLLAR FROM  
PACKAGE, UNROLL AND  
STRETCH TO ACTIVATE  
INSECTICIDE GENERATOR.  
AS THE COLLAR BEGINS  
TO WORK, A FINE WHITE  
POWDER WILL APPEAR  
ON THE SURFACE.

Do not use on puppies under 6 weeks old. Do not unroll  
collar until ready to use. Place the Hartz® InControl™  
Flea & Tick Collar for Dogs around the dog's neck,  
adjust for proper fit, and buckle in place. The collar must  
be worn loosely to allow for growth of the animal and to  
permit the collar to move around the neck. A properly fitted  
collar is one that, when fastened, will snugly slide over the  
pet's head. Leave 2 or 3 inches on the collar for extra  
adjustment and cut off and dispose of the extra length.  
Consult a veterinarian before using this product on  
dehydrated, aged or medicated animals. If used on  
pregnant or nursing dogs, do not replace collar until the  
puppies are at least 6 weeks old. Sensitivity may occur  
after using ANY pesticide product for pets. Some animals  
may become irritated by any collar if it is applied too  
tightly. If this occurs, loosen the collar. If irritation  
continues, remove the collar from the animal. If signs of  
sensitivity occur, bathe your pet with mild soap and rinse  
with large amounts of water. If signs continue, consult a  
veterinarian immediately. Do not use this product on  
animals simultaneously or within 30 days before or after  
treatment with or exposure to cholinesterase inhibiting  
drugs, pesticides, or similar chemicals. However, flea and  
tick collars may be immediately replaced. This collar is  
intended for use only as an insecticide/IGR generator. The  
collar will begin to kill fleas, ticks and flea eggs and larvae  
immediately. Its full protection against harder-to-kill ticks  
will be built up within a few days after being placed on the  
pet. For continuous protection  
under normal conditions replace the collar every 7 months.  
Under conditions where pets are exposed to severe flea  
and tick infestations, it may be necessary to replace the  
collar more frequently. Wetting will not impair the collar's  
effectiveness or the pet's protection. If the pet goes  
swimming or is out in the rain it is not necessary to remove  
the collar. The Hartz® InControl™ Flea & Tick Collar for  
Dogs may be worn with a regular collar.

**STORAGE & DISPOSAL:**  
Storage: Store at room temperature. Disposal: Do not reuse  
empty pouch. Place in trash or offer for recycling if available.

RM107351

7/8"  
[0.875"]

Hartz  
InControl™  
Flea & Tick Collar for Dogs

InControl™  
Flea & Tick Collar for Dogs

Hartz

7/8"

Job No:

[0.875"]  
Do not use PDF files for printing

27/32"