

1/4"
NO
COPY

6-5/16"

6-15/16"

1/4" GLUE LAP
NO VARNISH
1/8"
NO COPY



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

9235.410



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2008-2010 LIC. NO.

TYPE
5-3/8"

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE USE ONLY ON CATS

1. To kill fleas, ticks and lice, and to reduce itching and scratching due to insect bites, dust entire cat beginning at head and working back.
2. Make sure that the powder gets down to the skin.
3. Take care to treat feet and legs.

Recommended Dosage: Use approximately 1/3 ounce of powder for a small cat. Use approximately 1/2 ounce of powder for a large cat. There is sufficient powder in this container to treat a large cat 8 times. Lightly dust cat's bedding with approximately the same amount of powder. Repeat treatment of cat and bedding at weekly intervals if necessary.


STORAGE AND DISPOSAL:

Do not contaminate water, food or feed. Store in a cool, dry place. **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

ACTIVE INGREDIENT:

Tetrachlorvinphos	3.3%
OTHER INGREDIENTS	96.7%
TOTAL	100.0%

EPA Reg. No. 2596-78
EPA Est. No. 51435-WI-004



Hartz® and other trademarks are owned by The Hartz Mountain Corporation.

Made for ©The Hartz Mountain Corporation, 400 Plaza Drive, Secaucus, N.J. 07094 USA

504144

KEEP OUT OF REACH OF CHILDREN

CAUTION:

SEE SIDE PANEL FOR INGREDIENTS AND ADDITIONAL PRECAUTIONS.

NET WT:
4 oz (113g)

Hartz®

ULTRA GUARD™

Flea & Tick Powder for Cats



Kills:



Fleas



Ticks



Lice

for up to 7 days

Water Resistant

Hartz® UltraGuard™ Flea & Tick Powder for Cats
When your cat has fleas and ticks, your family and pet may be exposed to:

- Lyme disease
- Tapeworm infestation
- Flea bite anemia
- Flea allergy dermatitis

To learn more, visit www.hartz.com

For more information, call our flea and tick experts at 1-800-275-1414.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Not for use on kittens less than 12 weeks of age. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing, or medicated animals. Sensitivity may occur after using ANY pesticide product on pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. However, flea and tick collars may be immediately replaced.

FIRST AID – Contains an organophosphate that inhibits cholinesterase.

If swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

Note to Physicians and Veterinarians: This product contains an organophosphate and may cause cholinesterase inhibition. Cholinergic symptoms may include salivation, miosis, incoordination, muscle fasciculation and/or weakness, vomiting and diarrhea. Atropine is antidotal only if symptoms of cholinesterase inhibition are present. 2-PAM may be effective as an adjunct to atropine. Call 1-800-275-1414 or your local Poison Control Center for further information.